

< Back to results | 1 of 1

CSV exportDownloadPrintE-mailSave to PDFSave to listMore...>

Journal of Physics: Conference Series • Open Access • Volume 1306, Issue 1 • 9 September 2019 • Article number 012051 • 2nd International Conference on Mathematics: Education, Theory, and Application, ICMETA 2018 • Sukoharjo • 30 October 2018 through 31 October 2018 • Code 151630

Document type
Conference Paper • Bronze Open Access • Green Open Access

Source type
Conference Proceedings

ISSN
17426588

DOI
10.1088/1742-6596/1306/1/012051

View more

Optimizing Range Norm of the Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components

Siswanto^a✉, Suparwanto A.^b, Rudhito M.A.^c
Save all to author list

^a Department of Mathematics, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta, 57126, Indonesia
^b Department of Mathematics, Faculty of Mathematics and Natural Sciences, Gadjah Mada University, Yogyakarta, 55281, Indonesia
^c Program Study Mathematics in Education, Faculty of Teacher Training and Education, Sanata Dharma University, Yogyakarta, 55282, Indonesia

14

Views count

View all metrics

View PDFFull text options

Abstract

Indexed keywords

SciVal Topics

Metrics

Abstract

L_ε be the set of all real numbers and $\mathbb{R}_\epsilon = \cup \{ \epsilon \}$ whose $\epsilon = -\infty$. Max-plus algebra is the set \mathbb{R}_ϵ that is equipped two operations maximum and addition. It can be formed matrices in the size of m x n whose elements belong to \mathbb{R}_ϵ , called matrix over max-plus algebra. Optimizing range norm of the image set of matrix over max-plus algebra with prescribed components has been discussed. Interval Max-Plus Algebra is the set with $\epsilon = [\epsilon, \epsilon]$, is equipped with two operations maximum and addition. The set of all matrices in the size of m x n whose elements belong to $I(\mathbb{R}_\epsilon)$, called matrix over interval max-plus algebra. Optimizing range norm of the image set of matrix over interval max-plus algebra has been discussed. In this paper, we will discuss optimizing range norm of the image set of matrix over interval max-plus algebra with prescribed components. © 2019 IOP Publishing Ltd. All rights reserved.

Indexed keywords	>
SciVal Topics	>
Metrics	>

References (12)

View in search results format >

☐ All

CSV exportDownloadPrintE-mailSave to PDFCreate bibliography

1

Bacelli, F., Cohen, G., Older, G.J., Quadrat, J.P. (2001) *Synchronization and Linearity, An Algebra for Discrete Events System*. Cited 1211 times. (John Wiley and Sons)

2

Farlow, K.G. (2009) Max-Plus Algebra Master's Thesis submitted to the Faculty of the Virginia Polytechnic Institute and State University in partial fulfillment of the requirements for the degree of Masters in Mathematics

- ☐ 3 Tam, K.P.
(2010) *Optimizing and Approximating Eigenvectors in Max Algebra*. Cited 13 times.
A Thesis Submitted to The University of Birmingham for The Degree of Doctor of Philosophy (Ph.D)
-
- ☐ 4 Akian, M., Cohen, G., Gaubert, S., Quadrat, J.P., Viot, M.
(1994) *Max-Plus Algebra and Applications to System Theory and Optimal Control*
(Zurich, Switzerland: Proceedings of the International Congress of Mathematicians)
-
- ☐ 5 Rudhito, M.A.
Fuzzy NumberMax-Plus Algebra and Its Application to Fuzzy Scheduling and Queuing Network Problems. Cited 2 times.
(Indonesian: The Study Program S3 Mathematics Faculty of Mathematics and Natural Sciences : Gadjah Mada University, 2011)
-
- ☐ 6 Butkovic, P., Tam, K.P.
On Some Properties of the Image of a Max Linear Mapping
(2009) *Contemporary Mathematics*, 495, pp. 115-126. Cited 16 times.
-
- ☐ 7 Siswanto, Suparwanto, A., Rudhito, M.A.
(2013) *Minimizing Range Norm of the Image Set of Matrix over Interval Max-Plus Algebra*
(Indonesian: Proceedings of The National Seminar on Mathematics and Applications, Department of Mathematics, Faculty of Science and Technology, Airlangga University) 978-602-14413-0-5
-
- ☐ 8 Siswanto, Suparwanto, A., Rudhito, M.A.
Eigenvector Space of a Matrix of Interval Max-Plus Algebra
(2014) *Jurnal Matematika & Sains (JMS)*. FMIPA ITB, 19, pp. 8-15. Cited 5 times.
Indonesian
-
- ☐ 9 Siswanto, Suparwanto, A., Rudhito, M.A.
(2014) *Existence of Solution to the System of Linear Equations in Interval Max-Plus Algebra*
(Indonesian: Proceedings of the National Conference of Mathematics XVII, Institute of Technology Surabaya) 978-602-96426-3-6
-
- ☐ 10 Siswanto, Suparwanto, A., Rudhito, M.A.
Optimizing range norm of the image set of matrix over interval max-plus algebra ([Open Access](#))

(2016) *Far East Journal of Mathematical Sciences*, 99 (1), pp. 17-32. Cited 2 times.
http://www.pphmj.com/article.php?act=art_download&art_id=9502
doi: 10.17654/JMS099010017

[View at Publisher](#)
-
- ☐ 11 Kreinovich, V.
Why Intervals? Why Fuzzy Numbers? Towards a New Justification
(2007) *International Journal of Information Technology and Intelligent Computing*, 2.
-
- ☐ 12 Konigsberg, Z.R.
A Generalized Eigenmode Algorithm for Reducible Regular Matrices over the Max-Plus Algebra
(2009) *International Mathematical Forum*, 4, pp. 1157-1171. Cited 2 times.
-

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Tutorials
- Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.



also developed by scimago

SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

Home

Journal Rankings

Country Rankings

Viz Tools

Help

About Us

Journal of Physics: Conference Series

<div>COUNTRY</div> <div>United Kingdom</div> <div>Universities and research institutions in United Kingdom</div>	<div>SUBJECT AREA AND CATEGORY</div> <div>Physics and Astronomy</div> <div>Physics and Astronomy (miscellaneous)</div>	<div>PUBLISHER</div> <div>IOP Publishing Ltd.</div>	<div>H-INDEX</div> <div>85</div>
<div>PUBLICATION TYPE</div> <div>Conferences and Proceedings</div>	<div>ISSN</div> <div>17426588, 17426596</div>	<div>COVERAGE</div> <div>2005-2020</div>	<div>INFORMATION</div> <div>Homepage</div> <div>How to publish in this journal</div> <div>jpcs@iopublishing.org</div>

SCOPE

The open access Journal of Physics: Conference Series (JPCS) provides a fast, versatile and cost-effective proceedings publication service.

Join the conversation about this journal

Quartiles

SJR

2006 2008 2010 2012 2014 2016 2018 2020

Total Documents

2005 2007 2009 2011 2013 2015 2017 2019

Citations per document

2005 2007 2009 2011 2013 2015 2017 2019

Total Cites

Self-Cites

2005 2007 2009 2011 2013 2015 2017 2019

External Cites per Doc

Cites per Doc

2005 2007 2009 2011 2013 2015 2017 2019

% International Collaboration

2005 2007 2009 2011 2013 2015 2017 2019

Citable documents

Non-citable documents

2005 2007 2009 2011 2013 2015 2017 2019

Cited documents

Uncited documents

2005 2007 2009 2011 2013 2015 2017 2019

Journal of Physics: Conference Series

Q4

Physics and Astronomy (miscellaneous)

best quartile

SJR 2020

0.21

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code:

SCImago Graphica

Explore, visually communicate and make sense of data with our new free tool.

Get it

https://www.scimagojr.com/journalsearch.php?q=130053&tip=sid&clean=0

1/14

Journal of Physics

Conference Series

The 11th Biennial Conference on
Classical and Quantum Relativistic
Dynamics of Particles and Fields

1239

VOLUME 1239 – 2019

4–7 June 2019
Mérida, Yucatán, Mexico

EDITOR
Marta Lind

The open access journal for conference proceedings
iopscience.org/jpcs

IOP Publishing

The Committee of 6th South East Asia Design Research International Conference (SEA-DR IC) 2018**Steering Committee**

Prof. Dr. Ir. Samsul Rizal, M. Eng (Rector of Syiah Kuala University)

Prof. Dr. Djufri, M. Si (Dean of FKIP Syiah Kuala University)

Prof. Dr. Maarten Dolk

Prof. Dr. R.K Sembiring

Dr. Y. Marpaung

Prof. Dr. Zulkardi, M. I. Kom, M.Sc.

Prof. Dr. Sutarto Hadi, M.Si., M. Sc.

Prof. Dr. Dian Armanto, M.Pd, M.Sc.

Prof. Dr. Ahmad Fauzan, M.Pd, M. Sc.

Prof. Dr. Turmudi, M.Ed, M.Sc.

Dr. M. Ikhsan, M.Pd. (Head of Mathematics Education Department of Syiah Kuala Univesity)

Prof. Berinderjeet Kaur

Prof. Lilia Halim

Organizing Committee

Conference Chair : Dr. Rahmah Johar, M.Pd.

Conference Co-Chair : Dr. Anwar, M.Pd.

Secretary : Dr. Mailizar, M.Ed.

Vice Secretary : Elizar, Ph.D.

Treasurer : Dr. Cut Morina Zubainur, M.Pd.

Vice Treasurer : Suhartati, M.Pd.

Scientific Committee

Scientific Committee Chair : Prof. Dr. Marwan, M. Si

Scientific Committee Member :

Prof. Maarten Dolk

Prof. Rohaida Mohd. Saat

Prof. Dr. Musri Musman, M.Sc.

Prof. Dr. Ratu Ilma Indra Putri, M.Si.

Prof. Dr. Muchlisin Z.A, M.Sc.

Prof. Dr. Adlim, M.Sc.

Prof. Dr. Ahmad Fauzan, M.Sc.

Dr. Cathy Wissehr

Dr. Wanty Widjaja

Dr. Abdul Halim Abdullah

Dr. Ariyadi Wijaya, M.Sc.

Dr. Al Jupri, M.Sc.

Dr. Tatag Yuli Eko Siswono, M.Pd.

Dr. Yenita Roza, M.Sc.

Dr. Shintia Revina, M.Sc.

Dr. Neni Mariana, M.Sc.

Dr. Rooselina Ekawati, M.Sc.

Dr. Nasrullah Idris, M.Eng.

Dr. Hongki Julie, M.Pd.

Dr. Taufik Fuadi Abidin, M.Tech.

Dr. Rully Charitas Indra Prahmana

Dr. Supriatno, M.Si.
Dr. Said Munzir, M.Eng.Sc.
Dr. Suhartono, M.Sc.
Aysenur Alp
Zarlaida Fitri, M.Sc.
Veronika Fitri Rianasari, M.Sc.
Meliasari., M.Sc.
Zetra Hainul Putra, M.Sc.
Destina Wahyu Winarti, M.Sc.
Fridgo Tasman, M.Sc.
Fatimatul Khikmiyah, M.Sc.
Achmad Badrun Kurnia, M.Sc.
Mulia Putra, M.Pd., M.Ed.
Rita Novita, M.Pd.
Bustang Buhari, M.Sc.
Intan Kemala Sari, M.Pd.

Keynote Speakers

Prof. Berinderjeet Kaur	National Institute of Education, Singapore
Prof. Maarten Dolk	Utrecht University, the Netherlands
Prof. Lilia Halim	University Kebangsaan Malaysia, Malaysia
Dr. John Willison	the University of Adelaide, Australia

The 6th South East Asia Design Research International Conference (SEA-DR IC) 2018

**R Johar¹, C Morina¹, Anwar¹, Mailizar¹, Elizar¹, C Khairunnisak¹, R C I Prahmana²,
W Artika¹, L Vitoria¹, L Khairi¹, S Maulina¹ and M Ulfa¹**

¹Syiah Kuala University, Banda Aceh, 23111, Indonesia

²Universitas Ahmad Dahlan, Jl. Pramuka Kav. 5, Yogyakarta 55161, Indonesia

E-mail: rahmahjohar@fkip.unsyiah.ac.id

Preface

The South East Asia Design Research (SEA-DR) as a forum of design research in collaboration with Master Program of Mathematics Education, Syiah Kuala University, organized the 6th SEA-DR conference with the theme “Inspiring students to learn: Fostering innovative teaching and learning of science, mathematics and technology”.

This conference was an excellent opportunity for academics, researchers, teachers and students to share knowledge, experiences and research findings as well as to inspire the best practice of development research in the field of teaching mathematics, science, and technology.

We had four keynote speakers that were Prof. Berinderjeet Kaur, Prof. Maarten Dolk, Prof. Lilia Halim, and Dr. John Willison. We also had ten invited speakers and four keynote speakers in workshop sessions. Furthermore, there were 181 papers, including 148 oral presentation and 33 posters presentations. The 6th SEA-DR conference successfully attracted delegates from many countries. There were seven countries participating in this conference, including: Singapore, the Netherlands, Denmark, Australia, Malaysia, Brunei Darussalam, and Indonesia.

Finally, we would like to extend our gratitude for everyone involved for their contribution in the conference.



NOTICE: Ukraine: Click here to read IOP Publishing's statement.

Table of contents


Volume 1306
2019

◀ Previous issue Next issue ▶

The 2nd International Conference on Mathematics: Education, Theory, and Application 30-31 October 2018, Sukoharjo, Indonesia
Accepted papers received: 11 July 2019
Published online: 09 September 2019



Open all abstracts



Preface



OPEN ACCESS Preface	011001
✚ Open abstract  View article  PDF	


OPEN ACCESS Peer review statement	011002
✚ Open abstract  View article  PDF	



Papers



OPEN ACCESS The Relation between Almost Noetherian Module, Almost Finitely Generated Module and T -Noetherian Module A Faisol, B Surodjo and S Wahyuni	012001
✚ Open abstract  View article  PDF	



OPEN ACCESS Maternal Antibody Susceptible Vaccinated Infected Recovered (MSVIR) Model for Tetanus Disease and Its Applications in Indonesia Purnami Widyarningsih, Primadita Candrawati, Sutanto and Dewi Retno Sari Saputro	012002
✚ Open abstract  View article  PDF	



OPEN ACCESS Petri Net and Max-Plus Algebra Model in Bank Queue with Two Servers in Each Service Gevinda Arulia Martha, Siswanto and Tri Atmojo Kusmayadi	012003
✚ Open abstract  View article  PDF	

OPEN ACCESS Students' Intuition Characteristics in Solve Mathematical Problem In Stage Planning Arwanto, I Ketut Budayasa and Mega Teguh Budiarto	012004
✚ Open abstract  View article  PDF	

OPEN ACCESS Characteristic Polynomial of a Triangular and Diagonal Strictly Double \mathbb{R} -astic Matrices over Interval Max-Plus Algebra Aisyah Yvni Wulandari and Siswanto	012005
✚ Open abstract  View article  PDF	

OPEN ACCESS Super (a, d) - H -antimagic total labeling of edge corona product on cycle with path graph and cycle with cycle graph Arum Permata Sari, Titin Sri Martini and Vika Yugi Kurniawan	012006
✚ Open abstract  View article  PDF	

OPEN ACCESS Super (a, d) - H -antimagic total labeling on wheel edge corona product with a path and a cycle Ufii Nadzima and Titin Sri Martini	012007
✚ Open abstract  View article  PDF	

OPEN ACCESS On H -supermagic labeling of friendship edge corona product with path and sun edge corona product with path T Noviati, T S Martini and D Indriati	012008
✚ Open abstract  View article  PDF	



OPEN ACCESS H^- Supermagic labeling on $domino \odot P_n$ and $C_n \diamond S_n$ D A Anggraeni, T S Martini and T A Kusmayadi	012009
✚ Open abstract  View article  PDF	

OPEN ACCESS On Super $(a, d) - H^-$ Antimagic Total Labeling of Edge Corona Product and Corona Product of Gear Graph with Path and Cycle R Nistyawati and T S Martini	012010
✚ Open abstract  View article  PDF	

OPEN ACCESS A New Approach in Decision of Pareto Efficient Concept in Probabilistic Vendor – Buyer Supply – Chain Problem for Imperfect Quality with Lead – Free Demand R Setiawan	012011
✚ Open abstract  View article  PDF	

OPEN ACCESS (a, d) - H -Anti Magic Total Labeling on Double Cones Graph Mania Roswitha, Titin Sri Martini and Surya Aji Nugroho	012012
✚ Open abstract  View article  PDF	

OPEN ACCESS Estimation Claims Reserving Based on Archimedean Copula Lienda Noviyanti, R Hasna Afifah, A. Zanbar Soleh and Anna Chadidjah	012013
✚ Open abstract  View article  PDF	

OPEN ACCESS The local metric dimension of edge corona and corona product of cycle graph and path graph Elis Dyah Wulancar and Tri Atmojo Kusmayadi	012014
✚ Open abstract  View article  PDF	

OPEN ACCESS The local metric dimension of generalized broken fan graph and edge corona product of star graph and path graph	012015
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.	

Silfiatul Khoiriah and Tri Atmojo Kusmayadi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012016

Students' Analytical Thinking Profile Based on Reflective Cognitive Style in Solving Mathematics Problem

A Qolthahiriyus, I Sujadi and D Indriati

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012017

On the metric dimension of generalized broken fan graph and edge corona product of path with star graph

Makarima Sabila and Tri Atmojo Kusmayadi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012018

On the strong metric dimension of crossed prism graph and edge corona of cycle with path graph

Astri Yunitasari and Tri Atmojo Kusmayadi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012019

Nonlinear system identification model of the spread of TB disease using the genetic algorithm and multilayer perceptron

A Damayanti, P Subiyanto and A B Pratiwi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012020

Fuzzy modeling network type 2 and principal component analysis for the diagnosis of diabetic retinopathy

A Damayanti, S Maimunah and A B Pratiwi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012021

Metaheuristic Algorithms for Solving Multiple-Trips Vehicle Routing Problem with Time Windows (MTVRPTW)

A B Pratiwi, A Sasmito, Q S Istiqomah, M R Kurniawan and H Suprajitno

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012022

A Simulation of Shallow Water Wave Equation Using Finite Volume Method: Lax-Friedrichs Scheme

R. Setiyowati and Sumandi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012023

On Radical Property of Cross Polyomino Ideal

Yoshua Yonatan Hamonangan and Intan Muchtadi-Alamsyah

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012024

Think Pair Share (TPS) Model Using Science, Technology, Engineering, Mathematics (STEM) Approach in Mathematics Learning

I Werdiningsih, Budiyo and H Pratiwi

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012025

On Vertex Irregular Total k -labeling and Total Vertex Irregularity Strength of Lollipop Graphs

Siti' Aisyah Nur Nirmah and Diari Indriati

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012026

Mathematical problem solving based on Kolb's learning style

W A Rokhima, T A Kusmayadi and L Fitriana

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012027

Modelling of Hypertension Risk Factors Using Logistic Regression to Prevent Hypertension in Indonesia

Putri Andriani and Nur Chamidah

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012028

Student's Difficulties in Solving Higher Order Thinking Skills (HOTS) Problem for Mathematics Based on Verbalizer Cognitive Style

A Rahmawatiningrum, T A Kusmayadi and L Fitriana

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012029

Analysis of Concept Understanding Student in Class X Inequalities Material

D N Sari, B Usodo and I Pramudya

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012030

Server Room Temperature & Humidity Monitoring Based on Internet of Thing (IoT)

Moehammad Alvan Prastoyo Utomo, Abdul Aziz, Winarno and Bambang Harjito

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012031

On Edge Irregular Total k -labeling and Total Edge Irregularity Strength of Barbell Graphs

Melli Aftiana and Diari Indriati

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012032

Total Edge Irregularity Strength of Arithmetic Book Graphs

Lucia Ratnasari, Sri Wahyuni, Yeni Susanti, Diah Junia Eksi Palupi and Budi Surodjo

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012033

Spatial Lag Fixed Effect Panel Model with Weights Queen Contiguity for Economic Growth Data of ASEAN Member Countries

R D Wisnumurti, H Pratiwi and S S Handajani

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012034

Optimal control of a mathematical model for Japanese encephalitis transmission

Heni Kharismawati, Fatmawati and Windarto

[+ Open abstract](#)
[View article](#)
[PDF](#)

OPEN ACCESS

012035

On the metric dimension of amalgamation of sunflower and lollipop graph and cavenham graph



Satria Adhi Wijaya and Tri Atmojo Kusmayadi
+ Open abstract View article PDF

OPEN ACCESS 012036
Design of Student Worksheet Based On Discovery Learning to Improve the Ability of Mathematics Reasoning Students of Class VII Junior High School
F Marian and Suparman
+ Open abstract View article PDF

OPEN ACCESS 012037
Modelling of Diarrheal Sufferers Percentage and Certified Drink Water Providers in East Java Using Bivariate Binomial Negative Regression Method
N A P Rahmayanti, Y S Yonani and M F F Mardianto
+ Open abstract View article PDF

OPEN ACCESS 012038
The Validity of Algebraic Module with Living Values Education for Junior High School
Y Ningrum, T A Kusmayadi and L Fitriana
+ Open abstract View article PDF

OPEN ACCESS 012039
Graceful Chromatic Number of Unicyclic Graphs
R. Alfari, Dafik, R.M. Prihandini, R. Adawiyah, E. R. Albirri and I. H. Agustin
+ Open abstract View article PDF

OPEN ACCESS 012040
The γ -Dimension of Images of Bi-Lipschitz Function
S Wibowo, V Y Kurniawan and Siswanto
+ Open abstract View article PDF

OPEN ACCESS 012041
Indonesia's Inflation Analysis Using Hybrid Fourier - Wavelet Multiscale Autoregressive Method
Suparti, R Santoso, A Prahutama and A R Devi
+ Open abstract View article PDF

OPEN ACCESS 012042
Some properties on fuzzy chromatic number of union of fuzzy graphs through α -cut graphs coloring
I Rosyida, Widodo, Ch R Indrati and D Indriati
+ Open abstract View article PDF

OPEN ACCESS 012043
The Influence of Teacher's Conception of Teaching and Learning on Their Teaching Practice
Masduki, Suwarsono and M T Budiarto
+ Open abstract View article PDF

OPEN ACCESS 012044
Enhancing the Student's Reasoning Ability in Solving Real Number System Problems Using the Concept Map
R P Khotimah, Masduki and J Sungkono
+ Open abstract View article PDF

OPEN ACCESS 012045
Functional thinking profile of mathematics problems based on gender in senior high school
L N Pamungkas, T A Kusmayadi and L Fitriana
+ Open abstract View article PDF

OPEN ACCESS 012046
The r-dynamic chromatic number of corona of order two of any graphs with complete graph
Dafik, I H Agustin, D A R Wardani, B J Septory, A I Kristiana and E Y Kurniawati
+ Open abstract View article PDF

OPEN ACCESS 012047
The local antimagic total vertex coloring of graphs with homogeneous pendant vertex
Elsa Yuli Kurniawati, Ika Hesti Agustin, Dafik and Marsidi
+ Open abstract View article PDF

OPEN ACCESS 012048
Retinal Diseases Classification Using Levenberg-Marquath (LM) Learning Algorithm for Pi Sigma Network (PSN) and Principal Component Analysis (PCA) Methods
S Swasono, A Damayanti and A B Pratiwi
+ Open abstract View article PDF

OPEN ACCESS 012049
Classification of Javanese Language Level on Articles Using Multinomial Naive Bayes and N-Gram Methods
A P Ardhana, D E Cahyani and Winarno
+ Open abstract View article PDF

OPEN ACCESS 012050
Semiparametric approaches for modelling mortality with application to university employee data
Danardono, R S Utami, R Fitriani, Zulaela and S H Kartiko
+ Open abstract View article PDF

OPEN ACCESS 012051
Optimizing Range Norm of The Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components
Siswanto, A Suparwanto and M A Rudhito
+ Open abstract View article PDF

OPEN ACCESS 012052
Intensity of Teak Wood Production in Central Java Using Point Process by Counting Measure Approach
Respatiuluan, L Jatningsih, Y Susanti, S S Handayani, H Pratiwi, I Slamet and Hartatik
+ Open abstract View article PDF

JOURNAL LINKS

- Journal home
- Journal Scope
- Information for organizers
- Information for authors
- Contact us
- Reprint services from Curran Associates

PAPER • OPEN ACCESS

Optimizing Range Norm of The Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components

To cite this article: Siswanto *et al* 2019 *J. Phys.: Conf. Ser.* **1306** 012051

View the [article online](#) for updates and enhancements.

You may also like

- [Controlling optical responses through local dielectric resonance in nanometre metallic clusters](#)
Chen Liang-Liang, Gu Ying, Wang Li-Jin et al.
- [The strain and temperature scaling law for the critical current density of a jelly-roll Nb₃Al strand in high magnetic fields](#)
Simon A Keys, Norikiyo Koizumi and Damian P Hampshire
- [Robust matrices in the interval max-plus algebra](#)
Siswanto, Pangadi and S B Wiyono



The Electrochemical Society
Advancing solid state & electrochemical science & technology

242nd ECS Meeting

Oct 9 – 13, 2022 • Atlanta, GA, US

Abstract submission deadline: **April 8, 2022**

Connect. Engage. Champion. Empower. Accelerate.

MOVE SCIENCE FORWARD



Submit your abstract



Optimizing Range Norm of The Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components

Siswanto, A Suparwanto, and M A Rudhito

¹ Department of Mathematics, Faculty of Mathematics and Natural Sciences, Sebelas Maret University, Surakarta 57126, Indonesia.

² Department of Mathematics, Faculty of Mathematics and Natural Sciences, Gadjah Mada University, Yogyakarta 55281, Indonesia.

³ The Program Study Mathematics in Education, Faculty of Teacher Training and Education, Sanata Dharma University, Yogyakarta 55282, Indonesia.

E-mail : sis.mipa@staff.uns.ac.id

Abstract. Let \mathbb{R} be the set of all real numbers and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$ whose $\varepsilon = -\infty$. Max-plus algebra is the set \mathbb{R}_ε that is equipped two operations maximum and addition. It can be formed matrices in the size of $m \times n$ whose elements belong to \mathbb{R}_ε , called matrix over max-plus algebra. Optimizing range norm of the image set of matrix over max-plus algebra with prescribed components has been discussed. Interval Max-Plus Algebra is the set $I(\mathbb{R})_\varepsilon = \{x = [\underline{x}, \bar{x}] | \underline{x}, \bar{x} \in \mathbb{R}, \varepsilon < \underline{x} \leq \bar{x}\} \cup \{\varepsilon\}$ with $\varepsilon = [\varepsilon, \varepsilon]$, is equipped with two operations maximum ($\overline{\oplus}$) and addition ($\overline{\otimes}$). The set of all matrices in the size of $m \times n$ whose elements belong to $I(\mathbb{R})_\varepsilon$, called matrix over interval max-plus algebra. Optimizing range norm of the image set of matrix over interval max-plus algebra has been discussed. In this paper, we will discuss optimizing range norm of the image set of matrix over interval max-plus algebra with prescribed components.

1. Introduction

Let \mathbb{R} be the set of all real numbers and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$ where $\varepsilon = -\infty$. Max-Plus algebra is the set \mathbb{R}_ε that is equipped two operations maximum (\oplus) and addition (\otimes). It can be formed matrices in the size of $m \times n$ whose elements belong to \mathbb{R}_ε . Max-plus algebra has been used to model and analyze algebraically planning problems, communication, production system, queueing system with finite capacity, parallel computation, and traffic (Bacelli; *et, al.* [1]). Min-Plus algebra is the set $\mathbb{R}_{\varepsilon'} = \mathbb{R} \cup \{\varepsilon'\}$ where $\varepsilon' = -\infty$ which is equipped two operations minimum (\oplus') and addition (\otimes'). Tam [3] also discuss about complete max-plus algebra that is the $\overline{\mathbb{R}}_\varepsilon = \mathbb{R} \cup \{\varepsilon, \varepsilon'\}$ is equipped with operations \oplus and \otimes , while complete min-plus algebra is $\overline{\mathbb{R}}_{\varepsilon'} = \mathbb{R} \cup \{\varepsilon, \varepsilon'\}$ which is equipped operations \oplus' and \otimes' . Then $\overline{\mathbb{R}}_\varepsilon$ and $\overline{\mathbb{R}}_{\varepsilon'}$ are written as $\overline{\mathbb{R}}$.

The set of all matrices in the size of $m \times n$ over \mathbb{R}_ε , $\mathbb{R}_{\varepsilon'}$ and $\overline{\mathbb{R}}$ can be formed. It is called the set of all matrices over max-plus algebra if its components belong to \mathbb{R}_ε and denoted by $\mathbb{R}_\varepsilon^{m \times n}$. Moreover, if $n = 1$ we obtain the set of all vectors over max-plus algebra, namely $\mathbb{R}_\varepsilon^m = \{(x_1, x_2, \dots, x_m)^T | x_1, x_2, \dots, x_m \in \mathbb{R}_\varepsilon\}$. If $m = n$, $\mathbb{R}_\varepsilon^{n \times n}$ is idempotent semiring with operations \oplus and



\otimes (Farlow [2], Akian, *et. al.* [4], Konigsberg [12]). Tam [3] gives an example of application of max-plus algebra in the production system. The production matrix $A = (A_{ij}) \in \mathbb{R}_\varepsilon^{n \times n}$ with A_{ij} shows the time is used of the production process from machine j to i , while vector $x(k) = (x_i(k)) \in \mathbb{R}_\varepsilon^n$ where $x_i(k)$ is the starting time of i machine at the k^{th} stage. In this production process we obtain the equation $x(k+1) = A \otimes x(k)$.

One of the criteria is used by manufacturers is that the production process runs periodically with period λ , such that $x(k+1) = \lambda \otimes x(k)$. From equations $x(k+1) = A \otimes x(k)$ and $x(k+1) = \lambda \otimes x(k)$ are obtained $A \otimes x(k) = \lambda \otimes x(k)$. The eigenvalue and eigenvector problem of matrix A is problem of how to find eigenvalue λ and eigen vector $x(k)$ such that $A \otimes x(k) = \lambda \otimes x(k)$.

In the real life situation, there are several ways for the manufacturer to determine the starting time of each machine. One of the ways is by choosing eigenvector as the starting time, so that the system will immediately reach the steady state; that is the process works periodically with the eigenvalue as its periods.

However, in reality, there are more than one independent eigenvectors for the manufacturer to choose. In that case, a set of linear combination of the independent eigenvectors is constructed, and as such, the manufacturer needs an additional criteria to choose one element of the set. The additional criteria is by considering the difference between the largest and the smallest of the starting time of each machine. The difference is expressed as range norm of the starting time of each machine. Manufacturer can optimize the range norm of the starting time of each machine (Tam [3], Butkovic and Tam [6]). There are some determining factors of a manufacturer to optimize vector of range norm of the starting time of machine, that are the readiness of the raw material, the availability of the resources, and the distribution of the product. In solving the problems of a production system, Tam [3] has discussed optimizing range norm of the image set of matrix over max-plus algebra.

Besides the criteria that have been submitted previously, manufacturers can also determine the start time of the machine began to work on certain machines. In this case, means that there are components that have been determined at the initial time vector machine start working. Furthermore, manufacturers only determine the initial time for the machine left so initial time vector is an eigenvector with optimal range norm.

Based on Kreinovich's idea [11] about the probability to give an estimation of the period of time of a certain process and Rudhito [5] about generalization of max-plus algebra i.e. interval max-plus algebra, we will to expand concept in max-plus algebra which has been discussed by Tam [3] in interval max-plus algebra. Siswanto, *et. al.* [9] has discussed about existence of solution to the system of linear Equations in interval max-Plus algebra. Siswanto, *et. al.* [8] has discussed the optimizing range norm of the image set of matrix over interval max-plus algebra. In this paper, we will discuss how to minimize and maximize range norm of the image set of matrix over interval max-plus algebra with prescribed components. Specially, about optimizing range norm of the image set of matrix over interval max-plus algebra with one component is prescribed and another one is unprescribed.

Before discussing the result of this paper, several concepts which support the discussion are discussed. These are minimizing and maximizing range norm of the image set of a matrix over max-plus algebra with prescribed components. Other than that optimizing range norm of the image set of a matrix over interval max-plus algebra.

2. Preliminaries

2.1. Minimizing Range Norm

The following the concepts about the problem minimizing of range norm of eigenvector of matrices over maks-plus algebra will be presented.

Definition 2.1. If $x \in \mathbb{R}^m$ then we will denote the function $\delta(x) = \sum_{i \in M}^\oplus x_i - \sum_{i \in M}^\ominus x_i$ and called the range norm of x , i.e. range norm is largest value in x – smallest value in x .

Problem 2.1. Given a matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve $\delta(b)$ minimum, subject to $b \in \text{Im}(A)$.

Definition 2.2. If $x \in \mathbb{R}_\varepsilon^m$ then the function $\tilde{\delta}(x) = \sum_{x_i \neq \varepsilon}^\oplus x_i - \sum_{x_i \neq \varepsilon}^{\oplus'} x_i$ is the range norm of x after only considering the finite components.

Problem 2.2. Given a matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve $\tilde{\delta}(b)$ minimum, subject to $b \in \text{Im}(A)$.

Futhermore, the minimizing of a norm range eigenvector with prescribed components are presented.

Definition 2.3. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $1 \leq p \leq m - 1$. We will denote A_1^p to be the matrix that consist of the first p row(s) of A and A_2^p to be the matrix generated by deleting the first p row(s) of A .

Problem 2.3. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $c \in \text{Im}(A_1^p) \cap \mathbb{R}^p$, $1 \leq p \leq m - 1$. Find $d \in \mathbb{R}^{m-p}$ such that $\delta(b)$ minimum, subject to $b \in \text{Im}(A)$ and $b^T = (c^T, d^T)$.

Before we discussed the problem for all case of p , the two special cases for this problem when $p = 1$ and $p = m - 1$ are presented.

Theorem 2.4. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic and $c \in \text{Im}(A_1^p) \cap \mathbb{R}^p$, $1 \leq p \leq m - 1$. Suppose that $\bar{b} \in \mathbb{R}^m$ is an optimal solution to Problem 1.1 for A and $d \in \mathbb{R}^{m-p}$ is an optimal solution for Problem 2.3 for A and c , then $\delta(\bar{b}) \leq \delta(b)$

where $b^T = (c^T, d^T)$.

Theorem 2.4 guarantee lower bound of range norm which is obtained from optimal solution Problem 2.3. Using this result of the Problem 2.3, be obtained the solution of $p = 1$. Suppose that only the starting time of one machine is prescribed, then by Theorem 2.4 can be obtained the result as the following.

Theorema 2.5. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic and $c = (c_1) \in \mathbb{R}$ be the vector of prescribed component. Suppose that $b = A \otimes (A^* \otimes' 0)$ and $d = (c_1 \otimes b_1^{-1}) \otimes b$ where b_1 is the first component of b . Then the vector $(d_2, \dots, d_m)^T$ is an optimal solution to Problem 2.3.

Futhermore, the case of $p = m - 1$, used the following theorem.

Theorema 2.6. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic, $c \in \text{Im}(A_1^p) \cap \mathbb{R}^p$ and \tilde{d} be an optimal solution for Problem 2.3. Suppose $\bar{x} = (A_1^p)^* \otimes' c$ and $\bar{d} = A_2^p \otimes \bar{x}$ then $\tilde{d} \leq \bar{d}$.

2.2. Maximizing Range Norm

The following the concepts be related with the problem maximizing of eigenvector range norm of matrices over maks-plus algebra will be presented.

Problem 2.4. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve $\delta(b)$ maximum, subject to $b \in \text{Im}(A)$.

The maximizing range norm of the image set of a matrix over max-plus algebra with prescribed components will be presented.

Problem 2.5. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $c \in \text{Im}(A_1^p) \cap \mathbb{R}^p$, $1 \leq p \leq m - 1$, find $d \in \mathbb{R}^{m-p}$ such that $\delta(b)$ maximum, subject to $b \in \text{Im}(A)$ and $b^T = (c^T, d^T)$.

Before discussing the problem for all case of p , the two special cases for this problem when $p = 1$ and $p = m - 1$ are presented. First discussed the cases where only one component prescribed that is $p = 1$. In the case of this special, Problem 2.5 can be considered the same as it counterpart Problems 2.4. This can be seen in the following theorem.

Theorem 2.7. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $c = (c_1) \in \mathbb{R}$ be the prescribed component. Suppose that $b \in \text{Im}(A)$ such that b is an optimal solution to Problem 2.4 and $d = (c_1 \otimes b_1^{-1}) \otimes b$ where b_1 is the first component of b , then vector $(d_2, \dots, d_m)^T$ is an optimal solution to Problem 2.5.

Teorema 2.8. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic and non finite, i.e. $\exists i \in M, j \in N$ such that $a_{ij} = \varepsilon$. Suppose that $c = (c_1) \in \mathbb{R}$ is the vector of prescribed component then Problem 2.5 is unbounded.

Futhermore, we will consider the case when $p = m - 1$.

Theorem 2.9. Let $A \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic, $c \in \text{Im}(A_1^p) \cap \mathbb{R}^p$ and \tilde{d} be an optimal solution to Problem 2.5. Suppose that $\bar{x} = (A_1^p)^* \otimes' c$ and $\bar{d} = A_2^p \otimes \bar{x}$ then $\tilde{d} \leq \bar{d}$.

Besides the concepts above namely about max-plus algebra, interval min-plus algebra, complete interval max-plus algebra, and complete interval algebra min-plus (Rudhito [5]; Siswanto [7]), also are used the concepts about optimizing a range norm the set of image matrix over interval algebra max-plus (Siswanto [10]).

2.3. Minimizing Range Norm of The Image Set of Matrix over Interval Max Plus Algebra

Definition 2.10. Suppose that $x \in I(\mathbb{R})^m$ where $x \approx [\underline{x}, \bar{x}]$; $\underline{x}, \bar{x} \in \mathbb{R}^m$. The function $\delta(x) = [\min(\delta(\underline{x}), \delta(\bar{x})), \delta(\bar{x})]$ is called the range norm of x .

Definition 2.11. Given that matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$; $\underline{A}, \bar{A} \in \mathbb{R}_\varepsilon^{m \times n}$, then $\text{Im}(A) = \{A \otimes p \mid p \in I(\mathbb{R})_\varepsilon^n\}$.

Problem 2.6 Given that matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, solve $\delta(b)$ minimize, subject to $b \in \text{Im}(A)$, $b \approx [\underline{b}, \bar{b}]$.

Definisi 2.12. Given that matrix $b \in I(\mathbb{R})_\varepsilon^m$ with $b \approx [\underline{b}, \bar{b}]$. The function $\delta(b)$ is said minimum if and only if $\delta(\underline{b})$ and $\delta(\bar{b})$ are minimum.

Definition 2.13. Suppose that $x \in I(\mathbb{R})^m$ with $x \approx [\underline{x}, \bar{x}]$. The function $\tilde{\delta}(x) = [\min(\tilde{\delta}(\underline{x}), \tilde{\delta}(\bar{x})), \tilde{\delta}(\bar{x})]$ is the range norm of x , after only considering the finite components.

Problem 2.7. Given that matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, $A \approx [\underline{A}, \bar{A}]$. Solve $\tilde{\delta}(b)$ minimize, subject to $b \in \text{Im}(A)$, $b \approx [\underline{b}, \bar{b}]$.

Definition 2.14. Given that matrix $b \in I(\mathbb{R})_\varepsilon^m$ with $b \approx [\underline{b}, \bar{b}]$. The function $\tilde{\delta}(b)$ is said minimum if and only if $\tilde{\delta}(\underline{b})$ and $\tilde{\delta}(\bar{b})$ are minimum.

Definition 2.15. Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, $A \approx [\underline{A}, \bar{A}]$ is called double $I(\mathbb{R})$ -astic if A is double \mathbb{R} -astic for every $A \in [\underline{A}, \bar{A}]$.

Theorem 2.16. The matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ where $A \approx [\underline{A}, \bar{A}]$ is double $I(\mathbb{R})$ -astic if and only if \underline{A} is double \mathbb{R} -astic.

Lemma 2.17. Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ where $A \approx [\underline{A}, \bar{A}]$. If $A = [A_{ij}]$ be doubly $I(\mathbb{R})$ -astic, $b \approx [\underline{b}, \bar{b}]$, $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' v_0)$, $\bar{b} = \alpha \otimes (\bar{A} \otimes (\bar{A}^* \otimes' v_0))$ where $v_0 \in \mathbb{R}^m$ be a vector whose every components is equal to a component 0 and $\alpha = \max_i ((\underline{A} \otimes (\underline{A}^* \otimes' v_0))_i - (\bar{A} \otimes (\bar{A}^* \otimes' v_0))_i) \geq 0$ then

- i. $b \leq ([0, \alpha], [0, \alpha], \dots, [0, \alpha])^T$,
- ii. $\underline{b}_i = 0$ for some $i \in M$,
- iii. $\bar{b}_j = \alpha$ for some $j \in M$.

Lemma 2.18. If $x, y \in I(\mathbb{R})^m$ and $\alpha \in I(\mathbb{R})$ where $x \approx [\underline{x}, \bar{x}]$, $y \approx [\underline{y}, \bar{y}]$ and $a \approx [\underline{a}, \bar{a}]$ then $\delta(x) = \delta(a \otimes x)$.

Theorem 2.19. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be double $I(\mathbb{R})$ -astic and $v_0 \in \mathbb{R}^m$ be a vector whose every components is equal to a constant 0 then $b \approx [\underline{b}, \bar{b}]$ is a solution for Problem 2.6 where $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' v_0)$, $\bar{b} = \alpha \otimes (\bar{A} \otimes (\bar{A}^* \otimes' v_0))$ and $\alpha = \max_i ((\underline{A} \otimes (\underline{A}^* \otimes' v_0))_i - (\bar{A} \otimes (\bar{A}^* \otimes' v_0))_i)$.

To determine solution of Problem 2.7 the next theorem is needed :

Theorem 2.20. Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, $A \approx [\underline{A}, \bar{A}]$ double $I(\mathbb{R})$ -astic and $v_a \approx [v_{\underline{a}}, v_{\bar{a}}] \in I(\mathbb{R})^m$ be a vector whose every components is equal to an interval $a \approx [\underline{a}, \bar{a}]$. If $x \in I(\mathbb{R})_\varepsilon^n$ such that $b = A \otimes x$, $b \approx [\underline{b}, \bar{b}] \in \text{Im}(A)$ is solution of Problem 2.7 then $\forall j \in N$ satisfy the conditions :

- a. $x_j = \varepsilon$ or
- b. $x_j = \left[(\underline{A}^* \otimes' v_{\underline{a}})_{j'}, (\bar{A}^* \otimes' v_{\bar{a}})_{j'} \right]$.

2.4. Maximizing Range Norm of The Image Set of Matrix over Interval Max-Plus Algebra

Problem 2.8 Given that matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, solve : maximize $\delta(b)$ subject to $b \in \text{Im}(A)$, $b \approx [\underline{b}, \bar{b}]$.

Definition 2.22. The function $\delta(b)$ is called maximum if and only if $\delta(\bar{b})$ and $\delta(\underline{b})$ are maximum.

Theorem 2.23. If $A \in I(\mathbb{R})^{m \times n}$ then

$$\delta(b) \leq \left[\min \left(\max_{j=1,2,\dots,n} \delta(\underline{A}_j), \max_{j=1,2,\dots,n} \delta(\bar{A}_j) \right), \max_{j=1,2,\dots,n} \delta(\bar{A}_j) \right] \text{ for every } b \in \text{Im}(A).$$

Theorem 2.24. Suppose that $M = \{1, 2, \dots, m\}$ and $N = \{1, 2, \dots, n\}$. If $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and there are $i \in M, j \in N$ such that $a_{ij} = \varepsilon$ then Problem 2.8 unbounded.

3. Results and Discussion

This section will discuss the results of this research about optimizing range norm of the image set of matrix over interval max-plus algebra with some components are prescribed.

3.1. Minimizing Range Norm

Some definitions and theorems also problems the minimizing of a norm range the image set of matrix where one component is prescribed and one component is unprescribed, given as follows.

Definition 3.1. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ and $1 \leq p \leq m - 1$. We will denote A_1^p to be the matrix that consists of the first p row (rows) of A and A_2^p to be the matrix generated by deleting the first p row (rows) of A .

Problem 3.1. Given $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ and $c \in \text{Im}(A_1^p) \cap I(\mathbb{R})^p$, $1 \leq p \leq m - 1$. Find $d \in I(\mathbb{R})^{m-p}$ such that $\delta(b)$ minimum subject to $b \in \text{Im}(A)$ and $b^T = (c^T, d^T)$.

To solve this problem, started by the two special cases for $p = 1$ and $p = m - 1$. Let us consider the following theorem.

Theorem 3.2 Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and $c \in \text{Im}(A_1^p) \cap I(\mathbb{R})^p$, $1 \leq p \leq m - 1$. Suppose that e is an optimal solution to Problem 2.6 for A and $d \in I(\mathbb{R})^{m-p}$ is an optimal solution to Problem 3.1 for A and c then $\delta(e) \leq \delta(b)$ where $b^T = (c^T, d^T)$.

Proof. Suppose that $A \approx [\underline{A}, \bar{A}]$, $b \approx [\underline{b}, \bar{b}]$, $c \approx [\underline{c}, \bar{c}]$, $d \approx [\underline{d}, \bar{d}]$ and $e \approx [\underline{e}, \bar{e}]$. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and $c \in \text{Im}(A_1^p) \cap I(\mathbb{R})^p$, $1 \leq p \leq m - 1$. Therefore $\underline{A}, \bar{A} \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic and $\underline{c} \in \text{Im}(\underline{A}_1^p) \cap \mathbb{R}^p$, $\bar{c} \in \text{Im}(\bar{A}_1^p) \cap \mathbb{R}^p$. Suppose e is an optimal solution to Problem 2.6 for A and $d \in I(\mathbb{R})^{m-p}$ is an optimal solution to Problem 3.1 for A and c . Since $e \in \text{Im}(A)$, so $\underline{e} \in \text{Im}(\underline{A})$, $\bar{e} \in \text{Im}(\bar{A})$. with \underline{e} is an optimal solution to Problem 2.1 for \underline{A} and \bar{e} is an optimal solution to Problem 2.1 for \bar{A} . While $\underline{d} \in \mathbb{R}^{m-p}$ is an optimal solution to Problem 2.3 for \underline{A} and \underline{c} and $\bar{d} \in \mathbb{R}^{m-p}$ is an optimal solution to Problem 2.3 for \bar{A} and \bar{c} . According to Theorem 2.4, $\delta(\underline{e}) \leq \delta(\underline{b})$ and $\delta(\bar{e}) \leq \delta(\bar{b})$ where $\underline{b}^T = (\underline{c}^T, \underline{d}^T)$ and $\bar{b}^T = (\bar{c}^T, \bar{d}^T)$. As a result $\delta(e) \leq \delta(b)$ where $b^T = (c^T, d^T)$. ■

Theorem 3.2 provides a lower bound for the range norm of the optimal solution to Problem 3.1. Suppose that only the starting time of one machine is prescribed then by Theorem 2.20 and Theorem 3.2 are obtained results as follows.

Theorem 3.3 Let $A \approx [\underline{A}, \bar{A}] \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic, $c = ([\underline{c}_1, \bar{c}_1]) \in I(\mathbb{R})$ be the vector of prescribed components, $v_0 \in \mathbb{R}^m$ be a vector whose every component is equal to a constant 0. If $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' v_0)$, $\bar{b} = \alpha \otimes (\bar{A} \otimes (\bar{A}^* \otimes' v_0))$ where $\alpha = \max_i ((\underline{A} \otimes (\underline{A}^* \otimes' v_0))_i - (\bar{A} \otimes (\bar{A}^* \otimes' v_0))_i) \geq 0$, $\underline{d} = (\underline{c}_1 \otimes \underline{b}_1^{-1}) \otimes \underline{b}$, $\bar{d} = (\bar{c}_1 \otimes \bar{b}_1^{-1}) \otimes \bar{b}$ when $\underline{d} \leq \bar{d}$ then $d_1 = (d_2, d_3, \dots, d_m)$ is an optimal solution to Problem 3.1.

Proof. Since $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic then \underline{A}, \bar{A} be doubly \mathbb{R} -astic. Since $c = (c_1) = ([\underline{c}_1, \bar{c}_1]) \in I(\mathbb{R})$ then $\underline{c} = (\underline{c}_1)$, $\bar{c} = (\bar{c}_1) \in \mathbb{R}$ the vector of prescribed components. If $\underline{b} = \underline{A} \otimes$

$(\underline{A}^* \otimes' v_0)$ and $\bar{b} = \alpha \otimes (\bar{A} \otimes (\bar{A}^* \otimes' v_0))$ where $\alpha = \max_i ((\underline{A} \otimes (\underline{A}^* \otimes' v_0))_i - (\bar{A} \otimes (\bar{A}^* \otimes' v_0))_i) \geq 0$, $\underline{d} = (\underline{c}_1 \otimes \underline{b}_1^{-1}) \otimes \underline{b}$, $\bar{d} = (\bar{c}_1 \otimes \bar{b}_1^{-1}) \otimes \bar{b}$ when $\underline{d} \leq \bar{d}$. According to Theorem 2.5, the vector $(\underline{d}_2, \dots, \underline{d}_m)^T$ is an optimal solution to Problem 2.3 for \underline{c}_1 and \underline{A} , while $(\bar{d}_2, \dots, \bar{d}_m)^T$ is an optimal solution to Problem 2.3 for \bar{c}_1 and \bar{A} . Therefore vector $(\underline{d}_2, \dots, \underline{d}_m)^T$ where $\underline{d}_2 \approx [\underline{d}_2, \bar{d}_2], \dots, \underline{d}_m \approx [\underline{d}_m, \bar{d}_m]$ is an optimal solution to Problem 3.1 for c and A . ■

Next, in the case of $p = m - 1$ we needed to find the fisibel solution for d . Given $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and $c \in \text{Im}(A^p) \cap I(\mathbb{R})^p$ be the vector of prescribed components that $c \approx [c, \bar{c}]$, then there are $\hat{x} \in I(\mathbb{R})_\varepsilon^n$ where $\hat{x} = (A_1^{m-1})^* \otimes' c$ such that $c = A^{m-1} \otimes \hat{x}$. By using \hat{x} is obtained $g = A_2^{m-1} \otimes \hat{x}$. According to equation,

$$A \otimes \hat{x} = \begin{pmatrix} A_1^{m-1} \\ A_2^{m-1} \end{pmatrix} \otimes \hat{x} = \begin{pmatrix} A_1^{m-1} \otimes \hat{x} \\ A_2^{m-1} \otimes \hat{x} \end{pmatrix} = \begin{pmatrix} c \\ g \end{pmatrix} = b,$$

that g is the optimal solution to Problem 3.1.

Suppose that \underline{L} and \bar{L} represent the minimum value of \underline{c} and \bar{c} , while \underline{U} dan \bar{U} represent the maximum value of \underline{c} dan \bar{c} , that is

- $\underline{L} = \min_{i=1, \dots, m-1} \underline{c}_i$ and $\bar{L} = \min_{i=1, \dots, m-1} \bar{c}_i$,
- $\underline{U} = \max_{i=1, \dots, m-1} \underline{c}_i$ and $\bar{U} = \max_{i=1, \dots, m-1} \bar{c}_i$.

Since $g \approx [\underline{g}, \bar{g}]$, therefore there are three cases for \underline{g} , i.e.

- $\underline{L} \leq \underline{g} \leq \underline{U}$ where $\delta(\underline{b}) = \delta(\underline{c}) = \underline{U} - \underline{L}$,
- $\underline{g} < \underline{L}$ where $\delta(\underline{b}) = \underline{L} - \underline{g}$,
- $\underline{U} < \underline{g}$ where $\delta(\underline{b}) = \underline{g} - \underline{U}$.

Also there are three cases for \bar{g} , i.e.

- $\bar{L} \leq \bar{g} \leq \bar{U}$ where $\delta(\bar{b}) = \delta(\bar{c}) = \bar{U} - \bar{L}$,
- $\bar{g} < \bar{L}$ where $\delta(\bar{b}) = \bar{L} - \bar{g}$,
- $\bar{U} < \bar{g}$ where $\delta(\bar{b}) = \bar{g} - \bar{U}$.

According to the discussion in max-plus algebra, casses 1 and 2 and also casses 1' and 2' are solution to Problem 2.3 for $p = m - 1$. While casses 3 and 3', may be there are \underline{g} and \bar{g} such that $\delta(\underline{b})$ and $\delta(\bar{b})$ are optimal. The probability there exis \underline{g} and \bar{g} are determined the same way in max-plus algebra.

Next, from combination casses 1 and 1', cases 1 dan 2', cases 2 and 1', and also cases 2 and 2' can be obtained an optimal solution to Problem 3.1. The next theorem which is indicated upper bound of optimal solution to Problem 3.1.

Theorem 3.4. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic, $c \in \text{Im}(A^p) \cap \mathbb{R}^p$ and $f \in I(\mathbb{R})^{m-p}$ is an optimal solution to Problem 3.1. Suppose $\underline{x} = (\underline{A}_1^p)^* \otimes' \underline{c}$ and $\bar{x} = (\bar{A}_1^p)^* \otimes' \bar{c}$. If $\underline{g} = \underline{A}_2^p \otimes \underline{x}$ and $\bar{g} = \alpha \otimes (\bar{A}_2^p \otimes \bar{x})$ with $\alpha = \max_i ((\underline{A}_2^p \otimes' \underline{x})_i - ((\bar{A}_2^p \otimes' \bar{x}))_i)$ then $g \approx [\underline{g}, \bar{g}]$ and $f \leq g$.

Proof. Suppose $A \approx [\underline{A}, \bar{A}]$, $c \approx [\underline{c}, \bar{c}]$ dan $f \approx [\underline{f}, \bar{f}]$. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic, $c \in \text{Im}(A^p) \cap \mathbb{R}^p$ and $f \in I(\mathbb{R})^{m-p}$ is an optimal solution to Problem 3.1. Therefore $\underline{A}, \bar{A} \in \mathbb{R}_\varepsilon^{m \times n}$, $\underline{c} \in \text{Im}(\underline{A}_1^p) \cap \mathbb{R}^p$, $\bar{c} \in \text{Im}(\bar{A}_1^p) \cap \mathbb{R}^p$ with \underline{f} is an optimal solution to Problem 2.3 for \underline{A} and \underline{c} , while \bar{f} is optimal solution to Problem 2.3 for \bar{A} and \bar{c} . Suposse $\underline{x} = (\underline{A}_1^p)^* \otimes' \underline{c}$, $\bar{x} = (\bar{A}_1^p)^* \otimes' \bar{c}$, According to

Theorem 2.9, if $\underline{g} = \underline{A}_2^p \otimes \underline{x}$, $\overline{g'} = \overline{A}_2^p \otimes \overline{x}$ then $\underline{f} \leq \underline{g}$ and $\overline{f} \leq \overline{g'}$. Since $\overline{g} = \alpha \otimes (\overline{A}_2^p \otimes \overline{x}) = \alpha \otimes \overline{g'}$ where $\alpha = \max_i \left((\underline{A}_2^p \otimes' \underline{x})_i - ((\overline{A}_2^p \otimes' \overline{x}))_i \right)$ such that $\underline{g} \approx [\underline{g}, \overline{g}]$ and $\underline{f} \leq \underline{g}$. ■

3.2. Maximizing range norm

In the section will discuss about maximize a norm range of the image set of a matrix with some elements vector had been determined. Used an assumption that same as minimize a norm range that elements a vector that determined finite, image of the matrix the top namely A_1^p is $c \in \text{Im}(A_1^p) \cap I(\mathbb{R})^p$ and the elements a vector that will be determined also finite.

Problem 3.2. Given $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ and $c \in \text{Im}(A_1^p) \cap I(\mathbb{R})^p$, $1 \leq p \leq m-1$. Find $d \in I(\mathbb{R})^{m-p}$ such that $\delta(b)$ maximum subject to $b \in \text{Im}(A)$ and $b^T = (c^T, d^T)$.

As in minimizing problem, to solve this problem, started by the two special cases $p = 1$ and $p = m-1$. Based on a case for $p = m-1$, it can be discussed in the case of common. Given the following theorem.

Theorem 3.5. Let $A \approx [\underline{A}, \overline{A}] \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic, $c = ([\underline{c}_1, \overline{c}_1]) \in I(\mathbb{R})$ be a vector of prescribed components. If $b \in \text{Im}(A)$ is an optimal solution to Problem 2.8 and $\underline{d} = (\underline{c}_1 \otimes \underline{b}_1^{-1}) \otimes \underline{b}$, $\overline{d} = (\overline{c}_1 \otimes \overline{b}_1^{-1}) \otimes \overline{b}$ when $\underline{d} \leq \overline{d}$ then $e = (c, d_1^1)^T \in \text{Im}(A)$ is an solution to Problem 3.2 with d_1^1 is the vector $d = (d_1, d_2, \dots, d_m) \approx [\underline{d}, \overline{d}]$ by deleting the first component.

Proof. Let $A \approx [\underline{A}, \overline{A}] \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic, so $\underline{A}, \overline{A}$ be doubly \mathbb{R} -astic. Then, $c = ([\underline{c}_1, \overline{c}_1]) \in I(\mathbb{R})$ be the vector of prescribed components. So $\underline{c} = (\underline{c}_1)$, $\overline{c} = (\overline{c}_1) \in \mathbb{R}$ be the vector of prescribed components. Since $b \in \text{Im}(A)$ is an optimal solution to Problem 2.8, so $\underline{b} \in \text{Im}(\underline{A})$ is an optimal solution to Problem 2.4 for \underline{A} , while $\overline{b} \in \text{Im}(\overline{A})$ is an optimal solution to Problem 2.4 for \overline{A} . Since, $\underline{d} = (\underline{c}_1 \otimes \underline{b}_1^{-1}) \otimes \underline{b}$, $\overline{d} = (\overline{c}_1 \otimes \overline{b}_1^{-1}) \otimes \overline{b}$ and $\underline{d} \leq \overline{d}$. According to Theorem 2.7, $\underline{e} = (\underline{c}_1, \underline{d}_1^1)^T \in \text{Im}(\underline{A})$ is a optimal solution to Problem 2.5 for \underline{c}_1 and \underline{A} with \underline{d}_1^1 is the vector $\underline{d} = (\underline{d}_1, \underline{d}_2, \dots, \underline{d}_m) \approx [\underline{d}, \overline{d}]$ by deleting the first component and $\overline{e} = (\overline{c}_1, \overline{d}_1^1)^T \in \text{Im}(\overline{A})$ is a solution to Problem 2.5 for \overline{c}_1 and \overline{A} with \overline{d}_1^1 is the vector $\overline{d} = (\overline{d}_1, \overline{d}_2, \dots, \overline{d}_m) \approx [\underline{d}, \overline{d}]$ by deleting the first component. Therefore, $e = (c, d_1^1)^T \in \text{Im}(A)$ is an solution to Problem 3.2 for c_1 and A with d_1^1 is the vector $d = (d_1, d_2, \dots, d_m) \approx [\underline{d}, \overline{d}]$ by deleting the first component. ■

Theorem 3.6. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and $\exists i \in M, j \in N$ such that $a_{ij} = \varepsilon$. If $c = (c_1) \in I(\mathbb{R})$ is the vector of prescribed then Problem 3.2 has unbounded solution.

Proof. Suppose $A \approx [\underline{A}, \overline{A}]$ and $a_{ij} = [\underline{a}_{ij}, \overline{a}_{ij}]$. Let $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and there are $i \in M, j \in N$ such that $a_{ij} = \varepsilon$. Therefore $\underline{A}, \overline{A} \in \mathbb{R}_\varepsilon^{m \times n}$ be doubly \mathbb{R} -astic and there are $i \in M, j \in N$ such that $\underline{a}_{ij} = \varepsilon$ and $\overline{a}_{ij} = \varepsilon$. Because $c = (c_1) \in I(\mathbb{R})$ is the vector of prescribed. Suppose $c \approx [\underline{c}, \overline{c}]$ so $\underline{c} = \underline{c}_1$, $\overline{c} = \overline{c}_1$. According to Theorem 2.8 there is $\underline{d} \in \mathbb{R}^{m-1}$ so the Problems 2.5 for \underline{c} and \underline{A} have unbounded solution and there is $\overline{d} \in \mathbb{R}^{m-1}$ so the Problems 2.5 for \overline{c} and \overline{A} have unbounded solution. As a result Problems 3.2 have unbounded solution. ■

Next, in the case of $p = m-1$ we need to find the fisibel solution for d . Given matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ be doubly $I(\mathbb{R})$ -astic and $c \in \text{Im}(A^p) \cap I(\mathbb{R})^p$ as constrain vector that $c \approx [\underline{c}, \overline{c}]$, then $\exists \hat{x} \in I(\mathbb{R})_\varepsilon^n$ where $\hat{x} = (A_1^{m-1})^* \overline{\otimes}' c$ such that $c = A^{m-1} \overline{\otimes} \hat{x}$. Using the vector \hat{x} will be obtained, $g = A_2^{m-1} \overline{\otimes} \hat{x}$.

It appears that g is a fisibel solution for Problem 2.2 from the following equation,

$$A \overline{\otimes} \hat{x} = \begin{pmatrix} A_1^{m-1} \\ A_2^{m-1} \end{pmatrix} \overline{\otimes} \hat{x} = \begin{pmatrix} A_1^{m-1} \overline{\otimes} \hat{x} \\ A_2^{m-1} \overline{\otimes} \hat{x} \end{pmatrix} = \begin{pmatrix} c \\ g \end{pmatrix} = b.$$

Suppose that \underline{L} and \bar{L} represent the minimum value of \underline{c} and \bar{c} , while \underline{U} dan \bar{U} represent the maximum value of \underline{c} dan \bar{c} , that is

- a. $\underline{L} = \min_{i=1, \dots, m-1} \underline{c}_i$ and $\bar{L} = \min_{i=1, \dots, m-1} \bar{c}_i$,
- b. $\underline{U} = \max_{i=1, \dots, m-1} \underline{c}_i$ and $\bar{U} = \max_{i=1, \dots, m-1} \bar{c}_i$.

Since $g \approx [\underline{g}, \bar{g}]$, therefore there are three cases for \underline{g} , i.e.

1. $\underline{L} \leq \underline{g} \leq \underline{U}$ where $\delta(\underline{b}) = \delta(\underline{c}) = \underline{U} - \underline{L}$,
2. $\underline{g} < \underline{L}$ where $\delta(\underline{b}) = \underline{L} - \underline{g}$,
3. $\underline{U} < \underline{g}$ where $\delta(\underline{b}) = \underline{g} - \underline{U}$.

Also there are three cases for \bar{g} , i.e.

- 1'. $\bar{L} \leq \bar{g} \leq \bar{U}$ where $\delta(\bar{b}) = \delta(\bar{c}) = \bar{U} - \bar{L}$,
- 2'. $\bar{g} < \bar{L}$ where $\delta(\bar{b}) = \bar{L} - \bar{g}$,
- 3'. $\bar{U} < \bar{g}$ where $\delta(\bar{b}) = \bar{g} - \bar{U}$.

According to the discussion in max-plus algebra, casses 1 and 3 and also casses 1' and 3' are solution of Problem 2.5 for $p = m - 1$. While casses 2 and 2', may be there are \underline{g} and \bar{g} such that $\delta(\underline{b})$ and $\delta(\bar{b})$ are optimal. The probability there exis \underline{g} and \bar{g} are determined the same way in max-plus algebra.

Next, from combination casses 1 and 1', cases 1 dan 3', cases 3 and 1', and also cases 3 and 3' can be obtained an optimal solution for Problem 3.2. Next, the theorem which is indicated upper bound of optimal solution for Problem 3.2.

Theorem 2.7. Let $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ be double $I(\mathbb{R})$ -astic $c \in Im(A^p) \cap I(\mathbb{R})^p$ and f is an optimal solution to problem 2.2. Suppose $\underline{y} = (\underline{A}_1^p)^* \otimes' \underline{c}$ and $\bar{y} = (\bar{A}_1^p)^* \otimes' \bar{c}$. If $\underline{g} = \underline{A}_2^p \otimes \underline{y}$ and $\bar{g} = \alpha \otimes (\bar{A}_2^p \otimes \bar{y})$ with $\alpha = \max \left(\left((\underline{A}_2^p \otimes' \underline{y})_i - \left((\bar{A}_2^p \otimes' \bar{y})_i \right) \right) \right)$ then $g \approx [\underline{g}, \bar{g}]$ and $f \leq g$.

Proof. Let $A \approx [\underline{A}, \bar{A}]$, $c \approx [\underline{c}, \bar{c}]$ dan $f \approx [\underline{f}, \bar{f}]$. Since $A \in Ie(\mathbb{R})_{\varepsilon}^{m \times n}$ be doubly $I(\mathbb{R})$ -astic so $\underline{A}, \bar{A} \in \mathbb{R}_{\varepsilon}^{m \times n}$ be doubly \mathbb{R} -astic, $c \in Im(A^p) \cap \mathbb{R}^p$ so $\underline{c} \in Im(\underline{A}_1^p) \cap \mathbb{R}^p$, $\bar{c} \in Im(\bar{A}_1^p) \cap \mathbb{R}^p$. Since $f \in I(\mathbb{R})^{m-p}$ is an optimal solution to Problem 3.2 so $\underline{f} \in \mathbb{R}^{m-p}$ is an optimal solution to Problem 2.5 for \underline{c} and \underline{A} , $\bar{f} \in \mathbb{R}^{m-p}$ is an optimal solution to Problem 2.5 for \bar{c} and \bar{A} . Let $\underline{y} = (\underline{A}_1^p)^* \otimes' \underline{c}$ and $\bar{y} = (\bar{A}_1^p)^* \otimes' \bar{c}$. According to Theorem 2.9, if $\underline{g} = \underline{A}_2^p \otimes \underline{y}$, $\bar{g}' = \bar{A}_2^p \otimes \bar{y}$ then $\underline{f} \leq \underline{g}$ and $\bar{f} \leq \bar{g}'$. Because $\bar{g} = \alpha \otimes (\bar{A}_2^p \otimes \bar{y}) = \alpha \otimes \bar{g}'$ with $\alpha = \max_i \left(\left((\underline{A}_2^p \otimes' \underline{y})_i - \left((\bar{A}_2^p \otimes' \bar{y})_i \right) \right) \right)$ then $g \approx [\underline{g}, \bar{g}]$ and $f \leq g$. ■

4. Conclusion

Based on the results of discussion, we obtain conclusion how to,

- a. Minimizing range norm of the set of matrix over interval max-plus algebra with component prescribed.
- b. Maximizing range norm of the set of matrix over interval max-plus algebra with component prescribed.

References

- [1] Bacelli, F., G. Cohen, G. J. Older, and J. P. Quadrat, *Synchronization and Linearity, An Algebra for Discrete Events System*, John Wiley and Sons, 2001.

- [2] K. G. Farlow, *Max-Plus Algebra*, Master's Thesis submitted to the Faculty of the Virginia Polytechnic Institute and State University in partial fulfillment of the requirements for the degree of Masters in Mathematics, 2009.
- [3] K. P. Tam, *Optimizing and Approximating Eigenvectors in Max Algebra*, A Thesis Submitted to The University of Birmingham for The Degree of Doctor of Philosophy (Ph.D), (2010).
- [4] M. Akian, G. Cohen, S. Gaubert, J. P. Quadrat, and M. Viot, *Max-Plus Algebra and Applications to System Theory and Optimal Control*, Proceedings of the International Congress of Mathematicians, Zurich, Switzerland, (1994).
- [5] M. A. Rudhito, *Fuzzy NumberMax-Plus Algebra and Its Application to Fuzzy Scheduling and Queuing Network Problems*, The Study Program S3 Mathematics Faculty of Mathematics and Natural Sciences : Gadjah Mada University, 2011. (Indonesian).
- [6] P. Butkovic, and K. P. Tam, *On Some Properties of The Image of a Max Linear Mapping*, Contemporary Mathematics, (495) (2009), pp. 115 – 126.
- [7] Siswanto, A. Suparwanto, M. A. Rudhito, *Minimizing Range Norm Of The Image Set Of Matrix Over Interval Max-Plus Algebra*, Proceedings of The National Seminar on Mathematics and Applications, Department of Mathematics, Faculty of Science and Technology, Airlangga University; ISBN : 978-602-14413-0-5, (2013). (Indonesian).
- [8] Siswanto, A. Suparwanto, M. A. Rudhito. *Eigenvector Space of a Matrix of Interval Max-Plus Algebra*, Jurnal Matematika & Sains (JMS). FMIPA ITB, (19) (1) (2014), pp. 8 – 15. (Indonesian).
- [9] Siswanto, A. Suparwanto, M. A. Rudhito, *Existence of Solution to the System of Linear Equations in Interval Max-Plus Algebra*, Proceedings of the National Conference of Mathematics XVII, Institute of Technology Surabaya, ISBN : 978-602-96426-3-6, (2014). (Indonesian).
- [10] Siswanto, A. Suparwanto, M. A. Rudhito, *Optimizing Range Norm of The Image Set of Matrix Over Interval Max-Plus Algebra*, Far East Journal of Mathematical Sciences, (19) (2016), pp. 17 – 32.
- [11] V. Kreinovich, *Why Intervals ? Why Fuzzy Numbers ? Towards a New Justification*, International Journal of Information Technology and Intelligent Computing. (2) (1) (2007).
- [12] Z. R. Konigsberg, *A Generalized Eigenmode Algorithm for Reducible Regular Matrices over the Max-Plus Algebra*, International Mathematical Forum. (4) (24) (2009), pp. 1157 – 1171.