

Far East Journal of Mathematical Sciences
Volume 99, Issue 1, January 2016, Pages 17-32

Far East Journal of Mathematical Sciences • [Open Access](#) • Volume 99, Issue 1, Pages 17 - 32 • January 2016

Document type
Article • [Green Open Access](#)

Source type
Journal

ISSN
09720871

DOI
10.17654/MS099010017

Publisher
Pushpa Publishing House

Original language
English

[View less](#) ^

Optimizing range norm of the image set of matrix over interval max-plus algebra

Siswanto^{a, b} ; Suparwanto, Ari^b ; Rudhito, M. Andy^c 

 Save all to author list

^a Mathematics Department, Sebelas Maret University, Indonesia

^b Mathematics Department, Gadjah Mada University, Indonesia

^c Sanata Dharma University, Indonesia

245th percentile
Citations in Scopus

0.34
FWCI 

35
Views count  

[View all metrics >](#)

Full text options  Export 

- Abstract
- Author keywords
- SciVal Topics
- Metrics

Abstract


Let \mathbb{R} be the set of real numbers and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$, $\varepsilon = -\infty$. Maxplus algebra is the set \mathbb{R}_ε that is equipped with two operations maximum and addition. It can be formed matrices in the size of $m \times n$, whose elements belong to \mathbb{R}_ε , called matrix over max-plus algebra. Let (Formula presented) and $\varepsilon = \{\varepsilon, \varepsilon\}$. The set $I(\mathbb{R})_\varepsilon$ which is equipped with two operations maximum and addition is called interval max-plus algebra. It can be formed matrices in the size $m \times n$, whose elements belong to $I(\mathbb{R})_\varepsilon$, called matrices over interval max-plus algebra. Optimizing range norm of the image set of matrix over max-plus algebra has been discussed. In this paper, we discuss optimizing range norm of the image set of matrix over interval max-plus algebra. © 2016 Pushpa Publishing House, Allahabad, India.



Author keywords

Image set; Interval max-plus algebra; Range norm

Metrics [View all metrics >](#)

2 Citations in Scopus
45th percentile

0.34 Field-Weighted Citation
Impact 

 **PlumX Metrics** 
Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 2 documents

Closure of the simple image set of linear mapping interval max-plus

Siswanto , Kurniawan, V.Y. , Pangadi (2020) *Journal of Discrete Mathematical Sciences and Cryptography*

Optimizing Range Norm of the Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components

Siswanto , Suparwanto, A. , Rudhito, M.A. (2019) *Journal of Physics: Conference Series*

[View all 2 citing documents](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

Related documents

Optimizing Range Norm of the Image Set of Matrix over Interval Max-Plus Algebra with Prescribed Components

Siswanto , Suparwanto, A. , Rudhito, M.A. (2019) *Journal of Physics: Conference Series*

Solving a tropical optimization problem via matrix sparsification

Krivulin, N. (2015) *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*



Strongly regular matrices and simple image set in interval max-plus algebra

Siswanto , Suparwanto, A. , Rudhito, M.A. (2016) *JP Journal of Algebra, Number Theory and Applications*

[View all related documents based on references](#)

Find more related documents in Scopus based on:

Authors > Keywords >

 Your CSV file was successfully exported. 

References (12)

[View in search results format >](#)

☐ All

[RIS export](#)

[Print](#)

[E-mail](#)

[Save to PDF](#)

[Create bibliography](#)

☐ 1 Akian, M., Cohen, G., Gaubert, S., Quadrat, J.P., Viot, M.
Max-plus algebra and applications to system theory and optimal control
(1994) *Proceedings of the International Congress of Mathematicians*. Cited 19 times.
Zurich, Switzerland

☐ 2 Andersen, M.H.
Max-plus algebra: Properties and applications
(2002) *A Thesis Submitted to the Department of Mathematics in Partial Fulfillment of the Requirements for the Degree of Master of Science in Mathematics*. Cited 4 times.

☐ 3 Bacelli, F., Cohen, G., Older, J.G., Quadrat, J.P.
(2001) *Synchronization and Linearity*. Cited 604 times.
John Wiley & Sons, New York

☐ 4 Butkovic, P., Tam, K.P.
On some properties of the image of a max linear mapping
(2009) *Contemp. Math*, 495, pp. 115-126. Cited 16 times.

☐ 5 Cuninghame-Green, R.A., Butkovič, P.
Bases in max-algebra [\(Open Access\)](#)

(2004) *Linear Algebra and Its Applications*, 389 (1-3), pp. 107-120. Cited 44 times.
doi: 10.1016/j.laa.2004.03.022

[View at Publisher](#)

☐ 6 Farlow, K.G.
(2009)
Max-plus algebra, Master's Thesis Submitted to the Faculty of the Virginia Polytechnic
Institute and State University in Partial Fulfillment of the Requirements for the Degree of
Masters in Mathematics

☐ 7 Konigsberg, Z.R.
A generalized eigenmode algorithm for reducible regular matrices over the max-plus
algebra
(2009) *Int. Math. Forum*, 4 (24), pp. 1157-1171. Cited 2 times.

☐ 8 Kreinovich, V.
Why intervals? Why fuzzy numbers? Towards a new justification

(2007) *Proceedings of the 2007 IEEE Symposium on Foundations of Computational
Intelligence, FOCI 2007*, art. no. 4233894, pp. 113-119. Cited 4 times.
ISBN: 1424407036; 978-142440703-3
doi: 10.1109/FOCI.2007.372156

[View at Publisher](#)

☐ 9 Rudhito, M.A.
Fuzzy number max-plus algebra and its application to fuzzy scheduling and queuing
network problems
(2011) *The Study Program S3 Mathematics Faculty of Mathematics and Natural
Sciences*. Cited 2 times.
Gadjah Mada University, Indonesian)

10

Sergeev, S.
Max-plus definite matrix closures and their eigenspaces (Open Access)

(2007) *Linear Algebra and Its Applications*, 421 (2-3 SPEC. ISS.), pp. 182-201. Cited 13 times.
doi: 10.1016/j.laa.2006.02.038

View at Publisher

11

Siswanto, Suparwanto, A., Rudhito, M.A.
Minimizing range norm of the image set of matrix over interval max-plus algebra
(2013) *Proceedings of the National Seminar on Mathematics and Applications*
Department of Mathematics, Faculty of Science and Technology, Airlangga University, Indonesian)

12

Tam, K.P.
(2010) *Optimizing and Approximating Eigenvectors in Max Algebra Thesis*. Cited 13 times.
A Thesis Submitted to The University of Birmingham for the Degree of Doctor of Philosophy (Ph.D.)


© Copyright 2015 Elsevier B.V., All rights reserved.

SciVal Topic Prominence ⓘ

Topic: Max-Plus Algebra | Tropical | Semimodule


Prominence percentile: 79.094 ⓘ

also developed by scimago:


SCIMAGO INSTITUTIONS RANKINGS

SJR


Scimago Journal & Country Rank



[Home](#)
[Journal Rankings](#)
[Country Rankings](#)
[Viz Tools](#)
[Help](#)
[About Us](#)


Far East Journal of Mathematical Sciences



Discontinued in Scopus as of 2017









COUNTRY India  Universities and research institutions in India	SUBJECT AREA AND CATEGORY Mathematics ↳ Mathematics (miscellaneous)	PUBLISHER University of Allahabad	H-INDEX 20
PUBLICATION TYPE Journals	ISSN 09720871	COVERAGE 2008-2017	INFORMATION Homepage How to publish in this journal fjms@pphmj.com

SCOPE

The Far East Journal of Mathematical Sciences (FJMS) is aimed at to provide an outlet to original research papers and survey articles of current interest in all areas of Pure and Applied Mathematics, Statistics, Theoretical Mechanics, Mathematical Physics, Theoretical Computer Science, Mathematical Biology and Financial Mathematics. Application oriented work for users of Mathematics is also encouraged. Being aware with ever-expansion of the subject, the journal remains open for upcoming newer topics as well.

 Join the conversation about this journal

 Quartiles




← Show this widget in your own website


Just copy the code below and paste within your html code:

`<a href="https://www.scimagojr.com/journalsearch.php?q=17900156722&tip=..."`



SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



ISSN 0972-0871

Far East Journal of Mathematical Sciences (FJMS)

Volume 134, Number 1, 2022



Pushpa Publishing House

Vijaya Niwas, 198 Mumfordganj
Prayagraj 211002, INDIA

<http://pphmj.com/journals/fjms.htm>
fjms@pphmj.com & arun@pphmj.com



Journal Menu

- Journal Home
- Editorial Board
- Guidelines for Authors
- Indexing
- Contents
- Subscribe
- Publication Ethics and Publication Malpractice Statement

Categories

- Computer and Communications
- Economics and Business
- Electrical, Energy and Physics
- Life Sciences
- Materials Engineering
- Mathematics
- New Publications 2022
- Statistics
- All Journals



Far East Journal of Mathematical Sciences (FJMS)

Editorial Board

Professor K. K. Azad (Editor-in-Chief)

Retired Professor & Head, Department of Mathematics, University of Allahabad, India
kkazad@pphmj.com
https://scholar.google.co.in/citations?user=r2_5P_EAAAAJ&hl=en

Research Interest

1. General Topology
2. Dynamical Systems

[ONLINE SUBMISSION](#)

Professor Ahmet Sinan Cevik

Department of Mathematics, Selcuk University, Turkey
sinan.cevik@selcuk.edu.tr ; sinan.cevik@su.edu.tr
<http://www.ahmetsinancevik.com/>

Research Interest

1. Combinatorial Group
2. Semigroup Theory
3. Algebraic and Spectral Graph Theory

[ONLINE SUBMISSION](#)

Professor Alison Marr

Department of Mathematics, Southwestern University, USA
marra@southwestern.edu
<https://www.southwestern.edu/live/profiles/25772-alison-marr>

Research Interest

1. Graph Theory
2. Graph Labeling
3. Recreational Mathematics

[ONLINE SUBMISSION](#)

Professor Ashish K. Srivastava

Department of Mathematics and Statistics, Saint Louis University, USA
ashish.srivastava@slu.edu
<https://mathstat.slu.edu/~srivastava/>

Research Interest

1. Noncommutative Ring Theory

[ONLINE SUBMISSION](#)

Professor Chelliah Selvaraj

Department of Mathematics, Periyar University, India
selvavlr@yahoo.com
<https://periyaruniversity.irins.org/profile/117061>

Research Interest

1. Particular Ring theory
2. Homological Algebra
3. Representation Theory of Algebras

[ONLINE SUBMISSION](#)

Professor Claudio Cuevas

Departamento de Matematica, Universidade Federal de Pernambuco, Brazil
cch@dmf.ufpe.br
<https://livedna.net/?dna=55.9636>

Research Interest


1. Difference Equations
2. Periodicity and Ergodicity
3. Dispersive Estimates
4. Fractional Differential Equations
5. Integral and Integro-Differential Equation

[ONLINE SUBMISSION](#)

Professor D. S. Sankar

School of Applied Sciences and Mathematics, Universiti Teknologi Brunei, Brunei Darussalam
duraisamy.sankar@utb.edu.bn
<http://www.utb.edu.bn/academics/people/professor-dr-duraisamy-sambasivam-sankar/>

Research Interest

[LATEST ISSUE](#)[SUBMIT AN ARTICLE](#)[SEARCH WITHIN JOURNALS](#)

Journal Stats

Publication count: 4091

Citation count: 10391

h10-index: 265

h-index: 32

Impact Factor: 3.22

Downloads: 1220519

Views: 4571185

Downloads/article: 298.34

Citations/article: 2.54

1. Ordinary Differential Equations
2. Partial Differential Equations
3. Mathematical Economics

[ONLINE SUBMISSION](#)

Professor E. Thandapani

Ramanujan Institute for Advanced Study in Mathematics, University of Madras, India
 ethandapani@unom.ac.in
<https://www.unom.ac.in/index.php?route=department/department/profile&deptid=48&facultyid=265>

Research Interest

1. Differential Equation (ODE & PDE)
2. Difference Equations

[ONLINE SUBMISSION](#)

Professor Haruhide Matsuda

Department of Mathematics, Shibaura Institute of Technology, Japan
 hmatsuda@sic.shibaura-it.ac.jp
<https://dl.acm.org/profile/81100347557>

Research Interest

1. Graph Theory

[ONLINE SUBMISSION](#)

Professor Hong-Xu Li

Department of Mathematics, Sichuan University, China
 hoxuli@sohu.com
<http://en.ahu.edu.cn/math/2020/1021/c6474a247132/page.htm>

Research Interest

1. Functional Differential Equations
2. Ordinary Differential Equations
3. Almost Periodicity
4. Nonlinear Functional Analysis

[ONLINE SUBMISSION](#)

Professor I. S. Rakhimov

Department of Mathematics, Universiti Teknologi MARA (UiTM), Malaysia
 isamiddin@fskm.uitm.edu.my ; isamiddin@uitm.edu.my
<https://fskm.uitm.edu.my/v5/index.php/en/corporate/staff-directory/academic-staff/39-corporate/staf-directory/academic-staff/mathematics/185-isamiddin-rakhimov?ml=1>

Research Interest

1. Structural Theory of Finite-Dimensional Algebras
2. Lie and Leibniz Algebras
3. Dialgebras
4. Commutative Algebras
5. Algebraic Geometry
6. Invariant Theory

[ONLINE SUBMISSION](#)

Professor Jin-Lin Liu

Department of Mathematics, Yangzhou University, China
 jlliu@yzu.edu.cn
<https://dl.acm.org/profile/81458650741>

Research Interest

1. Geometric Function Theory

[ONLINE SUBMISSION](#)

Professor Lingyun Gao

Department of Mathematics, Jinan University, China
 tgaoly@jnu.edu.cn

Research Interest

1. Complex Analysis
2. Complex Differential (difference) Equations

[ONLINE SUBMISSION](#)

Professor Manabu Hagiwara

Department of Mathematics, University of Chiba, Japan
 hagiwara@math.s.chiba-u.ac.jp
<http://www.math.s.chiba-u.ac.jp/~hagiwara/>

Research Interest

1. Mathematical Approach to Coding Theory
2. Information Theory
3. Combinatorics, and its application to industry

[ONLINE SUBMISSION](#)

Professor Pu Zhang

Department of Mathematics, Shanghai Jiao Tong University, China
 pzhang@sjtu.edu.cn
<https://en.zhiyuan.sjtu.edu.cn/en/faculty/43/detail>

Research Interest

1. Algebra Representation Theory
2. Quantum Groups

[ONLINE SUBMISSION](#)
Professor Qing-Wen Wang

Department of Mathematics, Shanghai University, China
 wqw@staff.shu.edu.cn
<https://www.scicol.shu.edu.cn/PDF/WangQingWen.htm>

Research Interest

1. Linear Algebra
2. Matrix Theory
3. Operator Algebra

[ONLINE SUBMISSION](#)
Professor Sahsene Altinkaya

Department of Mathematics, Beykent University, Turkey
 sahsenealtinkaya@beykent.edu.tr
<https://avesis.uludag.edu.tr/sahsene>

Research Interest

1. Geometric Function Theory
2. Bi-univalent Functions

[ONLINE SUBMISSION](#)
Professor Satoru Takagi

Mathematics Department, Waseda University, Japan
 satoru@waseda.jp
<http://www.f.waseda.jp/satoru/index.html>

Research Interest

1. Partial Differential Equations
2. Functional Analysis

[ONLINE SUBMISSION](#)
Professor Sourav Das

Department of Mathematics, National Institute of Technology, Jamshedpur, India
 sourav.das@mt.iitr.ac.in
https://www.iitr.ac.in/~MT/Sourav_Das

Research Interest

1. Complex Analysis
2. Special Functions
3. Orthogonal Polynomials
4. Inequalities
5. Continued Fractions
6. Asymptotics

[ONLINE SUBMISSION](#)
Professor Vladimir Tulovsky

Department of Mathematics and Computer Science, Saint John's University, USA
 TULOVSKEV@stjohns.edu
<https://www.stjohns.edu/academics/faculty/vladimir-tulovsky>

Research Interest

1. Partial Differential Equations

[ONLINE SUBMISSION](#)
Professor Wei Dong Gao

Center for Applied Mathematics, Tianjin University, China
 weidong.gao@tju.edu.cn
<http://cam.tju.edu.cn/en/faculty/index.php?id=92>

Research Interest

1. Combinatorial Number Theory

[ONLINE SUBMISSION](#)
Professor Young Joo Lee

Department of Mathematics, Chonnam National University, South Korea
 leeyj@chonnam.ac.kr
<https://www.researchgate.net/profile/Young-Lee-84>

Research Interest

1. Complex analysis

2. Operator theory

[ONLINE SUBMISSION](#)**Professor Zhenlu Cui**

Department of Mathematics and Computer Science, Fayetteville State University, USA

zcui@uncfsu.edu

<https://www.uncfsu.edu/academics/colleges-schools-and-departments/college-of-health-science-and-technology/department-of-mathematics-and-computer-science/math-and-cs-faculty-and-staff/dr-zhenlu-cui>

Research Interest

1. Multiscale Modeling of Soft Matters
2. Perturbation Analysis
3. Numerical Analysis
4. Approximation Theory

[ONLINE SUBMISSION](#)

P-ISSN: 0972-0871

ISSN 0972-0871

**Far East
Journal
of
Mathematical
Sciences
(FJMS)**

Volume 134, Number 1, 2022

Pushpa Publishing House
Vijaya Vihar, 28B Bhamburda Road
Pune-411 004, India
http://www.ppshmj.com
ppshmj@gmail.com

LATEST ISSUE

SUBMIT AN ARTICLE

SEARCH WITHIN JOURNALS

Journal Stats

Publication count: 4091

Citation count: 10391

h10-index: 265

h-index: 32

Impact Factor: 3.22

Downloads: 1220519

Views: 4571185

Downloads/article: 298.34

Citations/article: 2.54

BACK

Far East Journal of Mathematical Sciences (FJMS)

Volume 99, Issue 1, Pages 1 - 152 (January 2016)

Articles 1 - 8 of 8 [1]

THE FRACTIONAL DIFFERENTIAL TRANSFORMATION METHOD FOR THE FRACTIONAL STURM-LIOUVILLE PROBLEM FOR FRACTIONAL BESSEL EQUATION

by: Sutrima, Marifatun and Isnandar Slamet

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 1 - 15 \(January 2016\)](#)

Downloads: 374 Views: 814

OPTIMIZING RANGE NORM OF THE IMAGE SET OF MATRIX OVER INTERVAL MAX-PLUS ALGEBRA

by: Siswanto, Ari Suparwanto and M. Andy Rudhito

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 17 - 32 \(January 2016\)](#)

Downloads: 395 Views: 1291

RIESZ BASIS PROPERTY AND EXPONENTIAL STABILITY OF A FLEXIBLE EULER-BERNOULLI BEAMS WITH A FORCE CONTROL IN ROTATION AND VELOCITY ROTATION

by: TourÃ© K. Augustin, Koua B. Jean Claude and Diop Fatou N.

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 33 - 60 \(January 2016\)](#)

Downloads: 376 Views: 1200

A TYPE OF CONFORMAL CURVATURE TENSOR

by: Jaeman Kim

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 61 - 74 \(January 2016\)](#)

Downloads: 399 Views: 1385

A CERTAIN FRACTIONAL DERIVATIVE OPERATOR FOR p-VALENT FUNCTIONS AND NEW CLASS OF ANALYTIC FUNCTIONS WITH NEGATIVE COEFFICIENTS

by: A. A. Amourah, Feras Yousef, Tariq Al-Hawary and M. Darus

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 75 - 87 \(January 2016\)](#)

Downloads: 400 Views: 1252

ON CERTAIN SUBCLASS OF STARLIKE ANALYTIC FUNCTIONS ASSOCIATED WITH RUSCHEWEYH DERIVATIVE AND A GENERALIZED MULTIPLIER TRANSFORMATION

by: Anessa Oshah and Maslina Darus

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 89 - 108 \(January 2016\)](#)

Downloads: 391 Views: 1137

A PROBABILISTIC INTEGRATED VENDOR-BUYER COOPERATIVE INVENTORY MODEL WITH IMPERFECT QUALITY ITEMS, CONTROLLABLE LEAD TIME, UNCERTAINTY DEMANDS, INSPECTION ERROR, SHORTAGE AND BACKORDERING ALLOWED

by: Rubono Setiawan and Triyanto

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 109 - 132 \(January 2016\)](#)

Downloads: 381 Views: 1318

SOME CONVERGENCE RESULTS FOR MODIFIED ABBAS AND NAZIR ITERATION PROCESS ON A NONLINEAR DOMAIN

by: Duangkamon Kitkuan

Far East Journal of Mathematical Sciences (FJMS)

[Volume 99, Issue 1, Pages 133 - 152 \(January 2016\)](#)

Downloads: 380 Views: 911

Journal Menu

- Journal Home
- Editorial Board
- Guidelines for Authors
- Indexing
- Contents
- Subscribe
- Publication Ethics and Publication Malpractice Statement

Content

- Volume 140 (2023)
- Volume 139 (2022)
- Volume 138 (2022)
- Volume 137 (2022)
- Volume 136 (2022)
- Volume 135 (2022)
- Volume 134 (2022)
- Volume 133 (2021)
- Volume 132 (2021)
- Volume 131 (2021)
- Volume 130 (2021)
- Volume 129 (2021)
- Volume 128 (2021)
- Volume 127 (2020)
- Volume 126 (2020)
- Volume 125 (2020)
- Volume 124 (2020)
- Volume 123 (2020)
- Volume 122 (2020)
- Volume 121 (2019)
- Volume 120 (2019)
- Volume 119 (2019)
- Volume 118 (2019)
- Volume 117 (2019)
- Volume 116 (2019)
- Volume 115 (2019)
- Volume 114 (2019)
- Volume 113 (2019)
- Volume 112 (2019)
- Volume 111 (2019)
- Volume 110 (2019)
- Volume 109 (2018)
- Volume 108 (2018)
- Volume 107 (2018)
- Volume 106 (2018)
- Volume 105 (2018)
- Volume 104 (2018)
- Volume 103 (2018)
- Volume 102 (2017)
- Volume 101 (2017)
- Volume 100 (2016)
- Volume 99 (2016)
- Volume 99, Issue 11
- Pg 1603 - 1778 (June 2016)
- Volume 99, Issue 12
- Pg 1779 - 1970 (June 2016)
- Volume 99, Issue 9
- Pg 1283 - 1441 (May 2016)
- Volume 99, Issue 10
- Pg 1443 - 1601 (May 2016)
- Volume 99, Issue 7
- Pg 959 - 1117 (April 2016)
- Volume 99, Issue 8
- Pg 1119 - 1281 (April 2016)
- Volume 99, Issue 5
- Pg 615 - 774 (March 2016)
- Volume 99, Issue 6
- Pg 775 - 958 (March 2016)



OPTIMIZING RANGE NORM OF THE IMAGE SET OF MATRIX OVER INTERVAL MAX-PLUS ALGEBRA

Siswanto^{1,2}, Ari Suparwanto² and M. Andy Rudhito³

¹Mathematics Department

Faculty of Mathematics and Natural Science

Sebelas Maret University

Indonesia

e-mail: sis.mipauns@yahoo.co.id

²Mathematics Department

Faculty of Mathematics and Natural Science

Gadjah Mada University

Indonesia

e-mail: ari_suparwanto@yahoo.com

³The Study Program in Mathematics Education

Faculty of Teacher Training and Education

Sanata Dharma University

Indonesia

e-mail: arudhito@yahoo.co.id

Abstract

Let \mathbb{R} be the set of real numbers and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$, $\varepsilon = -\infty$. Max-plus algebra is the set \mathbb{R}_ε that is equipped with two operations maximum and addition. It can be formed matrices in the size of $m \times n$,

Received: June 8, 2015; Revised: August 30, 2015; Accepted: September 16, 2015

2010 Mathematics Subject Classification: 15A80.

Keywords and phrases: interval max-plus algebra, range norm, image set.

Communicated by K. K. Azad

whose elements belong to \mathbb{R}_ε , called matrix over max-plus algebra.

Let $I(\mathbb{R})_\varepsilon = \{x = [\underline{x}, \bar{x}] \mid \underline{x}, \bar{x} \in \mathbb{R}, \varepsilon < \underline{x} \leq \bar{x}\} \cup \{\varepsilon\}$ and $\varepsilon = \{\varepsilon, \varepsilon\}$.

The set $I(\mathbb{R})_\varepsilon$ which is equipped with two operations maximum and addition is called interval max-plus algebra. It can be formed matrices in the size $m \times n$, whose elements belong to $I(\mathbb{R})_\varepsilon$, called matrices over interval max-plus algebra. Optimizing range norm of the image set of matrix over max-plus algebra has been discussed. In this paper, we discuss optimizing range norm of the image set of matrix over interval max-plus algebra.

I. Introduction

Max-plus algebra is the set $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$, of which \mathbb{R} are all the sets of real numbers and $\varepsilon = -\infty$ that is equipped with two operations \oplus (maximum) and \otimes (addition). Max-plus algebra has been used to model and analyze problems in planning, communication, production system, queueing system with finite capacity, parallel computation, and traffic using algebra [3] and also several other applications, i.e., in flight scheduling and airport problems [2] whereas min-plus algebra is the set $\mathbb{R}_{\varepsilon'} = \mathbb{R} \cup \{\varepsilon'\}$ and $\varepsilon' = \infty$ that is equipped with two operations \oplus' (minimum) and \otimes' (addition) [12]. Tam [12] also discussed complete max-plus algebra, of which the set $\overline{\mathbb{R}}_\varepsilon = \mathbb{R} \cup \{\varepsilon, \varepsilon'\}$ is equipped with operations \oplus and \otimes , and complete min-plus algebra, of which the set $\overline{\mathbb{R}}_{\varepsilon'} = \mathbb{R} \cup \{\varepsilon, \varepsilon'\}$ is equipped with two operations \oplus' and \otimes' . Furthermore, $\overline{\mathbb{R}}_\varepsilon$ and $\overline{\mathbb{R}}_{\varepsilon'}$ are written as $\overline{\mathbb{R}}$.

It can be formed a set of matrices in the size $m \times n$ of the sets \mathbb{R}_ε , $\mathbb{R}_{\varepsilon'}$, and $\overline{\mathbb{R}}$. The set of matrices with components in \mathbb{R}_ε is denoted by $\mathbb{R}_\varepsilon^{m \times n}$. The matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$ is called *matrix over max-plus algebra*. For $n = 1$, a set of vectors over max-plus algebra is obtained and written as \mathbb{R}_ε^m , that is $\mathbb{R}_\varepsilon^m = \{(x_1, x_2, \dots, x_m)^T \mid x_1, x_2, \dots, x_m \in \mathbb{R}_\varepsilon\}$ [5, 6]. Moreover, if $m = n$, then the set $\mathbb{R}_\varepsilon^{n \times n}$ equipped with operations \oplus and \otimes forms idempotent semiring [1, 7, 10].

Tam [12] illustrated an example of application of max-plus algebra in the production system. For example $A = (A_{ij}) \in \mathbb{R}_\varepsilon^{n \times n}$ is a production matrix, where A_{ij} shows the period of time of the production process from machine M_j to M_i , while $x(k) = (x_i(k)) \in \mathbb{R}_\varepsilon^n$ is a vector, where $x_i(k)$ is the starting time of machine M_i at the k th stage. In this production process, we have the equation $x(k+1) = A \otimes x(k)$. One of the criteria used by manufacturer is that the production process is expected to be continue periodically in certain periods, e.g., λ , so that the equation $x(k+1) = \lambda \otimes x(k)$ is obtained. From $x(k+1) = A \otimes x(k)$ and $x(k+1) = \lambda \otimes x(k)$, we get $A \otimes x(k) = \lambda \otimes x(k)$. The eigenvalue and eigenvector problem of matrix A is to determine the eigenvalue λ and the eigenvector $x(k)$ which satisfy the equation $A \otimes x(k) = \lambda \otimes x(k)$.

In the real life situation, there are several ways for the manufacturer to determine the starting time of each machine. One of the ways is by choosing eigenvector as the starting time, so that the system will immediately reach the steady state; that is the process continues periodically with the eigenvalue as its periods. However, in reality, there can exist more than one independent eigenvectors for the manufacturer to choose. In that case, a set of linear combination of the independent eigenvectors is constructed, and as such, the manufacturer needs an additional criterion to choose one element of the set. The additional criterion is by considering the difference between the largest and the smallest of the starting time of each machine. The difference is expressed as range norm of the starting time of each machine. Manufacturer can optimize (either minimize or maximize) the range norm of the starting time of each machine [4, 12]. The readiness of the raw material, the availability of the resources and the distribution of the product become the determining factors for the manufacturer to optimize the range norm of the starting time of each machine. In solving the problems of that production system, Tam [12] has discussed optimizing range norm of the image set of matrix over max-plus algebra.

Kreinovich [8] stated that any measurement is never 100% accurate. Thus, the value of the measurement \tilde{x} in general differs from the real value x . Specifically, in the case of interval uncertainty, after it is being measured and the result of the measurement \tilde{x} is obtained, the acquired information that the real value x is contained in the interval $x = [\tilde{x} - \Delta, \tilde{x} + \Delta]$.

Based on the above statement, the estimation on the period of time of a certain process can be given, for example the period of time of the process in a production system. Therefore, the period of time of a certain production process can be given in an interval of time.

According to the Kreinovich's idea [8] about the probability to give an estimation of the period of time of a certain process and Rudhito [9] about max-plus algebra generalization, i.e., interval max-plus algebra, we try to expand the concept in max-plus algebra, i.e., the optimizing range norm of the image set of matrix over interval max-plus algebra. Siswanto et al. [11] have investigated how to minimize range norm of the image set of matrix over interval max-plus algebra. In this research, we develop the range norm definition in [11]. Using the new range norm definition, we investigate the way to minimize and maximize range norm of the image set of matrix over interval max-plus algebra.

Before discussing the result of this paper, several concepts which support the discussion are observed as follows [12]:

Definition 1.1. Given that $a \in \mathbb{R}$, the conjugate of a is $a^* = a^{-1} = -a$. Suppose that $A \in \overline{\mathbb{R}}^{m \times n}$. Then the conjugate of matrix A is $A^* = (a_{ji}^*)$ or $A^* = -A^T$.

Definition 1.2. Given that $A \in \overline{\mathbb{R}}^{m \times n}$ and if matrix A at least has one finite element in each row, then matrix A is called *row \mathbb{R} -astic*. If matrix A at least has one finite element in each column, then matrix A is called *column \mathbb{R} -astic*. If matrix A at least has one finite element in each row and each column, then matrix A is called *double \mathbb{R} -astic*.

Definition 1.3. Suppose that $A \in \mathbb{R}_\varepsilon^{m \times n}$, $\text{Im}(A)$ is defined as $\text{Im}(A) = \{A \otimes x \mid x \in \mathbb{R}_\varepsilon^n\}$, namely the image set of matrix A .

Moreover, the definition of the range norm and the problem of optimizing range norm of the image set of matrix over max-plus algebra are given. The notations of $M = \{1, 2, \dots, m\}$ and $N = \{1, 2, \dots, n\}$ are used to simplify.

Definition 1.4. If $x \in \mathbb{R}^m$, then the function $\delta(x) = \sum_{i \in M}^\oplus x_i - \sum_{i \in M}^{\oplus'} x_i$ is called the *range norm* of x . In other words, the range norm of x = the largest value in x - the smallest value in x .

In this case, the norm term is a mere identification, in contrast to the norm defined in inner product space in general.

Problem 1.1. Given that matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve:

$$\begin{aligned} &\text{minimize } \delta(b) \\ &\text{subject to } b \in \text{Im}(A). \end{aligned}$$

Definition 1.5. If $x \in \mathbb{R}_\varepsilon^m$, then the function $\tilde{\delta}(x) = \sum_{\substack{i \in M \\ x_i \neq \varepsilon}}^\oplus x_i - \sum_{\substack{i \in M \\ x_i \neq \varepsilon}}^{\oplus'} x_i$

is the range norm of x , considering only the finite component of x .

Problem 1.2. Given that matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve:

$$\begin{aligned} &\text{minimize } \tilde{\delta}(b) \\ &\text{subject to } b \in \text{Im}(A). \end{aligned}$$

Problem 1.3. Given that matrix $A \in \mathbb{R}_\varepsilon^{m \times n}$, solve:

$$\begin{aligned} &\text{maximize } \delta(b) \\ &\text{subject to } b \in \text{Im}(A). \end{aligned}$$

Other than the above concepts, the concepts on interval max-plus algebra is also needed [9].

Closed interval x in \mathbb{R}_ε is a subset of \mathbb{R}_ε in the form $x = [\underline{x}, \bar{x}] = \{x \in \mathbb{R}_\varepsilon \mid \underline{x} \leq x \leq \bar{x}\}$. The interval x in \mathbb{R}_ε is known as interval max-plus. A number $x \in \mathbb{R}_\varepsilon$ can be expressed as interval $[x, x]$.

Definition 1.6. $I(\mathbb{R})_\varepsilon = \{x = [\underline{x}, \bar{x}] \mid \underline{x}, \bar{x} \in \mathbb{R}, \varepsilon < \underline{x} \leq \bar{x}\} \cup \{\varepsilon\}$ is formed with $\varepsilon = [\varepsilon, \varepsilon]$. On the set $I(\mathbb{R})_\varepsilon$, the operations \oplus and \otimes with $x \oplus y = [\underline{x} \oplus \underline{y}, \bar{x} \oplus \bar{y}]$ and $x \otimes y = [\underline{x} \otimes \underline{y}, \bar{x} \otimes \bar{y}]$ for every $x, y \in I(\mathbb{R})_\varepsilon$ are defined. Furthermore they are known as interval max-plus algebra and notated as $I(\mathfrak{R})_{\max} = (I(\mathbb{R})_\varepsilon; \oplus, \otimes)$.

Definition 1.7. The set of matrices in the size $m \times n$ with the elements in $I(\mathbb{R})_\varepsilon$ is notated as $I(\mathbb{R})_\varepsilon^{m \times n}$, namely:

$$I(\mathbb{R})_\varepsilon^{m \times n} = \{A = [A_{ij}] \mid A_{ij} \in I(\mathbb{R})_\varepsilon; \text{ for } i = 1, 2, \dots, m \text{ and } j = 1, 2, \dots, n\}.$$

Matrices which belong to $I(\mathbb{R})_\varepsilon^{m \times n}$ are known as matrices over interval max-plus algebra or shortly interval matrices.

Definition 1.8. For $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, matrices $\underline{A} = [\underline{A}_{ij}] \in \mathbb{R}_\varepsilon^{m \times n}$ and $\bar{A} = [\bar{A}_{ij}] \in \mathbb{R}_\varepsilon^{m \times n}$ are defined, each is known as lower bound matrix and upper bound matrix of the interval matrix A .

Definition 1.9. Given that interval matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, of which \underline{A} and \bar{A} each act as lower bound matrix and upper bound matrix of the interval matrix A . The matrix interval of A is defined, namely $[\underline{A}, \bar{A}] = \{A \in \mathbb{R}_\varepsilon^{m \times n} \mid \underline{A} \leq A \leq \bar{A}\}$ and $I(\mathbb{R}_\varepsilon^{m \times n})_b = \{[\underline{A}, \bar{A}] \mid A \in I(\mathbb{R})_\varepsilon^{m \times n}\}$.

The semimodule $I(\mathbb{R})_\varepsilon^{n \times n}$ over $I(\mathbb{R})_\varepsilon$ is isomorphic to the semimodule $I(\mathbb{R}_\varepsilon^{n \times n})_b$ over $I(\mathbb{R})_\varepsilon$ with the mapping of $f : I(\mathbb{R})_\varepsilon^{n \times n} \rightarrow I(\mathbb{R}_\varepsilon^{n \times n})_b$, $f(A) = [\underline{A}, \bar{A}]$, $\forall A \in I(\mathbb{R})_\varepsilon^{n \times n}$. The matrix interval $[\underline{A}, \bar{A}] \in I(\mathbb{R}_\varepsilon^{n \times n})_b$ is

known as matrix interval which corresponds to interval matrix $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$ and which is represented with $A \approx [\underline{A}, \overline{A}]$.

Definition 1.10. It is defined that

$$I(\mathbb{R})_{\varepsilon}^n = \{x = [x_1, x_2, \dots, x_n]^T \mid x_i \in I(\mathbb{R})_{\varepsilon}; i = 1, 2, \dots, n\}.$$

Set $I(\mathbb{R})_{\varepsilon}^n$ can be considered as set $I(\mathbb{R})_{\varepsilon}^{n \times 1}$. The elements of $I(\mathbb{R})_{\varepsilon}^n$ are called *interval vector* over $I(\mathbb{R})_{\varepsilon}$. Interval vector x corresponds to vector interval $[\underline{x}, \overline{x}]$, i.e., $x \approx [\underline{x}, \overline{x}]$.

The concept of interval min-plus algebra is defined in the same way with interval max-plus algebra concepts. Following are the definitions of complete interval max-plus algebra and complete interval min-plus algebra [11].

Definition 1.11. Complete interval max-plus algebra is set $I(\overline{\mathbb{R}})_{\varepsilon} = I(\mathbb{R})_{\varepsilon} \cup \{\varepsilon'\}$, $\varepsilon' = [\varepsilon', \varepsilon']$ that is completed with two operations $\overline{\oplus}$ and $\overline{\otimes}$, meanwhile complete interval min-plus algebra is set $I(\overline{\mathbb{R}})_{\varepsilon'} = I(\mathbb{R})_{\varepsilon'} \cup \{\varepsilon\}$ with the operations $\overline{\oplus}'$ and $\overline{\otimes}'$. Furthermore, $I(\overline{\mathbb{R}})_{\varepsilon} = I(\mathbb{R})_{\varepsilon'}$ are written as $I(\overline{\mathbb{R}})$.

In the same way as in the interval max-plus algebra and interval min-plus algebra, in complete interval max-plus algebra and complete interval min-plus algebra, can be defined as the set of matrices in the size $m \times n$ are notated as $I(\overline{\mathbb{R}})^{m \times n}$.

II. Main Results

In this section, the optimizing range norm of the image set of matrix over interval max-plus algebra is presented:

A. Minimizing the range norm of matrix image set over interval max-plus algebra

The following are definitions of the range norm of vector and the image set of matrix over interval max-plus algebra.

Definition 2.1. Suppose that $x \in I(\mathbb{R})^m$ with $x \approx [\underline{x}, \bar{x}]$, $\underline{x}, \bar{x} \in \mathbb{R}^m$. The function $I\delta(x) = [\min(\delta(\underline{x}), \delta(\bar{x})), \delta(\bar{x})]$ is called the *range norm* of x . In other words, the range norm of x is interval $[\min(\delta(\underline{x}), \delta(\bar{x})), \delta(\bar{x})]$, where $\min(\delta(\underline{x}), \delta(\bar{x})) = \text{minimum of } \delta(\underline{x}) \text{ or } \delta(\bar{x})$.

Definition 2.2. Given that matrix $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$, $\underline{A}, \bar{A} \in \mathbb{R}_{\varepsilon}^{m \times n}$, it is then defined that $\text{Im}(A) = \{A \bar{\otimes} p \mid p \in I(\mathbb{R})_{\varepsilon}^n\}$.

Based on the above definitions, the problem of minimizing the range norm of matrix image set can be formulated as follows:

Problem 2.1. Given that matrix $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$, solve:

$$\begin{aligned} & \text{minimize } I\delta(b) \\ & \text{subject to } b \in \text{Im}(A). \end{aligned}$$

Definition 2.3. Given that matrix $b \in I(\mathbb{R})_{\varepsilon}^m$ with $b \approx [\underline{b}, \bar{b}]$. The function $I\delta(b)$ is said *minimum* if and only if $\delta(\underline{b})$ and $\delta(\bar{b})$ are minimum.

If the image vector has infinite components, the definition about the range norm and the problem of minimizing the range norm of matrix image set can be formulated as follows:

Definition 2.4. Suppose that $x \in I(\mathbb{R})_{\varepsilon}^m$ with $x \approx [\underline{x}, \bar{x}]$. The function $\bar{I}\delta(x) = [\min(\tilde{\delta}(\underline{x}), \tilde{\delta}(\bar{x})), \tilde{\delta}(\bar{x})]$ is the range norm of x , after only considering the finite component.

Problem 2.2. Given that matrix $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$, solve:

$$\begin{aligned} & \text{minimize } \bar{I}\delta(b) \\ & \text{subject to } b \in \text{Im}(A). \end{aligned}$$

Definition 2.5. Given that matrix $b \in I(\mathbb{R})_{\varepsilon}^m$ with $b \approx [\underline{b}, \bar{b}]$. The function $\bar{I}\delta(b)$ is said *minimum* if and only if $\tilde{\delta}(\underline{b})$ and $\tilde{\delta}(\bar{b})$ are minimum.

Minimizing the range norm if its image vector is finite

The first we investigate minimizing the range norm if its image vector is finite, i.e., Problem 2.1. The next lemmas, definition and theorems are needed to answer Problem 2.1:

Lemma 2.6. *Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$. If $A = [A_{ij}]$ with $A_{ij} \neq [\varepsilon, \varepsilon]$ for each $i \in M$, $j \in N$; $b \approx [\underline{b}, \bar{b}]$ and $0 \approx [\underline{0}, \bar{0}]$ with $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{0})$ and $\bar{b} = \bar{A} \otimes (\bar{A}^* \otimes' \bar{0})$, then*

- (i) $b \leq 0$ and
- (ii) $b_i = 0$ for some $i \in M$.

Proof. Suppose that $b \approx [\underline{b}, \bar{b}]$ with $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{0})$ and $\bar{b} = \bar{A} \otimes (\bar{A}^* \otimes' \bar{0})$. Since $A = [A_{ij}]$ with $A_{ij} \neq [\varepsilon, \varepsilon]$ for each $i \in M$ and $j \in N$, $\underline{A} = [\underline{A}_{ij}]$ with $\underline{A}_{ij} \neq \varepsilon$ for each $i \in M$ and $j \in N$ and $\bar{A} = [\bar{A}_{ij}]$ with $\bar{A}_{ij} \neq \varepsilon$ for each $i \in M$ and $j \in N$. According to concept in max-plus algebra,

- (i) $\underline{b} \leq \underline{0}$ and $\bar{b} \leq \bar{0}$,
- (ii) $\underline{b}_i = \underline{0}$ and $\bar{b}_i = \bar{0}$ for some $i \in M$.

Since $b \approx [\underline{b}, \bar{b}]$ and $0 \approx [\underline{0}, \bar{0}]$,

- (i) $b \leq 0$,
- (ii) $b_i = 0$ for some $i \in M$. □

Lemma 2.7. *If $x, y \in I(\mathbb{R})^m$ and $a \in I(\mathbb{R})$ with $x \approx [\underline{x}, \bar{x}]$, $y \approx [\underline{y}, \bar{y}]$ and $a \approx [\underline{a}, \bar{a}]$, then*

- (i) $I\delta(x \oplus y) \leq I\delta(x) \oplus I\delta(y)$,
- (ii) $I\delta(x) = I\delta(a \otimes x)$.

Proof. Since $x \approx [\underline{x}, \bar{x}]$, $y \approx [\underline{y}, \bar{y}]$ and $a \approx [\underline{a}, \bar{a}]$, $x \oplus y \approx [\underline{x} \oplus \underline{y}, \bar{x} \oplus \bar{y}]$ and $a \otimes x \approx [\underline{a} \otimes \underline{x}, \bar{a} \otimes \bar{x}]$. Next, according to concept in max-plus algebra,

$$(i) \quad \delta(\underline{x} \oplus \underline{y}) \leq \delta(\underline{x}) \oplus \delta(\underline{y}) \text{ and } \delta(\bar{x} \oplus \bar{y}) \leq \delta(\bar{x}) \oplus \delta(\bar{y}),$$

$$(ii) \quad \delta(\underline{x}) = \delta(\underline{a} \otimes \underline{x}) \text{ and } \delta(\bar{x}) = \delta(\bar{a} \otimes \bar{x}).$$

Therefore,

(i)

$$\begin{aligned} I\delta(x \oplus y) &= [\min(\delta(\underline{x} \oplus \underline{y}), \delta(\bar{x} \oplus \bar{y})), \delta(\bar{x} \oplus \bar{y})] \\ &\leq [\min(\delta(\underline{x}) \oplus \delta(\underline{y}), \delta(\bar{x}) \oplus \delta(\bar{y})), \delta(\bar{x}) \oplus \delta(\bar{y})] \\ &= [\min(\delta(\underline{x}), \delta(\bar{x})), \delta(\bar{x})] \oplus [\min(\delta(\underline{y}), \delta(\bar{y})), \delta(\bar{y})] \\ &= I\delta(x) \oplus I\delta(y), \end{aligned}$$

(ii)

$$\begin{aligned} I\delta(x) &= [\min(\delta(\underline{x}), \delta(\bar{x})), \delta(\bar{x})] \\ &= [\min(\delta(\underline{a} \otimes \underline{x}), \delta(\bar{a} \otimes \bar{x})), \delta(\bar{a} \otimes \bar{x})] = I\delta(a \otimes x). \quad \square \end{aligned}$$

Definition 2.8. The interval matrix $A \in I(\mathbb{R})^{m \times n}$, $A \approx [\underline{A}, \bar{A}]$ is considered as double $I(\mathbb{R})$ -astic if A is double \mathbb{R} -astic for each $A \in [\underline{A}, \bar{A}]$.

Theorem 2.9. The matrix $A \in I(\mathbb{R})^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$ is double $I(\mathbb{R})$ -astic if and only if \underline{A} is double \mathbb{R} -astic.

Proof. It is known that $A \in I(\mathbb{R})^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$ is double $I(\mathbb{R})$ -astic. Based on the definition, A is double \mathbb{R} -astic for each $A \in [\underline{A}, \bar{A}]$. If $A = \underline{A}$, then \underline{A} is double \mathbb{R} -astic. It is known that \underline{A} is double \mathbb{R} -astic. Since \underline{A} is the lower bound matrix for the matrix interval $[\underline{A}, \bar{A}]$, $\underline{A} \leq \bar{A}$. Therefore, since it is a matrix \underline{A} double \mathbb{R} -astic, matrix A

double \mathbb{R} -astic for each $A \in [\underline{A}, \bar{A}]$, thus $A \in I(\mathbb{R})^{m \times n}$ double $I(\mathbb{R})$ -astic is obtained. \square

The following theorem presents solution to Problem 2.1:

Theorem 2.10. *Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ is double $I(\mathbb{R})$ -astic and $v_a \in I(\mathbb{R})^m$ is a vector whose each component is equal to $a \in I(\mathbb{R})$. Then $b \approx [\underline{b}, \bar{b}]$ is a solution for Problem 2.1 such that $\underline{b} \leq \underline{A} \otimes (\underline{A}^* \otimes' \underline{v}_a)$ and $\bar{b} = \bar{A} \otimes (\bar{A}^* \otimes' \bar{v}_a)$.*

Proof. Suppose that $A \approx [\underline{A}, \bar{A}]$, $v_a \approx [\underline{v}_a, \bar{v}_a]$ and $a \approx [\underline{a}, \bar{a}]$. Since $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ is double $I(\mathbb{R})$ -astic, $v_a \in I(\mathbb{R})^m$ and v_a is a vector whose each component is equal to $a \in I(\mathbb{R})$, which means that:

(i) Matrix $\underline{A} \in \mathbb{R}_\varepsilon^{m \times n}$ is double \mathbb{R} -astic and \underline{v}_a is a vector whose component is equal to $\underline{a} \in \mathbb{R}$. Based on concept in max-plus algebra, $\underline{A} \otimes (\underline{A}^* \otimes' \underline{v}_a)$ is a solution for Problem 1.1.

(ii) Matrix $\bar{A} \in \mathbb{R}_\varepsilon^{m \times n}$ is double \mathbb{R} -astic and \bar{v}_a is a vector whose component is equal to $\bar{a} \in \mathbb{R}$. Based on concept in max-plus algebra, $\bar{A} \otimes (\bar{A}^* \otimes' \bar{v}_a)$ is a solution for Problem 1.1.

Therefore, $[b, \bar{b}]$ with $\underline{b} \leq \underline{A} \otimes (\underline{A}^* \otimes' \underline{v}_a)$ and $\bar{b} = \bar{A} \otimes (\bar{A}^* \otimes' \bar{v}_a)$ is a solution for Problem 2.1. \square

Minimizing the range norm if its image vector is not finite

To solve Problem 2.2, the following theorem is used:

Theorem 2.11. *Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, where $A = [A_{ij}]$ for each $i \in M$ and $j \in N$ with $A_{ij} \neq [\varepsilon, \varepsilon]$, and $v_a \in I(\mathbb{R})^m$ is a vector whose each component is equal to constant interval a . If $x \in I(\mathbb{R})_\varepsilon^n$ such that $b' \leq$*

$A \otimes x$ is a solution for Problem 2.2, then $\forall j \in N$ satisfies:

a. $x_j = [\varepsilon, \varepsilon]$ or

b. $x_j = [\min((\underline{A}^* \otimes' \underline{v}_a)_j, (\bar{A}^* \otimes' \bar{v}_a)_j), (\bar{A}^* \otimes' \bar{v}_a)_j]_j$.

Proof. Suppose that $A \approx [\underline{A}, \bar{A}]$, $v_a \approx [\underline{v}_a, \bar{v}_a]$ and $a \approx [\underline{a}, \bar{a}]$. Since $A = [A_{ij}]$ for each $i \in M$ and $j \in N$ with $A_{ij} \neq [\varepsilon, \varepsilon]$, and $v_a \in I(\mathbb{R})^m$ is a vector whose each component is equal to constant interval a , it means that:

(i) Matrix $\underline{A} = [\underline{A}_{ij}]$ for each $i \in M$ and $j \in N$ with $\underline{A}_{ij} \neq \varepsilon$ and $\underline{v}_a \in \mathbb{R}^m$ is a vector whose each component is equal to constant \underline{a} .

(ii) Matrix $\bar{A} = [\bar{A}_{ij}]$ for each $i \in M$ and $j \in N$ with $\bar{A}_{ij} \neq \varepsilon$ and $\bar{v}_a \in \mathbb{R}^m$ is a vector whose each component is equal to constant \bar{a} .

Then, suppose that $x \approx [\underline{x}, \bar{x}]$, $b' \approx [\underline{b}', \bar{b}']$ and $x_j \approx [\underline{x}_j, \bar{x}_j]$. It is known that $x \in I(\mathbb{R})_{\varepsilon}^n$ such that $b' = A \otimes x$ is a solution for Problem 2.2, which according to concept in max-plus algebra means:

(iii) $\underline{x} \in \mathbb{R}_{\varepsilon}^n$ with $\underline{b}' \leq \underline{A} \otimes \underline{x}$ is the solution for Problem 1.2, so $\forall j \in N$ satisfies:

a'. $\underline{x}_j = \varepsilon$ or

b'. $\underline{x}_j = \min((\underline{A}^* \otimes' \underline{v}_a)_j, (\bar{A}^* \otimes' \bar{v}_a)_j)$.

(iv) $\bar{x} \in \mathbb{R}_{\varepsilon}^n$ with $\bar{b}' = \bar{A} \otimes \bar{x}$ is the solution for Problem 1.2, so $\forall j \in N$ satisfies:

a'. $\bar{x}_j = \varepsilon$ or

b'. $\bar{x}_j = (\bar{A}^* \otimes' \bar{v}_a)_j$.

Thus, from (i) and (iii) and also from (ii) and (iv), we have:

If $x \in I(\mathbb{R})_{\varepsilon}^n$ such that $b' = A \otimes x$ is the solution for Problem 2.2, then $\forall j \in N$ satisfies:

a. $x_j = [\varepsilon, \varepsilon]$ or

b. $x_j = [\min((\underline{A}^* \otimes' \underline{v}_{\underline{a}})_j, (\bar{A}^* \otimes' \bar{v}_{\bar{a}})_j), (\bar{A}^* \otimes' \bar{v}_{\bar{a}})_j]_j$. \square

B. Maximizing the range norm of matrix image set over interval max-plus algebra

The problem of maximizing the range norm of matrix image set can be formulated as follows:

Problem 2.3. Given that matrix $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$, solve:

$$\text{maximize } I\delta(b)$$

$$\text{subject to } b \in \text{Im}(A).$$

Definition 2.12. Given that matrix $b \in I(\mathbb{R})_{\varepsilon}^m$ with $b \approx [\underline{b}, \bar{b}]$. The function $I\delta(b)$ is said to be *maximum* if and only if $\delta(\bar{b})$ and $\delta(\underline{b})$ are maximum.

Maximizing the range norm if its matrix is finite

In this section, we discuss maximizing the range norm if its matrix is finite.

Theorem 2.13. Given that matrix $A \in I(\mathbb{R})^{m \times n}$ with $A = (A_1 \ A_2 \ \dots \ A_n)$ and $A_j \approx [\underline{A}_j, \bar{A}_j]$; $j = 1, 2, \dots, n$; then for each $b \in \text{Im}(A)$,

$$I\delta(b) \leq [\min(\max_{j=1,2,\dots,n} \delta(\underline{A}_j), \max_{j=1,2,\dots,n} \delta(\bar{A}_j)), \max_{j=1,2,\dots,n} \delta(\bar{A}_j)].$$

Proof. Suppose that $A \in I(\mathbb{R})^{m \times n}$ and $b \in \text{Im}(A)$ with $A \approx [\underline{A}, \bar{A}]$ and $b \approx [\underline{b}, \bar{b}]$, it means that $\underline{A}, \bar{A} \in \mathbb{R}^{m \times n}$ and $\underline{b} \in \text{Im}(\underline{A})$, $\bar{b} \in \text{Im}(\bar{A})$. Based

on concept in max-plus algebra, it is obtained that:

$$\begin{aligned}
\delta(\underline{b}) &\leq \max_{j=1, 2, \dots, n} \delta(\underline{A}_j) \\
&= \max_{j=1, 2, \dots, n} (\max_{i=1, 2, \dots, m} \underline{a}_{ij} - \min_{i=1, 2, \dots, m} \underline{a}_{ij}) \text{ and} \\
\delta(\bar{b}) &\leq \max_{j=1, 2, \dots, n} \delta(\bar{A}_j) \\
&= \max_{j=1, 2, \dots, n} (\max_{i=1, 2, \dots, m} \bar{a}_{ij} - \min_{i=1, 2, \dots, m} \bar{a}_{ij}).
\end{aligned}$$

Therefore,

$$\begin{aligned}
I\delta(b) &= [\min(\delta(\underline{b}), \delta(\bar{b})), \delta(\bar{b})] \\
&\leq [\min(\max_{j=1, 2, \dots, n} \delta(\underline{A}_j), \max_{j=1, 2, \dots, n} \delta(\bar{A}_j)), \max_{j=1, 2, \dots, n} \delta(\bar{A}_j)] \\
&= [\min(\max_{j=1, 2, \dots, n} (\max_{i=1, 2, \dots, m} \underline{a}_{ij} - \min_{i=1, 2, \dots, m} \underline{a}_{ij}), \\
&\quad \max_{j=1, 2, \dots, n} (\max_{i=1, 2, \dots, m} \bar{a}_{ij} - \min_{i=1, 2, \dots, m} \bar{a}_{ij})), \\
&\quad \max_{j=1, 2, \dots, n} (\max_{i=1, 2, \dots, m} \bar{a}_{ij} - \min_{i=1, 2, \dots, m} \bar{a}_{ij})]. \quad \square
\end{aligned}$$

Maximizing the range norm if its matrix is not finite

In the last section, we discuss maximizing the range norm if its matrix is not finite.

Theorem 2.14. *Given that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ is double $I(\mathbb{R})$ -astic and not finite, i.e., $\exists i \in M, j \in N$ such that $a_{ij} = [\varepsilon, \varepsilon]$, then Problem 2.3 is unbounded.*

Proof. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ with $A \approx [\underline{A}, \bar{A}]$ is double $I(\mathbb{R})$ -astic and infinite so $\exists i \in M, j \in N$ such that $a_{ij} = [\varepsilon, \varepsilon]$ which means that $\underline{A}, \bar{A} \in \mathbb{R}_{\varepsilon}^{m \times n}$ is double \mathbb{R} -astic and $\exists i \in M, j \in N$ such that $\underline{a}_{ij} = \varepsilon$ and

$\bar{a}_{ij} = \varepsilon$. Therefore, according to concept in max-plus algebra, Problem 1.3 is related to matrices \underline{A} and \bar{A} which have unbounded solution, namely:

a. Given that matrix $\underline{A} \in \mathbb{R}_\varepsilon^{m \times n}$, solve:

$$\text{maximize } \delta(\underline{b})$$

$$\text{subject to } \underline{b} \in \text{Im}(\underline{A}).$$

b. Given that matrix $\bar{A} \in \mathbb{R}_\varepsilon^{m \times n}$, solve:

$$\text{maximize } \delta(\bar{b})$$

$$\text{subject to } \bar{b} \in \text{Im}(\bar{A}).$$

Therefore, Problem 2.3, that is given matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, solve:

$$\text{maximize } I\delta(b)$$

$$\text{subject to } b \in \text{Im}(A)$$

has unbounded solution. □

III. Concluding Remarks

The main results are obtained the way to:

a. Minimizing the range norm of matrix image set over interval max-plus algebra, with its finite and not finite image vector.

These results are presented in Theorem 2.10 and Theorem 2.11.

b. Maximizing the range norm of matrix image set over max-plus algebra, with its finite and not finite matrix.

These results are presented in Theorem 2.13 and Theorem 2.14.

References

- [1] M. Akian, G. Cohen, S. Gaubert, J. P. Quadrat and M. Viot, Max-plus algebra and applications to system theory and optimal control, Proceedings of the International Congress of Mathematicians, Zurich, Switzerland, 1994.
- [2] M. H. Andersen, Max-plus algebra: properties and applications, A Thesis Submitted to the Department of Mathematics in Partial Fulfillment of the Requirements for the Degree of Master of Science in Mathematics, 2002.
- [3] F. Bacelli, G. Cohen, J. G. Older and J. P. Quadrat, Synchronization and Linearity, John Wiley & Sons, New York, 2001.
- [4] P. Butkovic and K. P. Tam, On some properties of the image of a max linear mapping, *Contemp. Math.* 495 (2009), 115-126.
- [5] R. A. Cuninghame-Green and P. Butkovic, Bases in max-algebra, *Linear Algebra Appl.* 389 (2004), 107-120.
- [6] K. G. Farlow, Max-plus algebra, Master's Thesis Submitted to the Faculty of the Virginia Polytechnic Institute and State University in Partial Fulfillment of the Requirements for the Degree of Masters in Mathematics, 2009,
- [7] Z. R. Konigsberg, A generalized eigenmode algorithm for reducible regular matrices over the max-plus algebra, *Int. Math. Forum* 4(24) (2009), 1157-1171.
- [8] V. Kreinovich, Why intervals? Why fuzzy numbers? Towards a new justification, *Int. J. Inform. Technol. Intell. Comput.* 2(1) (2007), 113-119.
- [9] M. A. Rudhito, Fuzzy number max-plus algebra and its application to fuzzy scheduling and queuing network problems, The Study Program S3 Mathematics Faculty of Mathematics and Natural Sciences, Gadjah Mada University, 2011 (Indonesian).
- [10] S. Sergeev, Max-plus definite matrix closures and their eigenspaces, *Linear Algebra Appl.* 421(2-3) (2007), 182-201.
- [11] Siswanto, A. Suparwanto and M. A. Rudhito, Minimizing range norm of the image set of matrix over interval max-plus algebra, Proceedings of the National Seminar on Mathematics and Applications, Department of Mathematics, Faculty of Science and Technology, Airlangga University, 2013 (Indonesian).
- [12] K. P. Tam, Optimizing and approximating eigenvectors in max algebra thesis, A Thesis Submitted to The University of Birmingham for the Degree of Doctor of Philosophy (Ph.D.), 2010.