




Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: M Andy Rudhito
Assignment title: Periksa similarity
Submission title: STRONGLY REGULAR MATRICES AND SIMPLE IMAGE SET IN I...
File name: R_MATRICES_AND_SIMPLE_IMAGE_SET_IN_INTERVAL_MAX-PL...
File size: 121.63K
Page count: 16
Word count: 4,603
Character count: 19,238
Submission date: 11-Apr-2022 08:47AM (UTC+0700)
Submission ID: 1807183449



STRONGLY REGULAR MATRICES AND SIMPLE IMAGE SET IN INTERVAL MAX-PLUS ALGEBRA

by Rudhito M Andy

Submission date: 11-Apr-2022 08:47AM (UTC+0700)

Submission ID: 1807183449

File name: R_MATRICES_AND_SIMPLE_IMAGE_SET_IN_INTERVAL_MAX-PLUS_ALGEBRA.pdf (121.63K)

Word count: 4603

Character count: 19238



STRONGLY REGULAR MATRICES AND SIMPLE IMAGE SET IN INTERVAL MAX-PLUS ALGEBRA

Siswanto^{1,2}, Ari Suparwanto² and M. Andy Rudhito³

¹Mathematics Department
Faculty of Mathematics and Natural Sciences
Sebelas Maret University
Indonesia
e-mail: sis.mipauns@yahoo.co.id

²Mathematics Department
Faculty of Mathematics and Natural Sciences
Gadjah Mada University
Indonesia
e-mail: ari_suparwanto@yahoo.com

³The Study Program in Mathematics Education
Faculty of Teacher Training and Education
Sanata Dharma University
Indonesia
e-mail: arudhito@yahoo.co.id

Abstract

4

Suppose that \mathbb{R} is the set of real numbers and $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$ with $\varepsilon = -\infty$. Max-plus algebra is the set \mathbb{R}_ε that is equipped with two operations, maximum and addition. Matrices over max-plus algebra

Received: September 7, 2015; Revised: September 25, 2015; Accepted: October 9, 2015

2010 Mathematics Subject Classification: 15A15, 05C50, 15A30.

Keywords and phrases: image set, strongly regular matrix, simple image set.

Communicated by K. K. Azad

are matrices whose elements belong to \mathbb{R}_ε . The set $I(\mathbb{R})_\varepsilon = \{x = [\underline{x}, \bar{x}] \mid \underline{x}, \bar{x} \in \mathbb{R}, \varepsilon < \underline{x} \leq \bar{x}\} \cup \{\varepsilon\}$ with $\varepsilon = [\varepsilon, \varepsilon]$ which is equipped with maximum and addition operations is known as interval max-plus algebra. Matrices over interval max-plus algebra are matrices whose elements belong to $I(\mathbb{R}_\varepsilon)$. The research about image set, strongly regular matrix and simple image set in max-plus algebra have been done. In this paper, we investigate image set, strongly regular matrix and simple image set in interval max-plus algebra.

1. Introduction

According to Baccelli et al. [1], max-plus algebra has been applied for modelling and analyzing problems in the area of planning, communication, production, queuing system with finite capacity, parallel computation and traffic. Max-plus algebra is the set of $\mathbb{R}_\varepsilon = \mathbb{R} \cup \{\varepsilon\}$ equipped with two operations, \otimes (maximum) and \oplus (addition), where \mathbb{R} is the set of real numbers with $\varepsilon = -\infty$ (Baccelli et al. [1]). According to Tam [7], max-plus algebra is a linear algebra over semiring \mathbb{R}_ε that is equipped with \oplus (maximum) and \otimes (addition) operations, while min-plus algebra is a linear algebra over semiring $\mathbb{R}_{\varepsilon'} = \mathbb{R} \cup \{\varepsilon'\}$, $\varepsilon' = \infty$ equipped with \oplus' (minimum) and \otimes' (addition) operations. Tam [7] also defined complete max-plus algebra and complete min-plus algebra. Complete max-plus algebra is a linear algebra over semiring $\mathbb{R}_{\varepsilon, \varepsilon'} = \mathbb{R}_\varepsilon \cup \{\varepsilon'\}$ equipped with \oplus (maximum) and \otimes (addition) operations, while complete min-plus algebra is a linear algebra over semiring $\mathbb{R}_{\varepsilon', \varepsilon} = \mathbb{R}_{\varepsilon'} \cup \{\varepsilon\}$ equipped with \oplus' (minimum) and \otimes' (plus) operations. Following the definition of max-plus algebra given by Baccelli et al. [1], we define min-plus algebra is the set of $\mathbb{R}_{\varepsilon'}$ equipped with \oplus' (minimum) and \otimes' (plus) operations. Besides that, we define complete max-plus algebra is the set $\mathbb{R}_{\varepsilon, \varepsilon'} = \mathbb{R}_\varepsilon \cup \{\varepsilon'\}$ equipped with \oplus (maximum) and \otimes (addition) operations, while complete min-plus algebra is the set $\mathbb{R}_{\varepsilon', \varepsilon} = \mathbb{R}_{\varepsilon'} \cup \{\varepsilon\}$ equipped with \oplus' (minimum) and \otimes' (plus) operations.

It shows that $\mathbb{R}_{\epsilon, \epsilon'} = \mathbb{R}_{\epsilon} \cup \{\epsilon'\} = \mathbb{R} \cup \{\epsilon, \epsilon'\} = \mathbb{R}_{\epsilon'} \cup \{\epsilon\} = \mathbb{R}_{\epsilon', \epsilon}$. Furthermore, $\mathbb{R}_{\epsilon, \epsilon'}$ and $\mathbb{R}_{\epsilon', \epsilon}$ are written as $\bar{\mathbb{R}}$. It can be formed a set of matrices in the size $m \times n$ of the sets \mathbb{R}_{ϵ} , $\mathbb{R}_{\epsilon'}$, and $\bar{\mathbb{R}}$. The set of matrices with components in \mathbb{R}_{ϵ} is denoted by $\mathbb{R}_{\epsilon}^{m \times n}$. The matrix $A \in \mathbb{R}_{\epsilon}^{m \times n}$ is called *matrix over max-plus algebra*. The matrix over min-plus algebra, over complete max-plus algebra and over min-plus algebra can also be defined (Tam [7]).

In 2010, Tam [7] discussed the linear system in max-plus algebra, image set and strongly regular matrix. Tam [7] and Butkovic [2, 3] mentioned that the solution of the linear system in max-plus algebra is closely related to image set, simple image set and strongly regular matrix. Butkovic [3-5] have shown the characteristics of the strongly regular matrix, that is matrix has strong permanent.

Rudhito [6] has generalized max-plus algebra into interval max-plus algebra and fuzzy number max-plus algebra. Moreover, Siswanto et al. [8] have discussed linear equation in interval max-plus algebra. Based on the discussion on the max-plus algebra that is connected to the system of linear equation, in this paper we discuss image set, strongly regular matrix and simple image set in interval max-plus algebra. Furthermore, it will also discuss matrix has strong permanent. The matrix has strong permanent is the characteristic of strongly regular matrix.

Before discussing the result of the research, the necessary concepts are discussed. These are interval max-plus algebra, system of linear equation in max-plus algebra and system of linear equation and the system of linear inequation in interval max-plus algebra.

Related to the system of linear equation $A \otimes x = b$, Tam [7] defined the solution set of the system of linear equation and image set of matrix A .

Definition 1.1. Given $A \in \mathbb{R}_{\epsilon}^{m \times n}$ and $b \in \mathbb{R}_{\epsilon}^m$. The solution set of the system of linear equation $A \otimes x = b$ is $S(A, b) = \{x \in \mathbb{R}_{\epsilon}^n \mid A \otimes x = b\}$ and the image set of matrix A is $Im(A) = \{A \otimes x \mid x \in \mathbb{R}_{\epsilon}^n\}$.

Based on the above definition, the following theorem and corollary are given.

Theorem 1.2. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $b \in \mathbb{R}_\varepsilon^m$. Vector $b \in \text{Im}(A)$ if and only if $S(A, b) \neq \emptyset$.

Corollary 1.3. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $b \in \mathbb{R}_\varepsilon^m$. Vector $b \in \text{Im}(A)$ if and only if $A \otimes (A^* \otimes b) = b$.

The following is the definitions of subspace and theorem which states that for $A \in \mathbb{R}_\varepsilon^{m \times n}$, $\text{Im}(A)$ is a subspace.

Definition 1.4. Suppose that $S \subseteq \mathbb{R}_\varepsilon^n$. If for each $x, y \in S$ and for each $\alpha \in \mathbb{R}_\varepsilon'$, $\alpha \otimes x \in S$ and $x \oplus y \in S$, then S is known as subspace of max-plus algebra.

Theorem 1.5. Suppose that $A \in \mathbb{R}_\varepsilon^{m \times n}$. If $\alpha, \beta \in \mathbb{R}$ and $u, v \in \text{Im}(A)$ then $\alpha \otimes u \oplus \beta \otimes v \in \text{Im}(A)$.

In max-plus algebra there are the definitions of strongly linearly independent and strongly regular matrix.

Definition 1.6. The vectors $A_1, \dots, A_n \in \mathbb{R}_\varepsilon^m$ are said to be *linearly dependent* if one of those vectors can be expressed as a linear combination of other vectors. Otherwise A_1, \dots, A_n are considered linearly independent.

Definition 1.7. The vectors $A_1, \dots, A_n \in \mathbb{R}_\varepsilon^m$ are said to be *strongly linearly independent* if there is $b \in \mathbb{R}^m$ as such so that b can be uniquely expressed as a linear combination of A_1, \dots, A_n .

Definition 1.8. For the vectors $A_1, \dots, A_n \in \mathbb{R}_\varepsilon^m$ which are strongly linearly independent, if $m = n$, then the matrix $A = (A_1, \dots, A_n)$ is known as strongly regular.

Below is the definition of simple image set.

Definition 1.9. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$. The simple image set of matrix A is defined as $S_A = \{b \in \mathbb{R}^m \mid A \otimes x = b \text{ has unique solution}\}$.

Theorem 1.10. Suppose $A \in \mathbb{R}_\varepsilon^{m \times n}$ is double \mathbb{R} -astic and $b \in \mathbb{R}^m$, then $|S(A, b)| \in \{0, 1, \infty\}$.

Definition 1.11. Suppose $A \in \mathbb{R}_\varepsilon^{m \times n}$, $T(A)$ is defined as $\{|S(A, b)| \mid b \in \mathbb{R}^m\}$.

Theorem 1.12. If $A \in \mathbb{R}_\varepsilon^{m \times n}$ is double \mathbb{R} -astic, then $T(A)$ is $\{0, \infty\}$ or $\{0, 1, \infty\}$.

Based on Definition 1.7, the column vectors of A are strongly linearly independent if and only if $1 \in T(A)$. In max-plus algebra, a matrix that has strong permanent is also defined (Butkovic [4]).

Definition 1.13. Given $A \in \mathbb{R}_\varepsilon^{m \times n}$ and $\pi \in P_n$, where P_n is the set of all permutation of $N = \{1, 2, \dots, n\}$. $\omega(A, \pi)$ is defined as $\prod_{j \in n}^\otimes A_{j, \pi(j)}$ and the permanent of matrix A is $\text{maper}(A) = \sum_{\pi \in P_n}^\oplus \omega(A, \pi)$.

Tam [7] defined that a matrix has strong permanent as given below.

Definition 1.14. The set of all permutation is written as $ap(A)$, and is defined as $ap(A) = \{\pi \in P_n \mid \text{maper}(A) = \omega(A, \pi)\}$. Matrix A is said to have strong permanent if $|ap(A)| = 1$.

Butkovic [4] and Tam [7] explained the connection between strong regular matrix and matrix has strong permanent as below.

Theorem 1.15. Suppose that $A \in \mathbb{R}_\varepsilon^{m \times n}$ is double \mathbb{R} -astic. Matrix A is strongly regular matrix if and only if matrix A has strong permanent.

In the following section, interval max-plus algebra, matrix over interval max-plus algebra and the system of linear equation in interval max-plus algebra is in (Rudhito [6], Siswanto et al. [8]) are presented.

Closed interval x in \mathbb{R}_ε is a subset of \mathbb{R}_ε in the form of $x = [\underline{x}, \bar{x}] = \{x \in \mathbb{R}_\varepsilon \mid \underline{x} \leq x \leq \bar{x}\}$. The interval x in \mathbb{R}_ε is known as *interval max-plus*. A number $x \in \mathbb{R}_\varepsilon$ can be expressed as interval $[x, x]$.

Definition 1.16. $I(\mathbb{R})_\varepsilon = \{x = [\underline{x}, \bar{x}] \mid \underline{x}, \bar{x} \in \mathbb{R}, \varepsilon < \underline{x} \leq \bar{x}\} \cup \{\varepsilon\}$ is formed, with $\varepsilon = [\varepsilon, \varepsilon]$. On the set $I(\mathbb{R})_\varepsilon$, the operations \oplus and \otimes with $x \oplus y = [\underline{x} \oplus \underline{y}, \bar{x} \oplus \bar{y}]$ and $x \otimes y = [\underline{x} \otimes \underline{y}, \bar{x} \otimes \bar{y}]$ for every $x, y \in I(\mathbb{R})_\varepsilon$ are defined. Furthermore they are known as *interval max-plus algebra* and notated as $I(\mathbb{R})_{\max} = (I(\mathbb{R})_\varepsilon; \oplus, \otimes)$.

Definition 1.17. The set of matrices in the size $m \times n$ with the elements in $I(\mathbb{R})_\varepsilon$ is notated as $I(\mathbb{R})_\varepsilon^{m \times n}$, namely:

$$I(\mathbb{R})_\varepsilon^{m \times n} = \{A = [A_{ij}] \mid A_{ij} \in I(\mathbb{R})_\varepsilon; \text{ for } i = 1, 2, \dots, m \text{ and } j = 1, 2, \dots, n\}.$$

Matrices which belong to $I(\mathbb{R})_\varepsilon^{m \times n}$ are known as *matrices over interval max-plus algebra* or shortly *interval matrices*.

Definition 1.18. For $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, matrix $\underline{A} = [\underline{A}_{ij}] \in \mathbb{R}_\varepsilon^{m \times n}$ and $\bar{A} = [\bar{A}_{ij}] \in \mathbb{R}_\varepsilon^{m \times n}$ are defined, as a lower bound matrix and an upper bound matrix of the interval matrix A , respectively.

Definition 1.19. Given that interval matrix $A \in I(\mathbb{R})_\varepsilon^{m \times n}$, and \underline{A} and \bar{A} are a lower bound matrix and an upper bound matrix of the interval matrix A , respectively. The matrix interval of A is defined, namely $[\underline{A}, \bar{A}] = \{A = \mathbb{R}_\varepsilon^{m \times n} \mid \underline{A} \leq A \leq \bar{A}\}$ and $I(\mathbb{R}_\varepsilon^{m \times n})_b = \{[\underline{A}, \bar{A}] \mid A \in (\mathbb{R})_\varepsilon^{m \times n}\}$.

Suppose that $A \in I(\mathbb{R})_\varepsilon^{m \times n}$ and $b = (b_1, \dots, b_m) \in I(\mathbb{R})_\varepsilon^m$. The equation system $A \otimes x = b$ is known as the *system of linear equation in interval*

max-plus algebra. Furthermore, $A \approx [\underline{A}, \bar{A}]$, $x \approx [\underline{x}, \bar{x}]$ and $b \approx [\underline{b}, \bar{b}]$ with $[\underline{A}, \bar{A}] \in I(\mathbb{R}_\varepsilon^{n \times n})_b$, $[\underline{x}, \bar{x}], [\underline{b}, \bar{b}] \in I(\mathbb{R}_\varepsilon^n)_b$. To find the solution of the system of linear equation in interval max-plus algebra, the solution of the system of linear equation in max-plus algebra $\underline{A} \otimes \underline{x} = \underline{b}$ and $\bar{A} \otimes \bar{x} = \bar{b}$ should be determined.

Definition 1.20. Given the system of linear equation $A \otimes x = b$ with $A \in I(\mathbb{R})_e^{m \times n}$ and $b \in I(\mathbb{R})_e^m$. $IS(A, b) = \{x \in I(\mathbb{R})_e^m \mid A \otimes x = b\}$ is defined.

Along with the discussion the system of linear equation in max-plus algebra, without loss of generality, the system of linear equation in interval max-plus algebra $A \otimes x = b$ with $A \in I(\mathbb{R})_e^{m \times n}$ double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})_e^m$ is discussed.

Theorem 1.21. Suppose that $A \in I(\mathbb{R})_e^{m \times n}$ double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})_e^m$. Vector $y \in IS(A, b)$ if and only if $\underline{y} \in S(\underline{A}, \underline{b})$, $\bar{y} \in S(\bar{A}, \bar{b})$ and $\underline{y} \leq \bar{y}$. Furthermore y can be expressed as $y \approx [\underline{y}, \bar{y}]$.

Each of the following two corollaries is a criterion of the system of linear equation which has solution and a criterion the system of linear equation which has unique solution. The notations of $M = \{1, 2, \dots, m\}$ and $N = \{1, 2, \dots, n\}$ are used to simplify.

Corollary 1.22. Suppose $A \in I(\mathbb{R})_e^{m \times n}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})_e^m$, then the following three statements are equivalent:

- $IS(A, b) \neq \emptyset$.
- $\hat{y} \in IS(A, b)$.
- $\bigcup_{j \in N} M_j(\underline{A}, \underline{b}) = M$ and $\bigcup_{j \in N} M_j(\bar{A}, \bar{b}) = M$.

Corollary 1.23. Suppose $A \in I(\mathbb{R})_e^{m \times n}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$, then $IS(A, b) = \{\hat{y}\}$ if and only if

- $\bigcup_{j \in N} M_j(\underline{A}, \underline{b}) = M$ and $\bigcup_{j \in N} M_j(\bar{A}, \bar{b}) = M$.
- $\bigcup_{j \in N'} M_j(\underline{A}, \underline{b}) \neq M$ for every $N' \subseteq N$, $N' \neq N$ and $\bigcup_{j \in N''} M_j(\bar{A}, \bar{b}) \neq M$ for every $N'' \subseteq N$, $N'' \neq N$.

2. Main Results

In this section, we present new results of the research. The results that are obtained, are related with the system of linear equation in interval max-plus algebra $A \otimes x = b$, $A \in I(\mathbb{R})_e^{m \times n}$ and $b \in I(\mathbb{R})^m$.

We will start the definition of the image set of matrix A .

Definition 2.1. Given $A \in I(\mathbb{R})_e^{m \times n}$. The image set of matrix A is $Im(A) = \{A \otimes x \mid x \in I(\mathbb{R})_e^n\}$.

Based on the definition of the image set of a matrix, a theorem which demonstrated the necessary and sufficient conditions of the elements of the image set is presented.

Theorem 2.2. Given $A \in I(\mathbb{R})_e^{m \times n}$ and $b \in I(\mathbb{R})_e^m$. Vector $b \in Im(A)$ if and only if $IS(A, b) \neq \emptyset$.

Proof. Suppose that $A \approx [\underline{A}, \bar{A}]$ and $b \approx [\underline{b}, \bar{b}]$ with $\underline{A}, \bar{A} \in \mathbb{R}_e^{m \times n}$ and $\underline{b}, \bar{b} \in \mathbb{R}_e^m$. Since $b \in Im(A)$, it means that $\underline{b} \in Im(\underline{A})$ and $\bar{b} \in Im(\bar{A})$. Therefore, based on Theorem 1.2, i.e., $S(\underline{A}, \underline{b}) \neq \emptyset$ and $S(\bar{A}, \bar{b}) \neq \emptyset$, then $IS(A, b) \neq \emptyset$. On the other hand, since $IS(A, b) \neq \emptyset$, $S(\underline{A}, \underline{b}) \neq \emptyset$ and $S(\bar{A}, \bar{b}) \neq \emptyset$. Thus, based on Theorem 1.2 that is $\underline{b} \in Im(\underline{A})$ and $\bar{b} \in Im(\bar{A})$, then $b \in Im(A)$ with $b \approx [\underline{b}, \bar{b}]$. \square

With regard to ³⁷Theorem 2.2, the following corollary is obtained:

Corollary 2.3. ¹¹Given $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ and $b \in I(\mathbb{R})_{\varepsilon}^m$. Vector $b \in \text{Im}(A)$ if and only if $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$.

Proof. Suppose that $A \approx [A, \overline{A}]$ and $b \approx [\underline{b}, \overline{b}]$. Since $b \in \text{Im}(A)$, ¹⁰it means that $\underline{b} \in \text{Im}(\underline{A})$ and $\overline{b} \in \text{Im}(\overline{A})$. Based on Corollary 1.3, $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$. On the other hand, since $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$, based on Corollary 1.3, $\underline{b} \in \text{Im}(\underline{A})$ and $\overline{b} \in \text{Im}(\overline{A})$ are obtained. Therefore, $b \approx [\underline{b}, \overline{b}] \in \text{Im}(A)$. \square

The following theorem presents that $\text{Im}(A)$ is a subspace of $I(\mathbb{R})_{\varepsilon}^m$.

Theorem 2.4. Given $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$, $\text{Im}(A)$ is a subspace of $I(\mathbb{R})_{\varepsilon}^m$.

Proof. Since $\text{Im}(A) = \{A \otimes x \mid x \in I(\mathbb{R})_{\varepsilon}^n\}$, $\text{Im}(A) \subseteq I(\mathbb{R})_{\varepsilon}^m$ and $\text{Im}(A) \neq \emptyset$. Taking $p, q \in \text{Im}(A)$ and $\alpha, \beta \in I(\mathbb{R})$, accordingly, $\alpha \otimes \underline{p} \oplus \beta \otimes \underline{q} \approx [\underline{\alpha} \otimes \underline{p} \oplus \underline{\beta} \otimes \underline{q}, \overline{\alpha} \otimes \underline{p} \oplus \overline{\beta} \otimes \underline{q}]$ with $\underline{\alpha} \otimes \underline{p} \oplus \underline{\beta} \otimes \underline{q} \in \text{Im}(\underline{A})$ and $\overline{\alpha} \otimes \underline{p} \oplus \overline{\beta} \otimes \underline{q} \in \text{Im}(\overline{A})$. It means that $\alpha \otimes \underline{p} \oplus \beta \otimes \underline{q} \in \text{Im}(A)$. In other words, $\alpha \otimes \underline{p} \oplus \beta \otimes \underline{q} \in \text{Im}(A)$ and hence $\text{Im}(A)$ is a subspace of $I(\mathbb{R})_{\varepsilon}^m$. \square

Several definitions and theorems are presented to support the discussion on strongly regular matrix.

Definition 2.5. ¹⁴The vectors $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^m$ is said to be linearly dependent if one of those vectors can be said to be as linear combination of other vectors. Otherwise then, A_1, \dots, A_n is said to be linearly independent.

Theorem 2.6. If the vectors $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^m$ with $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2]$, ..., $A_n \approx [\underline{A}_n, \overline{A}_n]$ are linearly dependent, then $\underline{A}_1, \underline{A}_2, \dots, \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, \dots, \overline{A}_n$ are linearly independent.

Proof. It is known that $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^m$ is linearly dependent. It means that one of those vectors can be stated as a linear combination of the others. Without losing the generality of the proof, for example $A_1 = \bigoplus_{i=2}^n (\alpha_i \otimes A_i)$ with $\alpha_i \in I(\mathbb{R})_m$, since $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2]$, ..., $A_n \approx [\underline{A}_n, \overline{A}_n]$, $\underline{A}_1 = \bigoplus_{i=2}^n (\underline{\alpha}_i \otimes \underline{A}_i)$ and $\overline{A}_1 = \bigoplus_{i=2}^n (\overline{\alpha}_i \otimes \overline{A}_i)$. In other words, $\underline{A}_1, \underline{A}_2, \dots, \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, \dots, \overline{A}_n$ are linearly dependent.

Definition 2.7. The vectors $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^m$ is said to be *strongly linearly independent* if there is $b \in I(\mathbb{R})^m$, b can be uniquely expressed as a linear combination of A_1, \dots, A_n , that is $b = \bigoplus_{i=1}^n (\alpha_i \otimes A_i)$, $\alpha_i \in I(\mathbb{R})$.

The following is the definition of strongly regular matrix.

Definition 2.8. Matrix $A = (A_1, A_2 \dots A_n) \in I(\mathbb{R})_{\varepsilon}^{n \times n}$ is said to be *strongly regular* if the vectors $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^n$ are strongly linearly independent.

Based on the definition of strongly linearly independent in the max-plus algebra, the following theorem is formulated.

Theorem 2.9. If the vectors $A_1, \dots, A_n \in I(\mathbb{R})_{\varepsilon}^m$ with $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2]$, ..., $A_n \approx [\underline{A}_n, \overline{A}_n]$ are strongly linearly independent, then $\underline{A}_1, \underline{A}_2, \dots, \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, \dots, \overline{A}_n$ are strongly linearly independent.

Proof. It is known that $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2]$, ..., $A_n \approx [\underline{A}_n, \overline{A}_n]$ are strongly linearly independent, then there is $b \approx [\underline{b}, \overline{b}] \in I(\mathbb{R})^m$ so that b can be uniquely expressed as linear combination of A_1, \dots, A_n . Suppose that $b = \bigoplus_{i=1}^n (\alpha_i \otimes A_i)$; since

$$\bigoplus_{i=1}^n (\alpha_i \otimes A_i) \approx \bigoplus_{i=1}^n (\underline{\alpha}_i \otimes \underline{A}_i),$$

$\oplus_{i=1}^n (\bar{\alpha}_i \otimes \bar{A}_i)$, $\underline{b} = \oplus_{i=1}^n (\underline{\alpha}_i \otimes \underline{A}_i)$ and $\bar{b} = \oplus_{i=1}^n (\bar{\alpha}_i \otimes \bar{A}_i)$. It means that \underline{b} can be uniquely expressed as a linear combination of $\underline{A}_1, \underline{A}_2, \dots, \underline{A}_n$ and \bar{b} can be uniquely expressed as a linear combination $\bar{A}_1, \bar{A}_2, \dots, \bar{A}_n$. In other words, $\underline{A}_1, \underline{A}_2, \dots, \underline{A}_n$ and $\bar{A}_1, \bar{A}_2, \dots, \bar{A}_n$ are strongly linearly independent. \square

The following is the definition of simple image set.

Definition 2.10. Given $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$. The set $IS_A = \{b \in I(\mathbb{R})^m \mid A \otimes x = b \text{ has a unique solution}\}$ is defined as a simple image set of matrix A .

The regularity properties of the matrix are closely related to the number of the solutions of the system of linear equation. It can be stated in the following definitions and theorems.

Theorem 2.11. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ is a double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$ is the number of the elements of $IS(A, b)$ is $|IS(A, b)| \in \{0, 1, \infty\}$.

Proof. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$. Suppose that $A \approx [\underline{A}, \bar{A}]$ and $b \approx [\underline{b}, \bar{b}]$ with each of \underline{A} and \bar{A} is double \mathbb{R} -astic, while $\underline{b}, \bar{b} \in \mathbb{R}^m$. Based on the Theorem 1.10, $|S(\underline{A}, \underline{b})| \in \{0, 1, \infty\}$ and $|S(\bar{A}, \bar{b})| \in \{0, 1, \infty\}$. Therefore, following the Definition 1.18, it is confirmed that $|IS(A, b)| \in \{0, 1, \infty\}$. \square

Definition 2.12. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$, $T(A)$ is defined as $\{|IS(A, b)| \mid b \in \mathbb{R}^m\}$.

Theorem 2.13. If $A \in I(\mathbb{R})_{\varepsilon}^{m \times n}$ is double $I(\mathbb{R})$ -astic, then $T(A) = \{0, \infty\}$ or $T(A) = \{0, 1, \infty\}$.

Proof. Suppose that $A \in I(\mathbb{R})_e^{m \times n}$, $A \approx [\underline{A}, \bar{A}]$ with $\underline{A} \in \mathbb{R}_e^{m \times n}$ and $\bar{A} \in \mathbb{R}_e^{m \times n}$. Based on the theorem for the case of max-plus algebra, $T(\underline{A}) = \{0, 1, \infty\}$ or $T(\underline{A}) = \{0, \infty\}$ and $T(\bar{A}) = \{0, \infty\}$ or $T(\bar{A}) = \{0, 1, \infty\}$. Therefore, $T(A) = \{0, \infty\}$ or $T(A) = \{0, 1, \infty\}$. \square

Based on Theorem 2.13, A is a strongly regular matrix if and only if $1 \in T(A)$. Then, the criteria of strongly regular matrix is discussed.

Definition 2.14. Given $A = (a_{ij}) \in I(\mathbb{R})_e^{n \times n}$ and $\pi \in P_n$, with P_n is the set of all permutations of $\{1, 2, \dots, n\}$. $\omega(A, \pi)$ is defined as $\bigotimes_{j \in n} a_{j, \pi(j)}$.

Theorem 2.15. If $A \in I(\mathbb{R})_e^{n \times n}$ double $I(\mathbb{R})$ -astic and $\pi \in P_n$ with $A \approx [\underline{A}, \bar{A}]$, then $\omega(A, \pi) = [\omega(\underline{A}, \pi), \omega(\bar{A}, \pi)]$.

Proof. If $A \in I(\mathbb{R})_e^{m \times n}$ and $\pi \in P_n$ with $A \approx [\underline{A}, \bar{A}]$, it means that

$$\begin{aligned} \omega(A, \pi) &= \bigotimes_{j \in n} a_{j, \pi(j)} = \bigotimes_{j \in n} [\underline{a}_{j, \pi(j)}, \bar{a}_{j, \pi(j)}] \\ &= [\bigotimes_{j \in n} \underline{a}_{j, \pi(j)}, \bigotimes_{j \in n} \bar{a}_{j, \pi(j)}] = [\omega(\underline{A}, \pi), \omega(\bar{A}, \pi)]. \quad \square \end{aligned}$$

Definition 2.16. The permanent of matrix A is defined as $\text{maper} \bigotimes_{\pi \in P_n} \omega(A, \pi)$.

Next discussion is matrix over interval max-plus algebra which has weak permanent and strong permanent.

Definition 2.17. Suppose that $A \in I(\mathbb{R})_e^{n \times n}$ double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \bar{A}]$. The set of all optimal permutations is $ap(A)$ with $ap(A) = \{\pi \in P_n \mid \text{maper}(\underline{A}) = \omega(\underline{A}, \pi) \text{ or } \text{maper}(\bar{A}) = \omega(\bar{A}, \pi)\}$. Matrix A is said to have strong permanent if $|ap(A)| = 1$ and it is said to have weak permanent if $|ap(\underline{A})| = 1$ and $|ap(\bar{A})| = 1$.

Theorem 2.18. Suppose that $A \in I(\mathbb{R})_e^{n \times n}$ is double $I(\mathbb{R})$ -astic. If matrix A has strong permanent, then matrix A has weak permanent.

Proof. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$, $A \approx [\underline{A}, \bar{A}]$ with $\underline{A} \in (\mathbb{R})_{\varepsilon}^{n \times n}$ and $\bar{A} \in \mathbb{R}_{\varepsilon}^{n \times n}$. Since A has strong permanent, it means that $|ap(A)| = 1$. Suppose that $\pi_1 \in ap(A)$, then $maper(\underline{A}) = \omega(\underline{A}, \pi_1)$ and $maper(\bar{A}) = \omega(\bar{A}, \pi_1)$. Therefore, $|ap(\underline{A})| = 1$ and $|ap(\bar{A})| = 1$. In other words, A has weak permanent. \square

Theorem 2.19. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$ is double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \bar{A}]$. Matrix A has weak permanent if and only if \underline{A} and \bar{A} have strong permanent.

Proof. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$, $A \approx [\underline{A}, \bar{A}]$ with $\underline{A} \in \mathbb{R}_{\varepsilon}^{n \times n}$ and $\bar{A} \in \mathbb{R}_{\varepsilon}^{n \times n}$. It is known that matrix A has weak permanent. It means that $|ap(\underline{A})| = 1$ and $|ap(\bar{A})| = 1$. Therefore, matrix \underline{A} and \bar{A} have strong permanent. On the other hand, suppose that \underline{A} and \bar{A} have strong permanent, it means that $|ap(\underline{A})| = 1$ and $|ap(\bar{A})| = 1$. Therefore, matrix A with $A \approx [\underline{A}, \bar{A}]$ has weak permanent. \square

Theorem 2.20. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$ is double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \bar{A}]$. Matrix A has strong permanent if and only if \underline{A} and \bar{A} have strong permanent as such that if $maper(\underline{A}) = \omega(\underline{A}, \pi_1)$ and $maper(\bar{A}) = \omega(\bar{A}, \pi_2)$, then $\pi_1 = \pi_2$.

Proof. Suppose that $A \in I(\mathbb{R})_{\varepsilon}^{n \times n}$ with $A \approx [\underline{A}, \bar{A}]$. It is known that matrix A has strong permanent. Since matrix A has strong permanent, it means that $|ap(A)| = 1$. Suppose that $\pi \in ap(A)$ it means that $\pi \in P_n$ so that $maper(\underline{A}) = \omega(\underline{A}, \pi)$ and $maper(\bar{A}) = \omega(\bar{A}, \pi)$. Thus, $|ap(\underline{A})| = 1$ and $|ap(\bar{A})| = 1$. Therefore, \underline{A} and \bar{A} have strong permanent. On the other hand, it is known that matrix \underline{A} and \bar{A} have strong permanent. If $maper(\underline{A}) =$

$\omega(\underline{A}, \pi_1)$ and $\text{maper}(\overline{A}) = \omega(\overline{A}, \pi_2)$, then $\pi_1 = \pi_2$. Therefore, $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$ with $\pi_1 \in |ap(\underline{A})|$, $\pi_2 \in |ap(\overline{A})|$ and $\pi_1 = \pi_2$. Since $ap(A) = \{\pi \in P_n \mid \text{maper}(\underline{A}) = \omega(\underline{A}, \pi) \text{ or } \text{maper}(\overline{A}) = \omega(\overline{A}, \pi)\}$, $|ap(A)| = 1$. Therefore, matrix A has strong permanent.

The following is theorem which has the characteristics of strongly regular matrix.

Theorem 2.21. Suppose that $A \in I(\mathbb{R})_e^{n \times n}$ is double $I(\mathbb{R})$ -astic. Matrix A is strongly regular if and only if matrix A has strong permanent.

Proof. Suppose that $A \in I(\mathbb{R})_e^{n \times n}$ is double $I(\mathbb{R})$ -astic, with $A \approx [\underline{A}, \overline{A}]$. It is known that A is strongly regular matrix. Suppose that A_1, \dots, A_n are the columns of matrix A . Since A is strongly regular matrix, there is $b \in I(\mathbb{R})^n$ so that b can be uniquely expressed as linear combination of A_1, \dots, A_n , that is $b = \bigoplus_{i=1}^n (x_i \otimes A_i)$, $x_i \in I(\mathbb{R})$. The equation $b = \bigoplus_{i=1}^n (x_i \otimes A_i) \Leftrightarrow A \otimes x = b$, is the system of linear equation in interval max-plus algebra which has a unique solution. Based on Corollary 1.23, $|ap(A)| = 1$. Therefore, matrix A has strong permanent. The converse, it is known that matrix A has strong permanent. Based on Theorem 2.20, matrix \underline{A} and \overline{A} have such strong permanent, so that if $\text{maper}(\underline{A}) = \omega(\underline{A}, \pi_1)$ and $\text{maper}(\overline{A}) = \omega(\overline{A}, \pi_2)$, $\pi_1 = \pi_2$. Therefore, matrix \underline{A} and \overline{A} are both strongly regular matrix, and thus, there are \underline{b} and \overline{b} which can be uniquely written as the linear combinations $\underline{b} = \bigoplus_{i=1}^n (x_i \otimes A_i)$ and $\overline{b} = \bigoplus_{i=1}^n (\bar{x}_i \otimes \overline{A}_i)$, in which A_1, A_2, \dots, A_n are the columns of matrix \underline{A} , while $\overline{A}_1, \overline{A}_2, \dots, \overline{A}_n$ are the columns of matrix \overline{A} . That $\underline{b} = \bigoplus_{i=1}^n (x_i \otimes A_i) \Leftrightarrow \underline{A} \otimes \underline{x} = \underline{b}$ and $\overline{b} = \bigoplus_{i=1}^n (\bar{x}_i \otimes \overline{A}_i) \Leftrightarrow \overline{A} \otimes \bar{x} = \overline{b}$. Thus, the system of linear equation $\underline{A} \otimes \underline{x} = \underline{b}$ and $\overline{A} \otimes \bar{x} = \overline{b}$ have

unique solution. Since $\text{maper}(\underline{A}) = \omega(\underline{A}, \pi_1)$ and $\text{maper}(\overline{A}) = \omega(\overline{A}, \pi_2)$, $\pi_1 = \pi_2$ so that $[b, \bar{b}] \approx b$. As the result, $[\underline{A} \otimes \underline{x}, \overline{A} \otimes \bar{x}] = [b, \bar{b}]$ and since $\overline{A \otimes x} \approx [\underline{A} \otimes \underline{x}, \overline{A} \otimes \bar{x}]$, $\overline{A \otimes x} = b$ or $b = \bigoplus_{i=1}^n (x_i \otimes A_i)$ with A_1, \dots, A_n as the columns of matrix A . It means that there is b which can be uniquely expressed as linear combination of A_1, \dots, A_n . This matrix A is known as strongly regular matrix. \square

3. Concluding Remarks

Based on the discussion, the following conclusions are obtained:

1. The necessary and sufficient condition of matrix $A \in I(\mathbb{R})_e^{n \times n}$, which is double $I(\mathbb{R})$ -astic is strongly regular matrix.
2. The system of linear equation in the interval max-plus algebra $A \otimes x = b$, $A \in I(\mathbb{R})_e^{n \times n}$ which is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^n$ has unique solution if and only if matrix A has strong permanent.

Acknowledgment

The authors thank the anonymous referees for their valuable suggestions which let to the improvement of the manuscript.

References

- [1] F. Baccelli, G. Cohen, J. G. Older and J. P. Quadrat, Synchronization and Linearity, John Wiley & Sons, New York, 2001.
- [2] P. Butkovic, Strongly regularity of matrices - a survey of results, Discrete Applied Mathematics 48 (1994), 45-68.
- [3] P. Butkovic, Simple image set of $(\max, +)$ linear mappings, Discrete Applied Mathematics 105 (2000), 73-86.
- [4] P. Butkovic, Max-algebra: the linear algebra of combinatorics?, Linear Algebra and Application 367 (2003), 313-335.

- [5] P. Butkovic, Max Linear Systems: Theory and Algorithm, Springer, London, 2010.
- [6] M. A. Rudhito, Fuzzy number max-plus algebra and its application to fuzzy scheduling and queuing network problems, Doctoral Program of Mathematics, Faculty of Mathematics and Natural Sciences, Gadjah Mada University, 2011 (Indonesian).
- [7] K. P. Tam, Optimizing and approximating eigenvectors in max algebra (Thesis), A Thesis Submitted to The University of Birmingham for the Degree of Doctor of Philosophy (Ph.D.), 2010.
- [8] Siswanto, A. Suparwanto, M. A. Rudhito, Existence of solution to the system of linear equations in interval max-plus algebra, Proceedings of the National Conference of Mathematics XVII, Institute of Technology Surabaya, 2014 (Indonesian).

STRONGLY REGULAR MATRICES AND SIMPLE IMAGE SET IN INTERVAL MAX-PLUS ALGEBRA

ORIGINALITY REPORT

24%

SIMILARITY INDEX

9%

INTERNET SOURCES

22%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|--|---|-----------------------------------|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">1</div> | <p style="color: red;">Lado Lenart. "Algebraic Way to Control Assembly Process", IFAC Proceedings Volumes, 1998</p> <p>Publication</p> | <p style="color: red;">2%</p> |
| <hr/> | | |
| <div style="background-color: magenta; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">2</div> | <p style="color: magenta;">Peter Butkovic. "Simple image set of (max,+) linear mappings", Discrete Applied Mathematics, 2000</p> <p>Publication</p> | <p style="color: magenta;">1%</p> |
| <hr/> | | |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">3</div> | <p style="color: purple;">Jianqin Zhou. "Generalized Partially Bent Functions", Future Generation Communication and Networking (FGCN 2007), 12/2007</p> <p>Publication</p> | <p style="color: purple;">1%</p> |
| <hr/> | | |
| <div style="background-color: teal; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">4</div> | <p style="color: teal;">etd.repository.ugm.ac.id</p> <p>Internet Source</p> | <p style="color: teal;">1%</p> |
| <hr/> | | |
| <div style="background-color: green; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">5</div> | <p style="color: green;">Peter Butkovič. "Chapter 6 Linear Independence and Rank. The Simple Image Set", Springer Science and Business Media LLC, 2010</p> <p>Publication</p> | <p style="color: green;">1%</p> |
-

6	Peter Butkovič, Hans Schneider, Sergeï Sergeev. "Recognizing Weakly Stable Matrices", SIAM Journal on Control and Optimization, 2012 Publication	1 %
7	docplayer.info Internet Source	1 %
8	Brunsch, Thomas, Laurent Hardouin, and Jörg Raisch. "Modelling Manufacturing Systems in a Dioid Framework", Industrial Information Technology, 2014. Publication	1 %
9	Rainer Burkard, David Shmoys, Uwe Zimmermann. "Combinatorial Optimization", Oberwolfach Reports, 2005 Publication	1 %
10	download.atlantis-press.com Internet Source	1 %
11	docplayer.net Internet Source	1 %
12	S. H. Nasser, J. L. Verdegay, F. Mahmoudi. "A New Method to Solve Fuzzy Interval Flexible Linear Programming Using a Multi-Objective Approach", Fuzzy Information and Engineering, 2021 Publication	1 %

13	P. Butkovič. "Permuted max-algebraic eigenvector problem is NP-complete", Linear Algebra and its Applications, 2008 Publication	1 %
14	Kendall Atkinson. "Linear Spaces", Texts in Applied Mathematics, 2009 Publication	<1 %
15	jims-a.org Internet Source	<1 %
16	mafiadoc.com Internet Source	<1 %
17	pages.mtu.edu Internet Source	<1 %
18	web.mat.bham.ac.uk Internet Source	<1 %
19	Butkovic, P.. "Permuted max-algebraic eigenvector problem is NP-complete", Linear Algebra and Its Applications, 20080415 Publication	<1 %
20	repository.ub.ac.id Internet Source	<1 %
21	Bart De Schutter, Bart De Moor. "The QR Decomposition and the Singular Value Decomposition in the Symmetrized Max-Plus Algebra Revisited", SIAM Review, 2002 Publication	<1 %

22

Butkovic, P.. "Max-algebra: the linear algebra of combinatorics?", Linear Algebra and Its Applications, 20030701

Publication

<1 %

23

Submitted to The University of Manchester

Student Paper

<1 %

24

math.sharif.edu

Internet Source

<1 %

25

Zhen Huang. "On time transfer in X-ray pulsar navigation", Science in China Series E Technological Sciences, 05/2009

Publication

<1 %

26

www.lyndhurstschools.net.schools.bz

Internet Source

<1 %

27

"Modelling and Performance Analysis of Cyclic Systems", Springer Science and Business Media LLC, 2020

Publication

<1 %

28

Juan Pablo Ibieta Jimenez. "Entropia topológica de emaranhamento em teorias de Higher Gauge Abelianas", Universidade de Sao Paulo, Agencia USP de Gestao da Informacao Academica (AGUIA), 2019

Publication

<1 %

29

coek.info

Internet Source

<1 %

30	Submitted to Loughborough University Student Paper	<1 %
31	id.123dok.com Internet Source	<1 %
32	Marianne Akian. "Topologies on lattice ordered groups, separation from closed downward sets and conjugations of type Lau", Optimization, 12/1/2003 Publication	<1 %
33	Submitted to Western Governors University Student Paper	<1 %
34	www.hf.uio.no Internet Source	<1 %
35	"Computer Aided Verification", Springer Science and Business Media LLC, 2010 Publication	<1 %
36	Woojin Kim, Tae-Sun Yu, Tae-Eog Lee. "Integrated Scheduling of a Dual-Armed Cluster Tool for Maximizing Steady Schedule Patterns", IEEE Transactions on Systems, Man, and Cybernetics: Systems, 2021 Publication	<1 %
37	journalofinequalitiesandapplications.springeropen.com Internet Source	<1 %
38	www.emis.de Internet Source	<1 %

39

A. F. Ivanov, A. N. Sharkovsky. "Chapter 5 Oscillations in Singularly Perturbed Delay Equations", Springer Science and Business Media LLC, 1992

Publication

<1 %

40

Daniel Belitiță, Karl-Hermann Neeb. "Nonlinear Completely Positive Maps and Dilation Theory for Real Involutive Algebras", Integral Equations and Operator Theory, 2015

Publication

<1 %

41

Leela-apiradee, Worrawate, Phantipa Thipwiwatpotjana, and Artur Gorka. "Closed form of L-localized solution set of max-plus interval linear system and its application on optimization problem", Journal of Computational and Applied Mathematics, 2017.

Publication

<1 %

42

Monika Molnárová. "Generalized matrix period in max-plus algebra", Linear Algebra and its Applications, 2005

Publication

<1 %

43

Semirings and Affine Equations over Them Theory and Applications, 2003.

Publication

<1 %

44

T. Brunsch, J. Raisch. "Modeling and control of high-throughput screening systems in a max-

<1 %

45 Vladimir Müller, Aljoša Peperko. "On the spectrum in max algebra", Linear Algebra and its Applications, 2015
Publication

46 K. Tebani, S. Amari, R. Kara. "Closed-loop control of constrained discrete event systems: application to a networked automation system", The International Journal of Advanced Manufacturing Technology, 2016
Publication

47 Lothar Seybold, Marcin Witczak, Paweł Majdzik, Ralf Stetter. "Towards Robust Predictive Fault-Tolerant Control for a Battery Assembly System", International Journal of Applied Mathematics and Computer Science, 2015
Publication

48 P. Butkovi. "Reducible Spectral Theory with Applications to the Robustness of Matrices in Max-Algebra", SIAM Journal on Matrix Analysis and Applications, 2010
Publication

49 Pierre Guillon, Zur Izhakian, Jean Mairesse, Glenn Merlet. "The ultimate rank of tropical matrices", Journal of Algebra, 2015

50 Steven Delvaux, Maurice Duits. "An Equilibrium Problem for the Limiting Eigenvalue Distribution of Rational Toeplitz Matrices", SIAM Journal on Matrix Analysis and Applications, 2010

Publication

51 Submitted to University of Warwick

Student Paper

52 W. D. Wallis. "Chapter 8 Matrices", Springer Science and Business Media LLC, 2003

Publication

53 Xavier Allamigeon. "Inferring Min and Max Invariants Using Max-Plus Polyhedra", Lecture Notes in Computer Science, 2008

Publication

54 Y. Shang. "Robust invariance in uncertain discrete event systems with applications to transportation networks", IMA Journal of Mathematical Control and Information, 06/01/2010

Publication

55 era.ed.ac.uk

Internet Source

56 Z A Banaszak. "Design of a performance repetitive manufacturing system", Proceedings of the Institution of Mechanical

57

Rob M.P. Goverde. "A delay propagation
algorithm for large-scale railway traffic
networks", Transportation Research Part C:
Emerging Technologies, 2010

Publication

<1 %

Exclude quotes On

Exclude matches < 5 words

Exclude bibliography On