

Strongly regular matrices and simple image set in interval max-plus algebra

<u>Siswanto</u>^{a, b} ⊠ ; <u>Suparwanto, Ari</u>^b ⊠ ; <u>Rudhito, M. Andy</u>^c ⊠

🖪 Save all to author list

^a Mathematics Department, Sebelas Maret University, Indonesia

^b Mathematics Department, Gadjah Mada University, Indonesia

^c Sanata Dharma University, Indonesia

2 28th percentile Citations in Scopus 18 Views count ⑦ ↗

View all metrics >

Full text options 🗸 🛛 🔂 Export

Abstract

Author keywords

SciVal Topics

Metrics

Abstract

Suppose that R is the set of real numbers and $R_{\varepsilon} = RU \{\varepsilon\}$ with $\varepsilon = -\infty$. Max-plus algebra is the set R_{ε} that is equipped with two operations, maximum and addition. Matrices over max-plus algebra are matrices whose elements belong to R_{ε} . The set (Formula presented). Which is equipped with maximum and addition operations is known as interval max-plus algebra. Matrices over interval max-

Related documents

Closure of the simple image set of linear mapping interval maxplus

Siswanto , Kurniawan, V.Y. , Pangadi (2020) Journal of Discrete Mathematical Sciences and Cryptography

Robust matrices in the interval max-plus algebra

Siswanto , Pangadi , Wiyono, S.B. (2019) Journal of Physics: Conference Series

The solution sets of maxalgebraic linear equation systems

Li, S.-J. , Wang, H.-L. (2019) Journal of Intelligent and Fuzzy Systems

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

AH

plus algebra are matrices whose elements belong to $I(R_{\epsilon})$. The research about image set. strongly regular matrix and simple image set in max-plus algebra have been done. In this paper, we investigate image set strongly regular matrix and simple image set in interval max-plus algebra. © 2016 Pushpa Publishing House, Allahabad, India.

Author keywords

Image set; Simple image set; Strongly regular matrix

SciVal Topics ()	~
Metrics	~

References (8)	View in search results format $ ightarrow$
 All CSV export ~ Print E-mail Create bibliography ¹ Baccelli, F., Cohen, G., Older, J.G., Quad (2001) Synchronization and Linearity. Co John Wiley & Sons, New York 	習 Save to PDF dat, J.P. ited 24 times.
 2 Butkovič, P. Strong regularity of matrices - a (1994) Discrete Applied Mathematics, 44 doi: 10.1016/0166-218X(92)00104-T View at Publisher 	a survey of results (<mark>Open Access</mark>) 8 (1), pp. 45-68. Cited 40 times.
 Butkovic, P. Simple image set of (max,+) line (2000) Discrete Applied Mathematics, 10 doi: 10.1016/S0166-218X(00)00212-2 View at Publisher 	ear mappings (<mark>Open Access</mark>) 05 (1-3), pp. 73-86. Cited 22 times.
 4 Butkovič, P. Max-algebra: The linear algebra (2003) Linear Algebra and Its Application doi: 10.1016/S0024-3795(02)00655-9 View at Publisher 	of combinatorics? (Open Access) <i>ns</i> , 367, pp. 313-335. Cited 106 times.
5 Butkovic, P.	

(2010) *Max Linear Systems: Theory and Algorithm*. Cited 345 times. Springer, London

6 Rudhito, M.A.

(2011) Fuzzy Number Max-Plus Algebra and Its Application to Fuzzy Scheduling and Queuing Network Problems, Doctoral Program of Mathematics, Faculty of Mathematics and Natural Sciences. Cited 2 times. Gadjah Mada University, Indonesian)

🗌 7 Tam, K.P.

(2010) *Optimizing and Approximating Eigenvectors in Max Algebra* (*Thesis*). Cited 13 times. A Thesis Submitted to The University of Birmingham for the Degree of Doctor of Philosophy (Ph.D

 8 Siswanto, A., Suparwanto, M.A.
 (2014) Rudhito, Existence of Solution to the System of Linear Equations in Interval Max-Plus Algebra, Proceedings of the National Conference of Mathematics XVII
 Institute of Technology Surabaya, Indonesian)

© Copyright 2016 Elsevier B.V., All rights reserved.

< Back to results | < Previous 11 of 12 Next >

 \checkmark Top of page

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

日本語に切り替える **切換到简体中文** 切換到繁體中文

Русский язык

Customer Service

Help Tutorials Contact us

ELSEVIER

Terms and conditions \neg Privacy policy \neg

Copyright © Elsevier B.V ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

RELX

JP Journal of Algebra, Number Theory and Applications



JP Journal of Algebra, Number Theory and Applications

Discontinued in Scopus as of 2017

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
India Universities and research institutions in India	Mathematics Algebra and Number Theory	Pushpa Publishing House	10
PUBLICATION TYPE	ISSN	COVERAGE	
Journals	09725555	2010-2017	
SCOPE			
Information not localized			
	${ig Q}$ Join the conversatio	on about this journal	

Quartiles

 \sim

B



See why IDC MarketScape named Adobe an e-signature leader.



4/11/22, 8:39 AM

JP Journal of Algebra, Number Theory and Applications



Get it

Metrics based on Scopus® data as of April 2021

A Amran Dalloul 2 years ago

Dear,

I would like to inquire whether this Journal still Scopus or not. Thank you and Best Regards, Amran Dalloul

reply



Melanie Ortiz 2 years ago

Dear Amran,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on 11 June 2020. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day. Best Regards, SCImago Team

A Alaoui noureddine 3 years ago

Why this journal is cancelled while the citation window is still active. I understand that the date indicated by Scopus, the journal is no longer indexed in its database. So I'm wondering if this journal is indexed or not. Thank you all. Sincerely

reply



Elena Corera 3 years ago

SCImago Team

SCImago Team

Dear Alaoui, thank you very much for your participation. Our data comes from Scopus/Elsevier, which offers an annual copy of their database. We understand that since the date indicated by Scopus/Elsevier the journal is no longer indexed in its database.



 \sim

See why IDC MarketScape named Adobe an e-signature leader.



Pushpa Publishing House

Reviewers | Book & Monographs | Conference | Contact Us

SEARCH

Home

Publication Ethics C

Guidelines Journals

Submit a Manuscript

P-ISSN: 0972-5555

Journal Menu

Journal Home

Editorial Board

Guidelines for Authors

Indexing

Contents

Subscribe

Publication Ethics and Publication Malpractice Statement

Categories

Artificial Intelligence

Biological and Agricultural

Chemistry and Materials

Computer and Communications

Economics and Business

Electrical, Energy and Physics

Life Sciences

.....

Mathematics

Oceanography and Marine Sci.

Statistics

Upcoming publications 2022

All Journals



Aims and Scope :

Call for Papers 2022

A Peer-Reviewed Open Access International Journal abbreviated JPANTA



The JP Journal of Algebra, Number Theory and Applications is a peer-reviewed international

journal. Original research papers theoretical, computational or applied, in nature, in any branch of Algebra and Number Theory are considered by the JPANTA. Together with the core topics in these fields along with their interplay, the journal promotes contributions in Diophantine Representation equations, theory. and Cryptography. Realising the need of wide range of information for any emerging area of potential research, the journal encourages the submission of related survey articles as well.

For preparing a manuscript in MS-Word: <u>Template</u>

Frequency :

Volume 53 (1) (2022), January

Volume 53 (2) (2022), February Volume 54 (2022), April

Volume 55 (2022), May Volume 56 (2022), July

Volume 57 (2022), August Volume 58 (2022), October

Volume 59 (2022), November An article after acceptance and complete formalities of publication will be put online immediately.



SUBMIT AN ARTICLE

Journal Stats

Publication count: 1020 Citation count: 2027 h10-index: 58 h-index: 17 Impact Factor: 1.73 Downloads: 294691 Views: 1322825 Downloads/article: 288.91 Citations/article: 1.99

Abstracting, Indexing and Reviews :



- Emerging Sources Citation Index (ESCI) (Clarivate Analytics: Web of Science)
- Zentralblatt MATH Database
- Mendeley (Elsevier Database)
- ProQuest Database is Now Part of Clarivate Analytics: Web of Science
- Pablons (Clarivate Analytics: Web of Science)
- ScienceGate Database
- Excellence in Research for Australia (ERA Journal 2018 ID: 41673)
- OTHER INDEXS

Impact Factor and CiteScore :

Google-based Journal Impact Factor: 1.73 Google-based CiteScore (Average Citations per Article): 3.02

Article Processing Charges (APC) :

An open-access journal provides wide range of accessibility to published articles and thus is beneficial to the contributors. However, on account of no subscriptions, etc., the journal suffers financially. To meet the publication standard, peer-review is necessary which requires a lot of correspondences. Also, proper composing and proof-reading costs are to be met. Maintenance of an efficient office is to be afforded. Therefore, we have the Article Processing Charge (APC) for an accepted paper. To defray the publication cost, authors are requested to arrange APC of their accepted papers at the rate of US\$ 30 per page. However, for authors in India this charge is Rs. 1000.00 per page.

Twenty five reprints together with a soft copy of a paper are provided to the corresponding author ex-gratis. Additional sets of reprints may be ordered at the time of proofs correction.

Editor-in-Chief :

Professor K. K. Azad, Retired Professor & Head, Department of Mathematics, Ex-Dean, Faculty of Science, University of Allahabad, INDIA

kkazad@pphmj.com https://scholar.google.co.in/citations? user=r2_5P_EAAAAJ&hl=en

Editor-in-Chief (Academic) :

Professor Istvan Gaal, Institute of Mathematics and Informatics, University of Debrecen, HUNGARY gaal.istvan @unideb.hu https://scholar.google.hu/citations? user=_EoPkcoAAAAJ&hl=en

Editorial Office :

Pushpa Publishing House, Vijaya Niwas 198, Mumfordganj, Prayagraj 211 002, INDIA jpanta@pphmj.com

Home Journals Books & Monographs Institutional Price List Refund Policy Disclaimer Policy Privacy Policy This website is best viewed at 1024x768 or higher resolution with Microsoft Internet Explorer 6 or newer. Shipping & Delivery Terms and Conditions

Copyright 2012



Research Interest 1. Ring and Module Theory 2. Homological Algebra

Professor Ercan Altınışık

Department of Mathematics, Gazi University, Turkey

ealtinisik@gazi.edu.tr https://avesis.gazi.edu.tr/ealtinisik

Research Interest 1. Linear Algebra

ONLINE SUBMISSIO

Professor Fujio Kubo

Department of Applied Mathematics, Hiroshima University, Japan remakubo@hiroshima-u.ac.jp https://www.researchgate.net/profile/Fujio-Kubo

Research Interest

1. Nonassociative Algebras

2. Algebraic Deformation Theory

ONLINE SUBMISSION

Professor Giulio Peruginelli

Department of Mathematics, University of Padova, Italy gperugin@math.unipd.it https://www.math.unipd.it/~gperugin/SitoWeb/cvItaPeruginelli.pdf

Research Interest

- 1. Commutative Algebra
- 2. Commutative Rings
- 3. Algebraic Number Theory

ONLINE SUBMISSION

Professor Ik Pyo Kim

Department of Mathematics Education, Daegu University, South Korea kimikpyo@daegu.ac.kr https://daegu.academia.edu/IkPyoKim

Research Interest

1. Linear Algebra

ONLINE SUBMISSION

Professor Istvan Gaal [Editor-in-Chief (Academic)]

Institute of Mathematics and Informatics, University of Debrecen, Hungary gaal.istvan @unideb.hu

 $https://scholar.google.hu/citations?user=_EoPkcoAAAAJ\&hl=en$

Research Interest

- 1. Number Theory
- 2. Commutative Ring
- 3. Algebraic Geometry
- 4. Matrix Theory

ONLINE SUBMISSION

Professor Jiyou Li

School of Mathematical Sciences, Shanghai Jiao Tong University, China lijiyou@sjtu.edu.cn https://scholar.google.com/citations?user=DGzjICEAAAAJ&hl=en

Research Interest

- 1. Combinatorial and Algebraic Number Theory
- 2. Coding Theory
- 3. Cryptography
- 4. Algebraic Combinatorics in particular enumerative combinatorics

ONLINE SUBMISSION

Professor Jose Rios Montes

Instituto de Matematicas, Universidad Nacional Autonoma de Mexico, Mexico jrios@matem.unam.mx

http://matematicos.matem.unam.mx/matematicos-r-z/matematicos-r/jose-rios

Research Interest

1. Ring and Module Theory

ONLINE SUBMISSION

Professor Kiyoshi Shirayanagi

Department of Information Science, Toho University, Japan kiyoshi.shirayanagi@is.sci.toho-u.ac.jp, shirayan@acm.org https://www.lab2.toho-u.ac.jp/sci/is/shirayanagi/index_en.html

Research Interest

- 1. Algebraic structures
- 2. Algorithm stabilization
- 3. Numeric and Symbolic computation

ONLINE SUBMISSION

Professor Liangyun Chen

School of Mathematics and Statistics, Northeast Normal University, China chenly640@nenu.edu.cn

Research Interest

- 1. Lie algebra
- 2. Lie superalgebra
- 3. Novkov algebra
- 4. Lie triple system

ONLINE SUBMISSION

Professor Mihran Papikian

Department Mathematics, Pennsylvania State University, USA papikian@psu.edu http://personal.psu.edu/mup17/

Research Interest 1. Algebraic Number Theory 2. Arithmetic Geometry

ONLINE SUBMISSION

Professor Qing-Wen Wang

Department of Mathematics, Shanghai University, China wqw@staff.shu.edu.cn https://www.scicol.shu.edu.cn/PDF/WangQingWen.htm

Research Interest1. Linear Algebra2. Matrix Theory3. Operator Algebra

ONLINE SUBMISSION

Professor Selvaraj Chelliah

Department of Mathematics, Periyar University, India selvavlr@yahoo.com https://periyaruniversity.irins.org/profile/43452

- Research Interest
- 1. Ring Theory
- 2. Homological Algebra
- 3. Representation Theory

Professor Taras Goy

Faculty of Mathematics and Computer Science, Vasyl Stefanyk Precarpathian National University, Ukraine taras.goy@pnu.edu.ua https://kdrpm.pnu.edu.ua/en/goy-taras/

Research Interest 1. Number Theory

- 2. Linear and Multilinear Algebra
- 3. Matrix Theory

online ocorrisoron

Professor Yang Zhang

Department of Mathematics, University of Manitoba, Canada yang.zhang@umanitoba.ca https://server.math.umanitoba.ca/~yangzhang/yangzhang/Research_interest.html

Research Interest

- 1. Ring and Module Theory
- 2. Noncommutative rings
- 3. Tensors
- 4. Matrix Theory
- 5. Hopf Algebra

ONLINE SUBMISSION

Professor Yangming Li

Department of Mathematics, Guangdong University of Education, China liyangming@gdei.edu.cn

Research Interest 1. Group Theory and Generalizations

ONLINE SUBMISSIO

Professor Yong-Gao Chen

School of Mathematical Sciences, Nanjing Normal University, China ygchen@njnu.edu.cn http://schools.njnu.edu.cn/math/person/yonggao-chen

- Research Interest
- Combinatorial Number Theory
 Analytic Number Theory

Professor Zhixiong Chen

School of Mathematics, Putian University, China ptczx@126.com https://www.researchgate.net/profile/Zhixiong-Chen-8

Research Interest

- 1. Cryptography-Algebraic
- 2. Number Theoretic
- 3. Pseudorandom Number Generators
- 3. Pseudorandom Sequences and Pseudorandomness
- 4. (Hyper-) Elliptic Curve Cryptography

ONLINE SUBMISSION

Professor Zhongshan Li

Department of Mathematics and Statistics, Georgia State University, USA zli@gsu.edu

Pushpa Publishing House

Research Interest	
1. Combinatorial Matrix Theory	
2. Sign Pattern Matrices	
3. Oriented Matroids	
4. Convex Polytopes	
5. Algebraic Graph Theory	

Home Journals Books & Monographs Institutional Price List Refund Policy Disclaimer Policy Privacy Policy Shipping & Delivery Terms and Conditions This website is best viewed at 1024x768 or higher resolution with Microsoft Internet Explorer 6 or newer.

Copyright 2012

Pushpa Publishing House

Reviewers | Book & Monographs | Conference | Contact Us

SEARCH

9

Home

Publication Ethics

Guidelines Jo

Journals Submit a

Submit a Manuscript

Journal Menu

Journal Home

Editorial Board

Guidelines for Authors

Indexing

Contents

Subscribe

Publication Ethics and Publication Malpractice Statement

Content

 E Volume 54 (2022) E Volume 53 (2022) E Volume 52 (2021) E Volume 51 (2021) E Volume 50 (2021) E Volume 49 (2021) E Volume 48 (2020) E Volume 47 (2020)
 E Volume 46 (2020) E Volume 45 (2020) E Volume 44 (2019)
 E Volume 43 (2019)
 E Volume 42 (2019) E Volume 41 (2019)
 E Volume 40 (2018) E Volume 39 (2017) ■ Volume 38 (2016)

JP Journal of Algebra, Number

Volume 38, Issue 1, Pages 1 -108 (February 2016)

Articles 1 - 8 of 8 [1]

INVERSE LIMIT OF LOCAL HOMOLOGY by: V. H. Jorge Pérez and C. H. Tognon JP Journal of Algebra, Number Theory and Applications Volume 38, Issue 1, Pages 1 - 18 (February 2016) Downloads: 391 Views: 1058

ON THE DIOPHANTINE

EQUATION $4q^x + 7^y = z^{2m}$ by: Somchit Chotchaisthit JP Journal of Algebra, Number Theory and Applications Volume 38, Issue 1, Pages 19 - 28 (February 2016) Downloads: 411 Views: 1224

THE REFINEMENT OF THE



METHOD by: Sawian Jaidee JP Journal of Algebra, Number Theory and Applications Volume 38, Issue 1, Pages 29 - 37 (February 2016) Downloads: 393 Views: 1320





LATEST ISSUE SUBMIT AN ARTICLE SEARCH WITHIN JOURNALS

Journal Stats

Publication count: 1020 Citation count: 2027 h10-index: 58 h-index: 17 Impact Factor: 1.73 Downloads: 294691 Views: 1322825 Downloads/article: 288.91 Citations/article: 1.99

Ξ	Volume 38, Issue 6	ST
	Pg 535 - 676 (December	ST
_	2016)	by: IP
Ξ	Volume 38, Issue 5	JI Th
	2016)	Vo
Ξ	Volume 38, Issue 4	<u>(Fe</u>
_	Pg 319 - 430 (August 2016)	Do
Ξ	Volume 38, Issue 3	AN
	Pg 213 - 318 (June 2016)	TH
=	Volume 38, Issue 2	AN
_	Pg 109 - 211 (April 2016)	РО
	Pa 1 - 108 (February 2016)	by
÷ 1	Volume 37 (2015)	He
F V	Volume 36 (2015)	Rie
	Volume 25 (2013)	JP Th
	Volume 33 (2014)	Vo
+ \	Volume 34 (2014)	<u>(Fe</u>
+ V	Volume 33 (2014)	Do
∓ \	Volume 32 (2014)	ST
+ 1	Volume 31 (2013)	M
+ 1	Volume 30 (2013)	SE
+ 1	Volume 29 (2013)	AL
÷ 1	Volume 28 (2013)	by: M
÷ 1	Volume 27 (2012)	JP
÷ 1	Volume 26 (2012)	Th
∓ \	Volume 25 (2012)	<u>Vo</u>
	Volume 24 (2012)	<u>(Fe</u> Do
	<i>k</i> -hanne 22 (2012)	
± `	volume 23 (2011)	ON
+ \	Volume 22 (2011)	PR
+ \	Volume 21 (2011)	KA FO
+ 7	Volume 20 (2011)	VA
+ 1	Volume 19 (2010)	by
÷ 1	Volume 18 (2010)	JP
+ 1	Volume 17 (2010)	1h Vo
÷ 1	Volume 16 (2010)	<u>(Fe</u>
⊞ \	Volume 15 (2009)	Do
V	Volume 14 (2009)	
	<i>L</i> -L	LE LE
+	volume 13 (2009)	

STRONGLY SEMIPRIME AND
STRONGLY NILPOTENT IDEALS
by: Ahmad Khaksari
JP Journal of Algebra, Number
Theory and Applications
Volume 38, Issue 1, Pages 39 - 43
<u>(February 2016)</u>
Downloads: 381 Views: 1206

AN ALGEBRAIC APPROACH TO THE NUMBER OF SOME ANTICHAINS IN THE

POWERSET 2ⁿ

by: AgustÃn Moreno Cañadas, Hernán Giraldo and Gabriel Bravo Rios JP Journal of Algebra, Number Theory and Applications <u>Volume 38, Issue 1, Pages 45 - 62</u> (February 2016) Downloads: 391 Views: 1229

TRONGLY REGULAR

MATRICES AND SIMPLE IMAGE SET IN INTERVAL MAX-PLUS ALGEBRA by: Siswanto, Ari Suparwanto and M. Andy Rudhito IP Journal of Algebra, Number Theory and Applications Volume 38, Issue 1, Pages 63 - 78 February 2016) Downloads: 408 Views: 1294

ON THE LOCAL AND GLOBAL PRINCIPLE FOR SYSTEMS OF RATIONAL HOMOGENEOUS FORMS IN A FINITE NUMBER OF VARIABLES

by: Lan Nguyen IP Journal of Algebra, Number Theory and Applications <u>Volume 38, Issue 1, Pages 79 - 94</u> (<u>February 2016)</u> Downloads: 398 Views: 1221

LEFT CONJUGACY CLOSED LEFT QUASIGROUPS WITH

4/11/22, 8:22 AM	Pushpa Publishing House		
E Volume 12 (2008)	PAIRWISE DISTINCT LEFT		
	TRANSLATIONS by: Juha Partala		
	JP Journal of Algebra, Number		
	Theory and Applications		
	<u>Volume 38, Issue 1, Pages 95 - 108</u> (February 2016)		
E Volume 7 (2007)	Downloads: 401 Views: 980		
	BACK		
E Volume 5 (2005)	BACK		
E Volume 3 (2003)			
E Volume 2 (2002)			
E Volume 1 (2001)			

HomeJournalsBooks & MonographsInstitutional Price ListRefund PolicyDisclaimer PolicyThis website is best viewed at 1024x768 or higher resolution with Microsoft Internet Explorer 6 or newer.Privacy PolicyShipping & DeliveryTerms and Conditions

Copyright 2012



STRONGLY REGULAR MATRICES AND SIMPLE IMAGE SET IN INTERVAL MAX-PLUS ALGEBRA

Siswanto^{1,2}, Ari Suparwanto² and M. Andy Rudhito³

¹Mathematics Department Faculty of Mathematics and Natural Sciences Sebelas Maret University Indonesia e-mail: sis.mipauns@yahoo.co.id

²Mathematics Department Faculty of Mathematics and Natural Sciences Gadjah Mada University Indonesia e-mail: ari_suparwanto@yahoo.com

³The Study Program in Mathematics Education Faculty of Teacher Training and Education Sanata Dharma University Indonesia e-mail: arudhito@yahoo.co.id

Abstract

Suppose that $\mathbb R$ is the set of real numbers and $\mathbb R_\epsilon=\mathbb R\bigcup\{\epsilon\}$ with

 $\epsilon=-\infty.$ Max-plus algebra is the set \mathbb{R}_ϵ that is equipped with two

operations, maximum and addition. Matrices over max-plus algebra

Keywords and phrases: image set, strongly regular matrix, simple image set. Communicated by K. K. Azad

Received: September 7, 2015; Revised: September 25, 2015; Accepted: October 9, 2015 2010 Mathematics Subject Classification: 15A15, 05C50, 15A30.

are matrices whose elements belong to \mathbb{R}_{ε} . The set $I(\mathbb{R})_{\varepsilon} = \{x = [\underline{x}, \overline{x}]\underline{x}, \overline{x} \in \mathbb{R}, \varepsilon < \underline{x} \le \overline{x}\} \cup \{\varepsilon\}$ with $\varepsilon = [\varepsilon, \varepsilon]$ which is equipped with maximum and addition operations is known as interval max-plus algebra. Matrices over interval max-plus algebra are matrices whose elements belong to $I(\mathbb{R}_{\varepsilon})$. The research about image set, strongly regular matrix and simple image set in max-plus algebra have been done. In this paper, we investigate image set, strongly regular matrix and simple image set in interval max-plus algebra.

1. Introduction

According to Baccelli et al. [1], max-plus algebra has been applied for modelling and analyzing problems in the area of planning, communication, production, queuing system with finite capacity, parallel computation and traffic. Max-plus algebra is the set of $\mathbb{R}_{\varepsilon} = \mathbb{R} \cup \{\varepsilon\}$ equipped with two operations, \otimes (maximum) and \otimes (addition), where \mathbb{R} is the set of real numbers with $\varepsilon = -\infty$ (Baccelli et al. [1]). According to Tam [7], max-plus algebra is a linear algebra over semiring \mathbb{R}_{ε} that is equipped with \oplus (maximum) and \otimes (addition) operations, while min-plus algebra is a linear algebra over semiring $\mathbb{R}_{\varepsilon'} = \mathbb{R} \bigcup \{\varepsilon'\}, \ \varepsilon' = \infty$ equipped with \oplus' (minimum) and \otimes' (addition) operations. Tam [7] also defined complete max-plus algebra and complete min-plus algebra. Complete max-plus algebra is a linear algebra over semiring $\mathbb{R}_{\varepsilon,\varepsilon'} = \mathbb{R}_{\varepsilon} \bigcup \{\varepsilon'\}$ equipped with \oplus (maximum) and \otimes (addition) operations, while complete min-plus algebra is a linear algebra over semiring $\mathbb{R}_{\epsilon',\epsilon} = \mathbb{R}_{\epsilon'} \bigcup \{\epsilon\}$ equipped with \oplus' (minimum) and \otimes' (plus) operations. Following the definition of max-plus algebra given by Baccelli et al. [1], we define min-plus algebra is the set of $\mathbb{R}_{\epsilon'}$ equipped with \oplus' (minimum) and \otimes' (plus) operations. Besides that, we define complete max-plus algebra is the set $\mathbb{R}_{\varepsilon,\varepsilon'} = \mathbb{R}_{\varepsilon} \bigcup \{\varepsilon'\}$ equipped with \oplus (maximum) and \otimes (addition) operations, while complete min-plus algebra is the set $\mathbb{R}_{\varepsilon',\varepsilon} = \mathbb{R}_{\varepsilon'} \cup \{\varepsilon\}$ equipped with \oplus' (minimum) and \otimes' (plus) operations. It shows that $\mathbb{R}_{\varepsilon,\varepsilon'} = \mathbb{R}_{\varepsilon} \cup {\varepsilon'} = \mathbb{R} \cup {\varepsilon, \varepsilon'} = \mathbb{R}_{\varepsilon'} \cup {\varepsilon'} = \mathbb{R}_{\varepsilon',\varepsilon}$. Furthermore, $\mathbb{R}_{\varepsilon,\varepsilon'}$ and $\mathbb{R}_{\varepsilon',\varepsilon}$ are written as \mathbb{R} . It can be formed a set of matrices in the size $m \times n$ of the sets \mathbb{R}_{ε} , $\mathbb{R}_{\varepsilon'}$, and \mathbb{R} . The set of matrices with components in \mathbb{R}_{ε} is denoted by $\mathbb{R}_{\varepsilon}^{m \times n}$. The matrix $A \in \mathbb{R}_{\varepsilon}^{m \times n}$ is called *matrix over max-plus algebra*. The matrix over min-plus algebra, over complete maxplus algebra and over min-plus algebra can also be defined (Tam [7]).

In 2010, Tam [7] discussed the linear system in max-plus algebra, image set and strongly regular matrix. Tam [7] and Butkovic [2, 3] mentioned that the solution of the linear system in max-plus algebra is closely related to image set, simple image set and strongly regular matrix. Butkovic [3-5] have shown the characteristics of the strongly regular matrix, that is matrix has strong permanent.

Rudhito [6] has generalized max-plus algebra into interval max-plus algebra and fuzzy number max-plus algebra. Moreover, Siswanto et al. [8] have discussed linear equation in interval max-plus algebra. Based on the discussion on the max-plus algebra that is connected to the system of linear equation, in this paper we discuss image set, strongly regular matrix and simple image set in interval max-plus algebra. Furthermore, it will also discuss matrix has strong permanent. The matrix has strong permanent is the characteristic of strongly regular matrix.

Before discussing the result of the research, the necessary concepts are discussed. These are interval max-plus algebra, system of linear equation in max-plus algebra and system of linear equation and the system of linear inequation in interval max-plus algebra.

Related to the system of linear equation $A \otimes x = b$, Tam [7] defined the solution set of the system of linear equation and image set of matrix *A*.

Definition 1.1. Given $A \in \mathbb{R}_{\varepsilon}^{m \times n}$ and $b \in \mathbb{R}_{\varepsilon}^{m}$. The solution set of the system of linear equation $A \otimes x = b$ is $S(A, b) = \{x \in \mathbb{R}_{\varepsilon}^{n} | A \otimes x = b\}$ and the image set of matrix A is $Im(A) = \{A \otimes x | x \in \mathbb{R}_{\varepsilon}^{n}\}$.

Based on the above definition, the following theorem and corollary are given.

Theorem 1.2. Given $A \in \mathbb{R}_{\varepsilon}^{m \times n}$ and $b \in \mathbb{R}_{\varepsilon}^{m}$. Vector $b \in Im(A)$ if and only if $S(A, b) \neq \emptyset$.

Corollary 1.3. Given $A \in \mathbb{R}^{m \times n}_{\varepsilon}$ and $b \in \mathbb{R}^{m}_{\varepsilon}$. Vector $b \in Im(A)$ if and only if $A \otimes (A^* \otimes' b) = b$.

The following is the definitions of subspace and theorem which states that for $A \in R_{\varepsilon}^{m \times n}$, Im(A) is a subspace.

Definition 1.4. Suppose that $S \subseteq \mathbb{R}^n_{\varepsilon}$. If for each $x, y \in S$ and for each $\alpha \in \mathbb{R}_{\varepsilon'}$, $\alpha \otimes x \in S$ and $x \oplus y \in S$, then S is known as subspace of maxplus algebra.

Theorem 1.5. Suppose that $A \in \mathbb{R}^{m \times n}_{\varepsilon}$. If $\alpha, \beta \in \mathbb{R}$ and $u, v \in Im(A)$ then $\alpha \otimes u \oplus \beta \otimes v \in Im(A)$.

In max-plus algebra there are the definitions of strongly linearly independent and strongly regular matrix.

Definition 1.6. The vectors $A_1, ..., A_n \in \mathbb{R}^m_{\varepsilon}$ are said to be *linearly dependent* if one of those vectors can be expressed as a linear combination of other vectors. Otherwise $A_1, ..., A_n$ are considered linearly independent.

Definition 1.7. The vectors $A_1, ..., A_n \in \mathbb{R}^m_{\varepsilon}$ are said to be *strongly linearly independent* if there is $b \in \mathbb{R}^m$ as such so that b can be uniquely expressed as a linear combination of $A_1, ..., A_n$.

Definition 1.8. For the vectors $A_1, ..., A_n \in \mathbb{R}^m_{\varepsilon}$ which are strongly linearly independent, if m = n, then the matrix $A = (A_1, ..., A_n)$ is known as strongly regular.

67

Below is the definition of simple image set.

Definition 1.9. Given $A \in \mathbb{R}^{m \times n}_{\varepsilon}$. The simple image set of matrix A is defined as $S_A = \{b \in \mathbb{R}^m | A \otimes x = b \text{ has unique solution}\}.$

Theorem 1.10. Suppose $A \in \mathbb{R}^{m \times n}_{\varepsilon}$ is double \mathbb{R} -astic and $b \in \mathbb{R}^{m}$, then $|S(A, b)| \in \{0, 1, \infty\}$.

Definition 1.11. Suppose $A \in \mathbb{R}^{m \times n}_{\varepsilon}$, T(A) is defined as $\{|S(A, b)| | b \in \mathbb{R}^{m}\}$.

Theorem 1.12. If $A \in \mathbb{R}^{m \times n}_{\varepsilon}$ is double \mathbb{R} -astic, then T(A) is $\{0, \infty\}$ or $\{0, 1, \infty\}$.

Based on Definition 1.7, the column vectors of *A* are strongly linearly independent if and only if $1 \in T(A)$. In max-plus algebra, a matrix that has strong permanent is also defined (Butkovic [4]).

Definition 1.13. Given $A \in \mathbb{R}^{m \times n}_{\varepsilon}$ and $\pi \in P_n$, where P_n is the set of all permutation of $N = \{1, 2, ..., n\}$. $\omega(A, \pi)$ is defined as $\prod_{j \in n}^{\otimes} A_{j, \pi(j)}$ and the permanent of matrix A is maper $(A) = \sum_{\pi \in P_n}^{\oplus} \omega(A, \pi)$.

Tam [7] defined that a matrix has strong permanent as given below.

Definition 1.14. The set of all permutation is written as ap(A), and is defined as $ap(A) = \{\pi \in P_n | maper(A) = \omega(A, \pi)\}$. Matrix *A* is said to have *strong permanent* if |ap(A)| = 1.

Butkovic [4] and Tam [7] explained the connection between strong regular matrix and matrix has strong permanent as below.

Theorem 1.15. Suppose that $A \in \mathbb{R}^{m \times n}_{\varepsilon}$ is double \mathbb{R} -astic. Matrix A is strongly regular matrix if and only if matrix A has strong permanent.

In the following section, interval max-plus algebra, matrix over interval max-plus algebra and the system of linear equation in interval max-plus algebra is in (Rudhito [6], Siswanto et al. [8]) are presented.

Closed interval x in \mathbb{R}_{ε} is a subset of \mathbb{R}_{ε} in the form of $x = [\underline{x}, \overline{x}] = \{x \in \mathbb{R}_{\varepsilon} | \underline{x} \le x \le \overline{x}\}$. The interval x in \mathbb{R}_{ε} is known as *interval max-plus*. A number $x \in \mathbb{R}_{\varepsilon}$ can be expressed as interval [x, x].

Definition 1.16. $I(\mathbb{R})_{\varepsilon} = \{x = [\underline{x}, \overline{x}] | \underline{x}, \overline{x} \in \mathbb{R}, \varepsilon < \underline{x} \le \overline{x}\} \cup \{\varepsilon\}$ is formed, with $\varepsilon = [\varepsilon, \varepsilon]$. On the set $I(\mathbb{R})_{\varepsilon}$, the operations $\overline{\oplus}$ and $\overline{\otimes}$ with $x \overline{\oplus} y = [\underline{x} \oplus \underline{y}, \overline{x} \oplus \overline{y}]$ and $x \overline{\otimes} y = [\underline{x} \oplus \underline{y}, \overline{x} \oplus \overline{y}]$ for every $x, y \in I(\mathbb{R})_{\varepsilon}$ are defined. Furthermore they are known as *interval max-plus algebra* and notated as $I(\mathfrak{R})_{\max} = (I(\mathbb{R})_{\varepsilon}; \overline{\oplus}, \overline{\otimes})$.

Definition 1.17. The set of matrices in the size $m \times n$ with the elements in $I(\mathbb{R})_{\varepsilon}$ is notated as $I(\mathbb{R})_{\varepsilon}^{m \times n}$, namely:

$$I(\mathbb{R})_{\varepsilon}^{m \times n} = \{A = [A_{ij}] | A_{ij} \in I(\mathbb{R})_{\varepsilon}; \text{ for } i = 1, 2, ..., m \text{ and } j = 1, 2, ..., n\}.$$

Matrices which belong to $I(\mathbb{R})^{m \times n}_{\varepsilon}$ are known as *matrices over interval max*plus algebra or shortly interval matrices.

Definition 1.18. For $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$, matrix $\underline{A} = [\underline{A}_{ij}] \in \mathbb{R}^{m \times n}_{\varepsilon}$ and $\overline{A} = [\overline{A}_{ij}] \in \mathbb{R}^{m \times n}_{\varepsilon}$ are defined, as a lower bound matrix and an upper bound matrix of the interval matrix A, respectively.

Definition 1.19. Given that interval matrix $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$, and \underline{A} and \overline{A} are a lower bound matrix and an upper bound matrix of the interval matrix A, respectively. The matrix interval of A is defined, namely $[\underline{A}, \overline{A}] = \{A = \mathbb{R}^{m \times n}_{\varepsilon} | \underline{A} \leq A \leq \overline{A}\}$ and $I(\mathbb{R}^{m \times n}_{\varepsilon})_{b} = \{[\underline{A}, \overline{A}] | A \in (\mathbb{R})^{m \times n}_{\varepsilon}\}.$

Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $b = (b_1, ..., b_m) \in I(\mathbb{R})^m_{\varepsilon}$. The equation system $A \otimes x = b$ is known as the system of linear equation in interval

max-plus algebra. Furthermore, $A \approx [\underline{A}, \overline{A}]$, $x \approx [\underline{x}, \overline{x}]$ and $b \approx [\underline{b}, \overline{b}]$ with $[\underline{A}, \overline{A}] \in I(\mathbb{R}_{\varepsilon}^{n \times n})_{b}$, $[\underline{x}, \overline{x}], [\underline{b}, \overline{b}] \in I(\mathbb{R}_{\varepsilon}^{n})_{b}$. To find the solution of the system of linear equation in interval max-plus algebra, the solution of the system of linear equation in max-plus algebra $\underline{A} \otimes \underline{x} = \underline{b} \operatorname{dan} \overline{A} \otimes \overline{x} = \overline{b}$ should be determined.

Definition 1.20. Given the system of linear equation $A \otimes x = b$ with $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $b \in I(\mathbb{R})^m_{\varepsilon}$. $IS(A, b) = \{x \in I(\mathbb{R})^m_{\varepsilon} | A \otimes x = b\}$ is defined.

Along with the discussion the system of linear equation in max-plus algebra, without loss of generality, the system of linear equation in interval max-plus algebra $A \otimes x = b$ with $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$ is discussed.

Theorem 1.21. Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$. Vector $y \in IS(A, b)$ if and only if $\underline{y} \in S(\underline{A}, \underline{b}), \ \overline{y} \in S(\overline{A}, \overline{b})$ and $\underline{y} \leq \overline{y}$. Furthermore y can be expressed as $y \approx [\underline{y}, \overline{y}]$.

Each of the following two corollaries is a criterion of the system of linear equation which has solution and a criterion the system of linear equation which has unique solution. The notations of $M = \{1, 2, ..., m\}$ and $N = \{1, 2, ..., n\}$ are used to simplify.

Corollary 1.22. Suppose $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$, then the following three statements are equivalent:

a. $IS(A, b) \neq \emptyset$. b. $\hat{y} \in IS(A, b)$. c. $\bigcup_{j \in N} M_j(\underline{A}, \underline{b}) = M \text{ and } \bigcup_{j \in N} M_j(\overline{A}, \overline{b}) = M$. **Corollary 1.23.** Suppose $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$, then $IS(A, b) = \{\hat{y}\}$ if and only if

a.
$$\bigcup_{j \in N} M_j(\underline{A}, \underline{b}) = M \text{ and } \bigcup_{j \in N} M_j(\overline{A}, \overline{b}) = M.$$

b. $\bigcup_{j \in N'} M_j(\underline{A}, \underline{b}) \neq M$ for every $N' \subseteq N$, $N' \neq N$ and $\bigcup_{j \in N''} M_j(\overline{A}, \overline{b}) = M$ for every $N'' \subseteq N$, $N'' \neq N$.

2. Main Results

In this section, we present new results of the research. The results that are obtained, are related with the system of linear equation in interval max-plus algebra $A \otimes x = b$, $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $b \in I(\mathbb{R})^m$.

We will start the definition of the image set of matrix A.

Definition 2.1. Given $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$. The image set of matrix A is $IIm(A) = \{A \otimes x \mid x \in I(\mathbb{R})^n_{\varepsilon}\}.$

Based on the definition of the image set of a matrix, a theorem which demonstrated the necessary and sufficient conditions of the elements of the image set is presented.

Theorem 2.2. Given $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $b \in I(\mathbb{R})^m_{\varepsilon}$. Vector $b \in IIm(A)$ if and only if $IS(A, b) \neq \emptyset$.

Proof. Suppose that $A \approx [\underline{A}, \overline{A}]$ and $b \approx [\underline{b}, \overline{b}]$ with $\underline{A}, \overline{A} \in \mathbb{R}^{m \times n}_{\varepsilon}$ and $\underline{b}, \overline{b} \in \mathbb{R}^{m}_{\varepsilon}$. Since $b \in IIm(A)$, it means that $\underline{b} \in Im(\underline{A})$ and $\overline{b} \in Im(\overline{A})$. Therefore, based on Theorem 1.2, i.e., $S(\underline{A}, \underline{b}) \neq \emptyset$ and $S(\overline{A}, \overline{b}) \neq \emptyset$, then $IS(A, b) \neq \emptyset$. On the other hand, since $IS(A, b) \neq \emptyset$, $S(\underline{A}, \underline{b}) \neq \emptyset$ and $S(\overline{A}, \overline{b}) \neq \emptyset$ and $S(\overline{A}, \overline{b}) \neq \emptyset$. Thus, based on Theorem 1.2 that is $\underline{b} \in Im(\underline{A})$ and $\overline{b} \in Im(\overline{A})$, then $b \in IIm(A)$ with $b \approx [\underline{b}, \overline{b}]$. With regard to Theorem 2.2, the following corollary is obtained:

Corollary 2.3. Given $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $b \in I(\mathbb{R})^m_{\varepsilon}$. Vector $b \in IIm(A)$ if and only if $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$.

Proof. Suppose that $A \approx [\underline{A}, \overline{A}]$ and $b \approx [\underline{b}, \overline{b}]$. Since $b \in IIm(A)$, it means that $\underline{b} \in Im(\underline{A})$ and $\overline{b} \in Im(\overline{A})$. Based on Corollary 1.3, $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$. On the other hand, since $\underline{b} = \underline{A} \otimes (\underline{A}^* \otimes' \underline{b})$ and $\overline{b} = \overline{A} \otimes (\overline{A}^* \otimes' \overline{b})$, based on Corollary 1.3, $\underline{b} \in Im(\underline{A})$ and $\overline{b} \in Im(\overline{A})$ are obtained. Therefore, $b \approx [\underline{b}, \overline{b}] \in IIm(A)$. \Box

The following theorem presents that IIm(A) is a subspace of $I(\mathbb{R})^m_{\varepsilon}$.

Theorem 2.4. Given
$$A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$$
, $IIm(A)$ is a subspace of $I(\mathbb{R})^{m}_{\varepsilon}$

Proof. Since $IIm(A) = \{A \otimes x \mid x \in I(\mathbb{R})^n_{\mathcal{E}}\}$, $IIm(A) \subseteq I(\mathbb{R})^m_{\mathcal{E}}$ and $Im(A) \neq \emptyset$. Taking $p, q \in IIm(A)$ and $\alpha, \beta \in I(\mathbb{R})$, accordingly, $\alpha \otimes \overline{p} \otimes \overline{\beta} \otimes \overline{q} \approx [\underline{\alpha} \otimes \underline{p} \oplus \underline{\beta} \otimes \underline{q}, \overline{\alpha} \otimes \overline{p} \oplus \overline{\beta} \otimes \overline{q}]$ with $\underline{\alpha} \otimes \underline{p} \oplus \underline{\beta} \otimes \underline{q} \in Im(\underline{A})$ and $\overline{\alpha} \otimes \overline{p} \oplus \overline{\beta} \otimes \overline{q} \in Im(\overline{A})$. It means that $\alpha \otimes \overline{p} \oplus \overline{\beta} \otimes \overline{q} \in Im(A)$. In other words, $\alpha \otimes \overline{p} \oplus \overline{\beta} \otimes \overline{q} \in Im(A)$ and hence IIm(A) is a subspace of $I(\mathbb{R})^m_{\mathcal{E}}$.

Several definitions and theorems are presented to support the discussion on strongly regular matrix.

Definition 2.5. The vectors $A_{1,}, ..., A_n \in I(\mathbb{R})^m_{\varepsilon}$ is said to be *linearly dependent* if one of those vectors can be said to be as *linear combination* of other vectors. Otherwise then, $A_1, ..., A_n$ is said to be *linearly independent*.

Theorem 2.6. If the vectors $A_1, ..., A_n \in I(\mathbb{R})^m_{\varepsilon}$ with $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2], ..., A_n \approx [\underline{A}_n, \overline{A}_n]$ are linearly dependent, then $\underline{A}_1, \underline{A}_2, ..., \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, ..., \overline{A}_n$ are linearly independent. **Proof.** It is known that $A_1, ..., A_n \in I(\mathbb{R})^m_{\varepsilon}$ is linearly dependent. It means that one of those vectors can be stated as a linear combination of the others. Without losing the generality of the proof, for example $A_1 = \overline{\oplus}^n_{i=2}(\alpha_i \otimes A_i)$ with $\alpha_i \in I(\mathbb{R})_m$, since $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2]$, ..., $A_n \approx [\underline{A}_n, \overline{A}_n]$, $\underline{A}_1 = \bigoplus_{i=2}^n (\alpha_i \otimes \underline{A}_i)$ and $\overline{A}_1 = \bigoplus_{i=2}^n (\overline{\alpha}_i \otimes \overline{A}_i)$. In other words, $\underline{A}_1, \underline{A}_2, ..., \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, ..., \overline{A}_n$ are linearly dependent.

Definition 2.7. The vectors $A_1, ..., A_n \in I(\mathbb{R})^m_{\varepsilon}$ is said to be *strongly linearly independent* if there is $b \in I(\mathbb{R})^m$, b can be uniquely expressed as a linear combination of $A_1, ..., A_n$, that is $b = \overline{\bigoplus}_{i=1}^n (\alpha_i \otimes A_i), \alpha_i \in I(\mathbb{R})$.

The following is the definition of strongly regular matrix.

Definition 2.8. Matrix $A = (A_1, A_2 \dots A_n) \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is said to be *strongly regular* if the vectors $A_1, \dots, A_n \in I(\mathbb{R})^n_{\varepsilon}$ are strongly linearly independent.

Based on the definition of strongly linearly independent in the max-plus algebra, the following theorem is formulated.

Theorem 2.9. If the vectors $A_{1,}, ..., A_n \in I(\mathbb{R})^m_{\varepsilon}$ with $A_1 \approx [\underline{A}_1, \overline{A}_1]$, $A_2 \approx [\underline{A}_2, \overline{A}_2], ..., A_n \approx [\underline{A}_n, \overline{A}_n]$ are strongly linearly independent, then $\underline{A}_1, \underline{A}_2, ..., \underline{A}_n$ and $\overline{A}_1, \overline{A}_2, ..., \overline{A}_n$ are strongly linearly independent.

Proof. It is known that $A_1 \approx [\underline{A}_1, \overline{A}_1], A_2 \approx [\underline{A}_2, \overline{A}_2], ..., A_n \approx [\underline{A}_n, \overline{A}_n]$ are strongly linearly independent, then there is $b \approx [\underline{b}, \overline{b}] \in I(\mathbb{R})^m$ so that bcan be uniquely expressed as linear combination of $A_1, ..., A_n$. Suppose that $b = \overline{\bigoplus}_{i=1}^n (\alpha_i \otimes A_i)$; since

$$\overline{\oplus}_{i=1}^{n} (\alpha_{i} \overline{\otimes} A_{i}) \approx [\oplus_{i=1}^{n} (\underline{\alpha}_{i} \otimes \underline{A}_{i}),$$

 $\oplus_{i=1}^{n} (\overline{\alpha}_{i} \otimes \overline{A}_{i})], \quad \underline{b} = \oplus_{i=1}^{n} (\underline{\alpha}_{i} \otimes \underline{A}_{i}) \text{ and } \overline{b} = \oplus_{i=1}^{n} (\overline{\alpha}_{i} \otimes \overline{A}_{i}), \text{ It means that } \underline{b} \text{ can be uniquely expressed as a linear combination of } \underline{A}_{1}, \underline{A}_{2}, ..., \underline{A}_{n} \text{ and } \overline{b} \text{ can be uniquely expressed as a linear combination } \overline{A}_{1}, \overline{A}_{2}, ..., \overline{A}_{n}. \text{ In other words, } \underline{A}_{1}, \underline{A}_{2}, ..., \underline{A}_{n} \text{ and } \overline{A}_{1}, \overline{A}_{2}, ..., \overline{A}_{n} \text{ are strongly linearly independent.}$

The following is the definition of simple image set.

Definition 2.10. Given $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$. The set $IS_A = \{b \in I(\mathbb{R})^m \mid A \otimes x = b \text{ has a unique solution}\}$ is defined as a simple image set of matrix A.

The regularity properties of the matrix are closely related to the number of the solutions of the system of linear equation. It can be stated in the following definitions and theorems.

Theorem 2.11. Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ is a double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^m$ is the number of the elements of IS(A, b) is $|IS(A, b)| \in \{0, 1, \infty\}$.

Proof. Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^{m}$. Suppose that $A \approx [\underline{A}, \overline{A}]$ and $b \approx [\underline{b}, \overline{b}]$ with each of \underline{A} and \overline{A} is double \mathbb{R} -astic, while $\underline{b}, \overline{b} \in \mathbb{R}^{m}$. Based on the Theorem 1.10, $|S(\underline{A}, \underline{b})| \in \{0, 1, \infty\}$ and $|S(\overline{A}, \overline{b})| \in \{0, 1, \infty\}$. Therefore, following the Definition 1.18, it is confirmed that $|IS(A, b)| \in \{0, 1, \infty\}$.

Definition 2.12. Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$, T(A) is defined as $\{|IS(A, b)| | b \in \mathbb{R}^{m}\}.$

Theorem 2.13. If $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic, then $T(A) = \{0, \infty\}$ or $T(A) = \{0, 1, \infty\}$.

Proof. Suppose that $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$, $A \approx [\underline{A}, \overline{A}]$ with $\underline{A} \in \mathbb{R}^{m \times n}_{\varepsilon}$ and $\overline{A} \in \mathbb{R}^{m \times n}_{\varepsilon}$. Based on the theorem for the case of max-plus algebra, $T(\underline{A}) = \{0, 1, \infty\}$ or $T(\underline{A}) = \{0, \infty\}$ and $T(\overline{A}) = \{0, \infty\}$ or $T(\overline{A}) = \{0, 1, \infty\}$. Therefore, $T(A) = \{0, \infty\}$ or $(A) = \{0, 1, \infty\}$.

Based on Theorem 2.13, A is a strongly regular matrix if and only if $1 \in T(A)$. Then, the criteria of strongly regular matrix is discussed.

Definition 2.14. Given $A = (a_{ij}) \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ and $\pi \in P_n$, with P_n is the set of all permutations of $\{1, 2, ..., n\}$. $\omega(A, \pi)$ is defined as $\overline{\otimes}_{j \in n} a_{j, \pi(j)}$.

Theorem 2.15. If $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ double $I(\mathbb{R})$ -astic and $\pi \in P_n$ with $A \approx [\underline{A}, \overline{A}]$, then $\omega(A, \pi) = [\omega(\underline{A}, x) \omega(\overline{A}, \pi)]$.

Proof. If $A \in I(\mathbb{R})^{m \times n}_{\varepsilon}$ and $\pi \in P_n$ with $A \approx [\underline{A}, \overline{A}]$, it means that

$$\begin{split} \omega(A, \pi) &= \overline{\otimes}_{j \in n} a_{j, \pi(j)} = \overline{\otimes}_{j \in n} [\underline{a}_{j, \pi(j)}, \overline{a}_{j, \pi(j)}] \\ &= [\otimes_{j \in n} \underline{a}_{j, \pi(j)}, \otimes_{j \in n} \overline{a}_{j, \pi(j)}] = [\omega(\underline{A}, \pi), \omega(\overline{A}, \pi)]. \end{split}$$

Definition 2.16. The permanent of matrix A is defined as maper $\overline{\otimes}_{\pi \in P_n} \omega(A, \pi)$.

Next discussion is matrix over interval max-plus algebra which has weak permanent and strong permanent.

Definition 2.17. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \overline{A}]$. The set of all optimal permutations is ap(A) with $ap(A) = \{\pi \in P_n | \operatorname{maper}(\underline{A}) = \omega(\underline{A}, \pi) \text{ or maper}(\overline{A}) = \omega(\overline{A}, \pi) \}$. Matrix A is said to have *strong permanent* if |ap(A)| = 1 and it is said to have *weak permanent* if $|ap(\underline{A})| = 1$.

Theorem 2.18. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic. If matrix A has strong permanent, then matrix A has weak permanent.

Proof. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$, $A \approx [\underline{A}, \overline{A}]$ with $\underline{A} \in (\mathbb{R})^{n \times n}_{\varepsilon}$ and $\overline{A} \in \mathbb{R}^{n \times n}_{\varepsilon}$. Since *A* has strong permanent, it means that |ap(A)| = 1. Suppose that $\pi_1 \in ap(A)$, then maper $(\underline{A}) = \omega(\underline{A}, \pi_1)$ and maper $(\overline{A}) = \omega(\overline{A}, \pi_1)$. Therefore, $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$. In other words, *A* has weak permanent.

Theorem 2.19. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \overline{A}]$. Matrix A has weak permanent if and only if \underline{A} and \overline{A} have strong permanent.

Proof. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$, $A \approx [\underline{A}, \overline{A}]$ with $\underline{A} \in \mathbb{R}^{n \times n}_{\varepsilon}$ and $\overline{A} \in \mathbb{R}^{n \times n}_{\varepsilon}$. It is known that matrix A has weak permanent. It means that $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$. Therefore, matrix \underline{A} and \overline{A} have strong permanent. On the other hand, suppose that \underline{A} and \overline{A} have strong permanent, it means that $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$ and $|ap(\overline{A})| = A$. Therefore, matrix A with $A \approx [\underline{A}, \overline{A}]$ has weak permanent.

Theorem 2.20. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic with $A \approx [\underline{A}, \overline{A}]$. Matrix A has strong permanent if and only if \underline{A} and \overline{A} have strong permanent as such that if maper $(\underline{A}) = \omega(\underline{A}, \pi_1)$ and maper $(\overline{A}) = \omega(\overline{A}, \pi_2)$, then $\pi_1 = \pi_2$.

Proof. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ with $A \approx [\underline{A}, \overline{A}]$. It is known that matrix A has strong permanent. Since matrix A has strong permanent, it means that |ap(A)| = 1. Suppose that $\pi \in ap(A)$ it means that $\pi \in P_n$ so that maper $(\underline{A}) = \omega(\underline{A}, \pi)$ and maper $(\overline{A}) = \omega(\overline{A}, \pi)$. Thus, $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$. Therefore, \underline{A} and \overline{A} have strong permanent. On the other hand, it is known that matrix \underline{A} and \overline{A} have strong permanent. If maper $(\underline{A}) =$

 $\omega(\underline{A}, \pi_1)$ and maper $(\overline{A}) = \omega(\overline{A}, \pi_2)$, then $\pi_1 = \pi_2$. Therefore, $|ap(\underline{A})| = 1$ and $|ap(\overline{A})| = 1$ with $\pi_1 \in |ap(\underline{A})|, \pi_2 \in |ap(\overline{A})|$ and $\pi_1 = \pi_2$. Since $ap(A) = \{\pi \in P_n | \text{maper}(\underline{A}) = \omega(\underline{A}, \pi) \text{ or maper}(\overline{A}) = \omega(\overline{A}, \pi)\}, |ap(A)| = 1$. Therefore, matrix A has strong permanent.

The following is theorem which has the characteristics of strongly regular matrix.

Theorem 2.21. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic. Matrix *A* is strongly regular if and only if matrix *A* has strong permanent.

Proof. Suppose that $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ is double $I(\mathbb{R})$ -astic, with $A \approx [\underline{A}, \overline{A}]$. It is known that A is strongly regular matrix. Suppose that $A_1, ..., A_n$ are the columns of matrix A. Since A is strongly regular matrix, there is $b \in I(\mathbb{R})^n$ so that b can be uniquely expressed as linear combination of $A_{1,...,A_n}$, that is $b = \overline{\bigoplus}_{i=1}^n (x_i \otimes A_i)$, $x_i \in I(\mathbb{R})$. The equation $b = \overline{\bigoplus}_{i=1}^{n} (x_i \otimes A_i) \Leftrightarrow A \otimes x = b$, is the system of linear equation in interval max-plus algebra which has a unique solution. Based on Corollary 1.23, |ap(A)| = 1. Therefore, matrix A has strong permanent. The converse, it is known that matrix A has strong permanent. Based on Theorem 2.20, matrix \underline{A} and \overline{A} have such strong permanent, so that if maper $(\underline{A}) = \omega(\underline{A}, \pi_1)$ and maper $(\overline{A}) = \omega(\overline{A}, \pi_2)$, $\pi_1 = \pi_2$. Therefore, matrix A and \overline{A} are both strongly regular matrix, and thus, there are b and \overline{b} which can be uniquely written as the linear combinations $\underline{b} = \oplus_{i=1}^{n} (\underline{x}_{i} \otimes \underline{A}_{i}) \text{ and } \overline{b} = \oplus_{i=1}^{n} (\overline{x}_{i} \otimes \overline{A}_{i}), \text{ in which } \underline{A}_{1}, \underline{A}_{2}, ..., \underline{A}_{n} \text{ are}$ the columns of matrix \underline{A} , while \overline{A}_1 , \overline{A}_2 , ..., \overline{A}_n are the columns of matrix \overline{A} . That $\underline{b} = \bigoplus_{i=1}^{n} (\underline{x}_{i} \otimes \underline{A}_{i}) \Leftrightarrow \underline{A} \otimes \underline{x} = \underline{b}$ and $\overline{b} = \bigoplus_{i=1}^{n} (\overline{x}_{i} \otimes \overline{A}_{i}), \Leftrightarrow \overline{A} \otimes \overline{x} =$ \overline{b} . Thus, the system of linear equation $\underline{A} \otimes \underline{x} = \underline{b}$ and $\overline{A} \otimes \overline{x} = \overline{b}$ have

77

unique solution. Since maper(\underline{A}) = $\omega(\underline{A}, \pi_1)$ and maper(\overline{A}) = $\omega(\overline{A}, \pi_2)$, $\pi_1 = \pi_2$ so that $[\underline{b}, \overline{b}] \approx b$. As the result, $[\underline{A} \otimes \underline{x}, \overline{A} \otimes \overline{x}] = [\underline{b}, \overline{b}]$ and since $A \otimes x \approx [\underline{A} \otimes \underline{x}, \overline{A} \otimes \overline{x}]$, $A \otimes \overline{x} = b$ or $b = \overline{\bigoplus}_{i=1}^n (x_i \otimes A_i)$ with $A_{1,i}, ..., A_n$ as the columns of matrix A. It means that there is b which can be uniquely expressed as linear combination of $A_1, ..., A_n$. This matrix A is known as strongly regular matrix.

3. Concluding Remarks

Based on the discussion, the following conclusions are obtained:

1. The necessary and sufficient condition of matrix $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$, which is double $I(\mathbb{R})$ -astic is strongly regular matrix.

2. The system of linear equation in the interval max-plus algebra $A \otimes x = b$, $A \in I(\mathbb{R})^{n \times n}_{\varepsilon}$ which is double $I(\mathbb{R})$ -astic and $b \in I(\mathbb{R})^{n}$ has unique solution if and only if matrix *A* has strong permanent.

Acknowledgment

The authors thank the anonymous referees for their valuable suggestions which let to the improvement of the manuscript.

References

- [1] F. Baccelli, G. Cohen, J. G. Older and J. P. Quadat, Synchronization and Linearity, John Wiley & Sons, New York, 2001.
- [2] P. Butkovic, Strongly regularity of matrices a survey of results, Discrete Applied Mathematics 48 (1994), 45-68.
- [3] P. Butkovic, Simple image set of (max, +) linear mappings, Discrete Applied Mathematics 105 (2000), 73-86.
- [4] P. Butkovic, Max-algebra: the linear algebra of combinatorics?, Linear Algebra and Application 367 (2003), 313-335.

Siswanto, Ari Suparwanto and M. Andy Rudhito

- [5] P. Butkovic, Max Linear Systems: Theory and Algorithm, Springer, London, 2010.
- [6] M. A. Rudhito, Fuzzy number max-plus algebra and its application to fuzzy scheduling and queuing network problems, Doctoral Program of Mathematics, Faculty of Mathematics and Natural Sciences, Gadjah Mada University, 2011 (Indonesian).
- [7] K. P. Tam, Optimizing and approximating eigenvectors in max algebra (Thesis), A Thesis Submitted to The University of Birmingham for the Degree of Doctor of Philosophy (Ph.D.), 2010.
- [8] Siswanto, A. Suparwanto, M. A. Rudhito, Existence of solution to the system of linear equations in interval max-plus algebra, Proceedings of the National Conference of Mathematics XVII, Institute of Technology Surabaya, 2014 (Indonesian).

78