



Certificate of Appreciation

3876/UN1/FBI/KSA/PK/02.00/2021

is awarded to

Yustina Sri Hartini

in recognition of his/her contribution as

PRESENTER

at The 7th International Conference
on Biological Sciences (ICBS 2021)
Faculty of Biology, Universitas Gadjah Mada,
Yogyakarta, Indonesia, October 14th-15th, 2021

Organizing Committee
Chairperson

Dr. Miftahul Ilmi, M.Si.

Faculty of Biology,
Universitas Gadjah Mada,

Dean

Prof. Dr. Budi S. Daryono, M.Agr.Sc.



UNIVERSITAS GADJAH MADA
FACULTY OF BIOLOGY



Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

Search



Series: [Advances in Biological Sciences Research](#)

Proceedings of the 7th International Conference on Biological Science (ICBS 2021)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)

[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)



It is our pleasure to welcome you to the 7th International Conference on Biological Sciences 2021. This conference is a biennially event held by the Faculty of Biology, Universitas Gadjah Mada, in collaboration with the Faculty of Applied Science and Technology, Universiti Tun Hussein Onn Malaysia. This year the conference's theme is "Contributing to Global Issues: Current Advances in Biodiversity and Health Research in the Face of Climate Change". Along with the theme, we have 7 conference topics ranging widely from biosystematics to bioengineering.




Please click [here](#) for the conference website.

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

Home Privacy Policy Terms of use   

Copyright © 2006-2022 Atlantis Press – now part of Springer Nature

Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

Search



Series: [Advances in Biological Sciences Research](#)

Proceedings of the 7th International Conference on Biological Science (ICBS 2021)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)

[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)

It is our pleasure to welcome you to the 7th International Conference on Biological Sciences 2021. This conference is a biennially event held by the Faculty of Biology, Universitas Gadjah Mada, in collaboration with the Faculty of Applied Science and Technology, Universiti Tun Hussein Onn Malaysia.

This year the conference's theme is "Contributing to Global Issues: Current Advances in Biodiversity and Health Research in the Face of Climate Change". Along with the theme, we have 7 conference topics ranging widely from biosystematics to bioengineering.

This year's conference is also the first time held online due to the global pandemic situation. However, it becomes a blessing in disguise, because the

conference becomes accessible to a wider audience and participants from all over the world. The number of participants registered is 223. Among them, 183 participants will present their research. Most of the participants are from Indonesia, but also we have participants from Malaysia, Philippines, Pakistan, and India.

Besides the participants, the online nature of this conference also allows us to invite speakers from Europe and Australia. There will be Prof. Craig Moritz from Australian National University, Prof. Datin Maryati Mohammed from Universiti Tun Hussein Onn, Malaysia, Dr. Ahmed Abd El Wahed from University of Leipzig, Germany, Prof. Joana Falcao Salles from University of Groningen Netherlands, and Dr. Nastiti Wijayanti from the Faculty of Biology, Universitas Gadjah Mada, Indonesia. Because of the time difference, the speakers from Europe will give their talks on the afternoon of the first and second day of the conference.

The output of this conference will be published in the Advances in Biological Sciences Research as a conference proceeding, the Indonesian Journal of Biotechnology, and the Journal of Tropical Biodiversity and Biotechnology.

Finally, I would like to acknowledge the Indonesian Biology Consortium and the Publishers and Publications Board of Universitas Gadjah Mada which supported this conference. And I also like to thank you for our sponsors which make this conference possible.

I hope this conference will be fruitful for everyone. We look forward to seeing you all at the next ICBS conference.

Dr. Miftahul Ilmi, S.Si., M.Si
Chairman of 7th ICBS 2021

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher

of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

[Home](#) [Privacy Policy](#) [Terms of use](#)



Copyright © 2006-2022 Atlantis Press – now part of Springer Nature



STEERING COMMITTEE:

1. Prof. Dr. Budi Setiadi Daryono, M.Agr.Sc. (Dean of Faculty of Biology UGM and the Chairman of KOBI)
2. Dr. Rina Sri Kasiamdari, (Vice Dean of Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia)
3. Dr. Niken Satuti Nur Handayani, M.Sc (Vice Dean of Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia)
4. Dr. Eko Agus Suyono, M.App.Sc (Vice Dean of Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia)

ORGANIZING COMMITTEE

- Chairman : Dr. Miftahul Ilmi, S.Si., M.Si. (Faculty of Biology, Universitas Gadjah Mada)
- Secretaries : Novita Yustinadiar, S.Si., M.Si. (Faculty of Biology, Universitas Gadjah Mada)
Mutiar Fauziana, S.Si. (Faculty of Biology, Universitas Gadjah Mada)
- Treasurers : Titin Fauziah, S.E., M.B.A (Faculty of Biology, Universitas Gadjah Mada)
Pardiso (Faculty of Biology, Universitas Gadjah Mada)
- Pre-conference workshops PI : Dr. Fajar Sofyantoro, M.Sc. (Faculty of Biology, Universitas Gadjah Mada)
Annas Rabani, S.Si., M.Sc. (Faculty of Biology, Universitas Gadjah Mada)
- Proceedings and Publications : Dr. Wahyu Aristyaning Putri (Faculty of Biology, Universitas Gadjah Mada)
Dr. Dwi Sendi Priyono, S.Si., M.Si. (Faculty of Biology, Universitas Gadjah Mada)
- Promotion : Indra Lesmana, S.Si., M.Sc. (Faculty of Biology, Universitas Gadjah Mada)

[EN] We use cookies to help our viewer get the best experience on our website. -- [ID]
Kami menggunakan cookie untuk membantu pengunjung kami mendapatkan pengalaman terbaik di situs web kami

I AGREE / SAYA SETUJU

Mada)

Muhammad Nabil, S.Si. (Faculty of Biology, Universitas Gadjah

Mada)

- Equipment : Mulyanto, S.T., M.M. (Faculty of Biology, Universitas Gadjah Mada)

EDITORIAL BOARD

1. Yujiao Liang, M.D., Ph.D (ryougyokukou1987@yahoo.co.jp) Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Japan <https://www.scopus.com/authid/detail.uri?authorId=56857022600>
2. Wahyu Aristyaning Putri, M.Sc., Ph.D (wahyuaristyaningputri@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57196200480>
3. Dr. Dwi Sendi Priyono, M.Si. (dwisendipriyono@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57204157332>
4. Dr. Alona Cuevas Linatoc (alona@uthm.edu.my) Fakulti Sains Gunaan dan Teknologi, UTHM, Malaysia <https://scholar.google.com/citations?hl=en&user=p12ysNEAAAAJ>, <https://www.scopus.com/authid/detail.uri?authorId=54888614000>
5. Prof. Richard Luke Daniels, B.S., Ph.D (ldaniels@collegeofidaho.edu) Biology Department, College of Idaho, America <https://scholar.google.com/citations?hl=en&user=cfV8nL0AAAAJ>,
6. Prof. Parvez Alam (parvez.alam@ed.ac.uk) School of Engineering, Institute for Materials and Processes, The University of Edinburgh, UK <https://scholar.google.com/citations?hl=en&user=uafT9kgAAAAJ>
7. Shao Lun, Ph.D (shaolunliu@gmail.com) Department of Life Science & Center for Ecology and Environment, Tunghai University, Taiwan <https://scholar.google.com/citations?hl=en&user=tdEoWGMAAAAJ>
8. Dr. Miftahul Ilmi, M.Si. (m.ilmil@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57190191276>

REVIEWER

1. Yujiao Liang, M.D., Ph.D (ryougyokukou1987@yahoo.co.jp) Center for Infectious Diseases, Kobe University Graduate School of Medicine, Kobe, Japan <https://www.scopus.com/authid/detail.uri?authorId=56857022600>

[EN] We use cookies to help our visitor get the best experience on our website. -- [ID]

Kami menggunakan cookie untuk membantu pengunjung kami mendapatkan pengalaman terbaik di situs web kami.

PAKEE / SAYA SETUJU

2. Wahyu Aristyaning Putri, M.Sc., Ph.D (wahyuaristyaningputri@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57196200480>
3. Dr. Dwi Sendi Priyono, M.Si. (dwisendipriyono@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57204157332>
4. Dr. Alona Cuevas Linatoc (alona@uthm.edu.my) Fakulti Sains Gunaan dan Teknologi, UTHM, Malaysia <https://scholar.google.com/citations?hl=en&user=p12ysNEAAAAJ>, <https://www.scopus.com/authid/detail.uri?authorId=54888614000>
5. Prof. Richard Luke Daniels, B.S., Ph.D (ldaniels@collegeofidaho.edu) Biology Department, College of Idaho, America <https://scholar.google.com/citations?hl=en&user=cfV8nL0AAAAJ>,
6. Prof. Parvez Alam (parvez.alam@ed.ac.uk) School of Engineering, Institute for Materials and Processes, The University of Edinburgh, UK <https://scholar.google.com/citations?hl=en&user=uafT9kgAAAAJ>
7. Shao Lun, Ph.D (shaolunliu@gmail.com) Department of Life Science & Center for Ecology and Environment, Tunghai University, Taiwan <https://scholar.google.com/citations?hl=en&user=tdEoWGMAAAAAJ>
8. Dr. Miftahul Ilmi, M.Si. (m.ilmi@ugm.ac.id) Department of Tropical Biology, Faculty of Biology Universitas Gadjah Mada, Yogyakarta, Indonesia <https://www.scopus.com/authid/detail.uri?authorId=57190191276>
9. Furzani Pa'ee (furzani@uthm.edu.my) Department of Technology and Natural Resources, Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia, Johor, Malaysia. <https://scholar.google.com/citations?hl=en&user=yFR5k2AAAAAJ>
10. Prof. Uma Ramakrishnan (uramakri@ncbs.res.in) National Center for Biological Sciences, Tata Institute of Fundamental Research, India <https://scholar.google.com/citations?hl=en&user=zGGbdkEAAAAJ>
11. Prof. Laurentz Frantz (laurent.frantz@qmul.ac.uk) Ludwig Maximilian University Munich (LMU) & Queen Mary University of London, UK <https://scholar.google.com/citations?hl=en&user=DCK3qbcAAAAJ>
12. Mukhlis Jamal Musa Holle, M.Sc. (mukhlis.holle@zoo.ox.ac.uk) University of Oxford, UK <https://scholar.google.com/citations?hl=en&user=KRVCCcgAAAAJ>
13. Abdul Rahman Siregar, M.Biotech (asiregar@dpz.eu) German Primate Center | DPZ · Infection Biology Unit (Infektionsbiologie Deutsches Primatenzentrum GmbH Leibniz-Institut für Primatenforschung) and Faculty of Biology and Psychology Georg-August-Universität Göttingen, Germany <https://scholar.google.com/citations?hl=en&user=TSsPJDEAAAAJ>
14. Ratoe Soraya, M.D. (dr.ratoesuraya@stu.kobe.ac.jp) Kobe University, Japan

[EN] We use cookies to help our viewer get the best experience on our website. -- [ID]

Kami menggunakan cookie untuk membantu pengunjung kami mendapatkan pengalaman terbaik di situs web kami.

I AGREE / SAYA SETUJU

Copyright 2022 © **Flatsome Theme**

[EN] We use cookies to help our viewer get the best experience on our website. -- [ID]
Kami menggunakan cookie untuk membantu pengunjung kami mendapatkan
pengalaman terbaik di situs web kami.

I AGREE / SAYA SETUJU

Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

Search



Series: [Advances in Biological Sciences Research](#)

Proceedings of the 7th International Conference on Biological Science (ICBS 2021)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)

[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)

Search

[+ Advanced search](#)

SEARCH

91 articles

Proceedings Article

Tomato Leaf Curl New Delhi Virus Associated with Yellow Mosaic Disease of Cucumber (*Cucumis sativus*) in Bengkulu, Indonesia

Mimi Sutrawati, Sipriyadi Sipriyadi, Yuni Kristina Serlyani Sihotang, Cindy

Margareth Hutasoit, Parwito Parwito, Ewa Aulia

Cucumber mosaic virus, Papaya ringspot virus, Squash mosaic virus, Cucurbit aphid-borne yellows virus, Cucurbit aphid-borne yellows virus Tobacco mosaic virus, Tomato yellow leaf curl New Delhi virus and Zucchini yellow mosaic virus were among the viruses that infected cucumbers in Java, according to...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Application of a New Drying Method on the Quality of Voucher Specimen

Mohd Amril Ramzi Bin Mohd Rahimi, Furzani Pa'ee

Drying the specimens is one of the crucial steps in plant preservation and the process of making Herbarium voucher specimens. This research was conducted to investigate the effect of the drying method on specimens in the laboratory using the standard method (oven) and drying in the field using a novel...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

From Invasive to Creative: Transforming *Salvinia molesta* in Taman Botani Sri Medan into Cosmeceutical Solid Soap

Nurul Syamimi Muzaini, Furzani Pa'ee

Salvinia molesta is invasive species and is also known as the worst weed that invaded Malaysia's native wide range of aquatic ecosystem. This study was conducted to identify the potential metabolites *Salvinia molesta* invasive plant species from family Salviniaceae in Taman Botani Sri Medan, Batu Pahat,...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Antioxidant Activity Evaluation of Agarwood *Aquilaria malaccensis* Lamk. Leaves Extract Using DPPH, FRAP and ABTS Assays

Aprilliani Prissilla Halim, Nastiti Wijayanti, Lisna Hidayati, Tri Rini Nuringtyas

Indonesia has been known to have large forest areas and rich in biodiversity, which could be the source of primary and secondary metabolites. Along with the technological development and research in medicine, agarwood, which was initially used only for topical body treatments, room fragrances, and religious...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Hematology Profile of Female Guinea Pig (*Cavia porcellus* (Linnaeus, 1758)) with Diet Variations

Naila Nabila Rahmani, Ayu Aziza Ar Rachid, Laksmindra Fitria

Guinea pigs (*Cavia porcellus* (Linnaeus, 1758)) is a rodent other than rats and mice that are commonly used in biomedical research because of its physiological similarity to humans. To meet the requirement as a standard laboratory animal, the care management of GP needs to be controlled, one of which...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Genetic Diversity of *Moringa Oleifera* on Poteran Island-Madura Based on Petiole Colors Using ISSR (*Inter Simple Sequence Repeat*) Method

Wirdhatul Muslihatin, Triono Bagus Saputro, Nur Isma Latifah, Chusnul Eka Safitri Himayani

Moringa oleifera, which is endemic of Pulau Poteran, Madura, is one of a genetic variation of *Moringa*. It is native Indonesian germplasm, which must be preserved to maintain the genetic biodiversity of *Moringa*. Known locally as kelor, the plant has four different color of petiole: red, white, purple,...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Hematology Profile of Guinea Pigs [*Cavia porcellus* (Linnaeus, 1758)] Based on Sex and Age

Laksmindra Fitria, Sri Lestari, Asti Nur Istiqomah, Novita Paradhita Wulandari, Anggadiah Shinta Wardani

Guinea pigs or *Cavia porcellus* (Linnaeus, 1758) is small herbivorous rodent from South America. Guinea pigs (GP) are commonly used as experimental animals, pets, food, and animal-assisted therapy (AAT). Based on these benefits, the health of GP must be considered. Blood is essential biological sample...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Molecular Comparison between Two Similar Asteraceae

Species (*Synedrella nodiflora* (L.) Gaertn. and *Eleutheranthera ruderalis* (Swartz) Sch._Bip) by the Use of *trnL*(UAA) - *trnF*(GAA) Intergenic Spacer

Murni Dwiati, Indrawati Indrawati, Agus Hery Susanto

Synedrella nodiflora (L.) Gaertn and *Eleutheranthera ruderalis* (Swartz) Sch.-Bip. are two Asteraceae species showing very similar morphological appearances. Both are broad-leaf weed species in some crops throughout many tropical areas. Nevertheless, the individual species can be potentially utilized...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Potential of Production and Characteristic of Oleoresin Tapped from *Dipterocarpus verrucosus* as Natural Ingredient for Multi Purposes

Andrian Fernandes, Rizki Maharani

Keruing (*Dipterocarpus*) is one of the Dipterocarpaceae family which produces oleoresin. *Dipterocarpus* species diversities will give the difference in oleoresin obtained from tapping, and its physical and chemical properties. This research aimed to determined the distribution of tapped oleoresin of D...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Metabolite Profiling of *Davallia* in The Mentawai Islands, West Sumatra, Indonesia

Mildawati Mildawati, Sobir Sobir, Sulistijorini Sulistijorini, Tatik

Chikmawati

This study revealed the metabolite compounds of the *Davallia* species in the Mentawai Islands, West Sumatra, through metabolite profiling. This study aimed to determine the chemical compounds in the leaves of the *Davallia* species. Leaf samples of three species, *D. denticulata* (Burm. f.) Kuhn var. *denticulata*,...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

In Vitro Anthelmintic Activity of *Limonia acidissima*, L. Leaves Aqueous Extract on *Haemonchus contortus* (Rudolphi, 1803)

Muh. Andhi Hardianto, Slamet Widiyanto

Mortality, loss of production, slowed growth, poor weight gain, and even death are all common economic losses caused by haemonchosis. Due to the emergence of anthelmintic resistance and the expensive expense of chemical anthelmintic treatments, medicinal plants have been investigated as potential anthelmintics....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Developing RAPD-derived SCAR (Sequence Characterized Amplified Region) Marker for Flowering Time in Chili Pepper

Estri Laras Arumingtyas, Bunga Rizky Elfa Agustina, Joni Kusnadi

The selection step of plant breeding is a crucial stage in the process of developing new varieties which usually take a considerable long time in a

conventional breeding. The development of molecular markers for selection has been carried out to reduce the time required. A Sequence Characterized Amplified...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

UV Protectant Ability of *Attacus atlas* L. (Lepidoptera: Saturniidae) Sericin Extract to Increase Nucleopolyhedrovirus Effectiveness against Beet Army Worm, *Spodoptera exigua* (Hübner) (Lepidoptera: Noctuidae)

Hana Widiawati, Sukirno Sukirno, Siti Sumarmi, Hari Purwanto, R.C. Hidayat Soesilohadi, Ignatius Sudaryadi

Spodoptera exigua (Lepidoptera: Noctuidae) is the common pest known for attacking shallot crop. Baculovirus (Nucleopolyhedrovirus: NPV) is a biological agent that is widely used as the pest control agent. However, the activity of NPV is deteriorated when applied in the field due to the influence of ultraviolet...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Effect of UV Radiation and Fruit Feedings (Banana and Guava) on the Survival Rate and Morphological Changes of Reproductive Organ of Fruit Fly (*Drosophila melanogaster* Meigen, 1830)

Hipny Alwandri, Nafisa Kusumawati, Ignatius Sudaryadi

The increasing ozone concentration in the upper stratosphere was not significant enough to be able to protect from the detrimental effect that

ultraviolet (UV) light radiates. One solution to counter this problem is by using antioxidants as protectants, such as those contained in fruits. In order to...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Diversity of Macroalgae in the Intertidal Zone of Gili Ketapang Beach, Probolinggo, Jawa Timur, Indonesia

Natasya Meri Auliadani, Faradilla Faradilla, Abdul Razaq Chasani

Indonesia is a country that has high potential in biodiversity, especially in the marine areas. As the primary producer of marine ecosystems, macroalgae provide food for many species. Macroalgae have very varied structures but are generally divided into three groups based on their color of thallus i.e....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Diversity of Insect Based on Growth Stages of Rice (*Oryza sativa* L. 'IR 64') at High Altitude in Kepurun Village, Manisrenggo Sub-district, Klaten District, Central Java

Aryo Seto Pandu Wiranto, Nindita Sabila Ningtyas, Regina Diah Rachmawati, Rahmatullah Rahmatullah, Sukirno Sukirno

Rice (*Oryza sativa* L.) is one of the biggest commodities in Indonesia. Rice is cultivated in monoculture, so it affects the diversity of insects present in rice fields. The purpose of this research is to study the diversity of insect present at different 'IR 64' rice growth stages including seedling...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Application of Rhizobacteria and NPK for Growth and Productivity of Sweet Corn (*Zea mays* L.)

M. Shovitri, S.K. Sugianto, N.D. Kuswytasari, N.H Alami, E. Zulaika

Soil fertility is one of the limiting factors for sweet corn crops. To overcome it, the wise use of fertilizer is a must to maintain long soil sustainability. Our previous study showed that a rhizobacteria consortium had a significant positive effect on crop growth and productivity; even the chemical...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Antibacterial Activity and Toxicity Study of Selected *Piper* Leave Extracts Against the Fish Pathogen (*Aeromonas hydrophila*)

Norashikin Anjur, Siti Fatimah Sabran, Hassan Md Daud, Nor Zalina Othman

The extensive use of antibiotics in aquaculture has resulted in the emergence of bacterial resistance strains. The medicinal importance of the herb such as *Piper betle*, *Piper sarmentosum*, and *Piper nigrum* evidently proved as one of the most promising commercial botanicals with earlier reported to possess...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Numerical Taxonomy of Marine Macroalgae Gracilariaceae from Southern Coast of Gunungkidul Based on Morpho-Anatomical and Phytochemical Characters

Adinda Nur Anisa, Abdul Razaq Chasani

Gracilariaceae is a macroalgal family of Rhodophyta which can be found abundantly in tropical waters, including in Southern Coasts of Gunungkidul D.I.Yogyakarta, Indonesia. Coral and sand dominate the substrate of Southern Coasts of Gunungkidul, so it is an ideal habitat for Gracilariaceae. Along with...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

A Year of COVID-19 Outbreak in Indonesia #2: Variant Development Based on *Spike (S)* Mutations

Nicholas Gerry Andreanto, Dwita Novitasari, Delia Wahyu Pangesti, Rizqi Layli Khusufi, Annasa Sabatia, Irwansyah Nur Oktafian, Nur Alfi Maghfirotus Sa'adah, Reni Krisdayana, Salsabila Kasta Hygiea Iswara, Dwi Listyorini

SARS-CoV-2 has infected millions of people in Indonesia and taken thousands of lives by bonding Spike (S) protein and Angiotensin-Converting Enzyme 2 (ACE2) human cell receptor. Spike gene has a higher mutation rate compared to other genes, which suggested to increase its virulence, transmission and...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Effect of UV Radiation and Treatment to Orange (*Citrus sinensis* L. Osbeck) Fruit Feeding on the Survival Rate and

Colony Sex-ratio of Fruit Fly (*Drosophila melanogaster* Meigen, 1830)

Ignatius Sudaryadi, Yulia Maulita Janah, Nafisa Kusumawati

Ultraviolet (UV) is a form of physical stress that forces living organisms to respond to the challenge of DNA alteration. UV light causes oxidative stress by causing the creation of reactive oxygen species (ROS). Antioxidants are substances that can interfere with the production of reactive oxygen species...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Diversity and Origin of Mammal Collection in Mini Zoo in Sleman, Yogyakarta Special Region

Dea Evani Amelia, Bambang Agus Suropto

The mini zoo that became a local tourist spot manifests the ex-situ conservation of mammal species. Animal welfare principles play an important role in long-term sustainability during the animal care and management period. Mini zoos in Sleman, Yogyakarta Special Region were assumed that their diversity...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Carbon Sequestration of Tree Community in Urban Green Spaces of Bekasi City, Indonesia

Kristian Briantama, Adi Basukriadi

Bekasi City is experiencing climate problems, one of which is a decrease in rainy days. The existence of the Patriot Bina Bangsa City Forest and Bekasi

City Park is expected to reduce the impact of climate change through the role of the tree communities in it as a carbon sink. The problem faced is the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Histological Observation, Identification, and Secondary Metabolites Content in Endophytic Fungi of Mahogany Tree (*Swietenia mahagoni* Jacq)

Utami Sri Hastuti, Sulisetijono Sulisetijono, Chomisatut Thoyibah, Siti Hartina Pratiwi, Khusnul Khotimah

The mahogany (*Swietenia mahagoni* Jacq.) tree is a sort of medicinal plant. The phytochemical analysis by previous research has proved that the mahogany bark methanol extract is contained i.e alkaloid, tannin, saponin, phenol hydroquinone, and flavonoid. Some medicinal plant that has mutualism symbiotic...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Species Diversity of Insects on Tobacco (*Nicotiana tabacum* L. 'Bligon' and 'Grompol') Plantation in Sokorini Village, Muntilan, Magelang, Central Java

Aryo Seto Pandu Wiranto, Siti Sumarmi

Tobacco is an agricultural export commodity important for Indonesia. Therefore, tobacco farming, especially in optimizing production, is an attractive point for research. The challenges in optimizing tobacco production are insect pests causing loss on tobacco production. This study aimed to identify...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Effect of Altitude on Odonata Biodiversity in the Paddy Field of Sleman Regency, Special Region of Yogyakarta

Nariswari Salsabiela, Amanda Novitasari, Agustina Citra Windianingsih, Reza Bagus Alfian, Anggun Setyaningrum, Besta Eins Yudharta, Okti Alfiyatus Safa'ah, Sukirno Sukirno

Odonata acts as a natural enemy in the paddy field ecosystem. The Odonata diversity is highly related to habitat condition. This research analyzed the effect of altitude on the diversity of Odonata in paddy field ecosystems in Sleman Regency, Special Region of Yogyakarta. This research was conducted...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Activity of Lactate Dehydrogenase in Sapera, Saanen, and Ettawa Crossbred Goats in the Different Physiological Statuses

Sarmin Sarmin, Irkham Widiyono, Devita Anggraeni

The enzyme that catalyzes the conversion of lactate to pyruvate is called lactate dehydrogenase (LDH), essential in producing energy present in cells for various tissues, including the liver, heart, and skeletal muscle. Although the increased LDH activity is due to vascular thrombosis, bleeding, and...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Computational Study of Natural Compounds in Melon Fruit (*Cucumis melo* L. 'GMP') as Inhibitor of Epidermal Growth Factor Receptor Protein

Wiko Arif Wibowo, Teuku Nanda Saifullah Sulaiman, Supriyadi Supriyadi, Budi Setiadi Daryono

'GMP' melon is a breeding cultivar that has a bitter taste and fragrant aroma. The bitter taste character indicates the presence of potential natural compounds that can be used as anti-cancer. This study aims to reveal the natural compounds of 'GMP' melon and its use as an anti-cancer computationally....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Occurrence of Pest, the Management of Zoological Museum Specimens Collection and Climate Change

Arney Sapaat, Siti Fatimah Sabran, Maryati Mohamed

This study aims to assess the occurrence of pests of zoological specimen collections at natural history museums, management of the specimens with climate change patterns in Malaysia. The objective of this study was to determine the causes that damaged zoological specimens and the effectiveness of the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Morphological Characters and Plant Pigments Content of

Three Varieties of Chrysanthemum Induced by Paclobutrazol Treatments

Intani Quarta Lailaty, Laurentius Hartanto Nugroho

Chrysanthemum is one of the favorite ornamental plants as a potted flower. Consumers currently prefer potted flowers with short stems, lush leaves, also uniform and compact flowers. It is necessary to form the potted flowers by applying paclobutrazol (PBZ). This research aimed to determine the effectiveness...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Potency of Red Betel (*Piper crocatum* Ruiz & Pav.) Methanolic Extract as α -Amylase and α -Glucosidase Inhibitor

Yustina Sri Hartini, Dewi Setyaningsih

Methanol is an effective menstruum for attracting compounds with various pharmacological activities from *Piper crocatum* Ruiz & Pav. Several studies reported that laboratory tests for reducing blood sugar content of red betel extract such as aqueous, ethanolic, and ethyl acetate extracts. This study...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Application of Deep (Machine) Learning for Phytoplankton Identification Using Microscopy Images

Arief Rachman, Aulia Salsabella Suwarno, Susanna Nurdjaman

As a hot spot of marine diversity, between 150 – 400 phytoplankton species have been reported in various Indonesian marine ecosystems. However,

phytoplankton identification in Indonesia is mainly made manually by a human expert, which is a time-consuming process with many limitations. Thus, this study...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Molecular Docking of Anthocyanin Compound as Anti-Hyperlipidemia Against PPAR α , HMG-CoA Reductase and ACAT Proteins

Noor Nailis Sa'adah, Elshinta Riantica, Awik Puji Dyah Nurhayati, Nova Maulidina Ashuri, Dewi Hidayati

Hyperlipidemia can increase the risk of cardiovascular diseases, such as coronary heart disease (CHD), the number one cause of death worldwide each year. Therefore, needed efforts to reduce the prevalence of cardiovascular disease, one of which is parijoto fruit (*Medinilla speciosa*), which has a high...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Application of Chitosan-*Aloe vera* Gel Based Coating on Postharvest Quality and Storability of Red Chili (*Capsicum annum* L.)

Yora Faramitha, Fitria Febriyanti, Tiana Fitrilia, Firda Dimawarnita, Siswanto Siswanto

Red chili pepper (*Capsicum annum* L.) is a high-demand horticultural commodity in Indonesia but is vulnerable to quality deterioration. Postharvest treatment is needed to maintain the quality and extend the shelf life of red chilies. One of the promising preservation technologies to

prolong the shelf-life...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Effect of UV-B Radiation Intensity on *Hsp70*, *mtHsc70-1*, and *cpHsc70-2* Gene Expression in *Synedrella nodiflora* (L.) Gaertn. Leaf Under Field Conditions

Maria Celinna, Andi Eko Maryanto, Andi Salamah

Increased expression of genes encoding Heat Shock Protein 70 (Hsp70) is one of the plant defense responses against UV-B stress. *Synedrella nodiflora* may be tolerant to relatively high UV-B intensity. The research has been conducted to compare expression levels of genes encoding cytosolic, mitochondrial,...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Isolation and Characterization of *Vanda Orchid Homeobox* Gene from *Vanda tricolor* var. *Suavis* Lindl. form Merapi

Viantius Ruben, Muhammad Dylan Lawrie, Endang Semiarti

Vanda tricolor var. *Suavis* Lindl. form Merapi is one of the crucial Indonesian orchid species. Due to natural disasters or deforestation of their natural habitat, the population of this orchid continues to decline and is threatened to be extinct. Therefore, a strategy for mass propagation of this plant...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Genetic Variations and Phenetic Relationships of Hairy Bittercress (*Cardamine hirsuta* L.) Accession in Java based on *Inter-Simple Sequence Repeat*

Tania Agnesa, Purnomo Purnomo, Budi Setiadi Daryono

Cardamine hirsuta from the Family Brassicaceae is a potential plant that can be used as an object of comparative study with *Arabidopsis thaliana* (L.) Heynh. Plant comparative studies involving closely related species with similar traits are used to investigate the genetic pathways underlying morphological...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Molecular Docking Simulation of Trisindolina 1 Compound Against Pi3k Protein in Hepatocellular Carcinoma

Evira Nadila Oktyasti, Awik Puji Dyah Nurhayati

The increased cancer burden globally, from 12.7 million new cases in 2008 to a predicted 22.2 million in 2030, makes cancer a critical global problem with high unmet medical needs. Hepatocellular carcinoma (HCC) is the fifth most frequently diagnosed cancer worldwide, and HCC is the third leading cause...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Determination of Lectin Genes in Superior Mutant of Rodent Tuber Bogor Accession (*Typhonium flagelliforme*) Based on

PCR Amplification

Nesti Fronika Sianipar, Reflinur Reflinur, Muhammad Dzulkifly Ashan, Khoirunnisa Assidqi, Dwityantari Widyaningrum, Ragapadmi Purnamaningsih

Rodent tuber (*Typhonium flagelliforme*) is one of the Indonesian herbs that has not yet been developed as an anticancer drug. The local name rodent tuber plant has produced superior mutant strains with higher levels and bioactive compounds. Compared to the mother plant, GCMS analysis detected four new...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Effectiveness of N-Hexane and Ethanol Extract of Giant Calotrope (*Calotropis gigantea* L.) Leaves as Insecticide Against Shallot Pest *Spodoptera exigua* (Hübner)

Elvian Indah Nilamsari, L. Hartanto Nugroho, Sukirno Sukirno

Shallot (*Allium ascalonicum*) is one of the high-value plants. However, shallot productivity has been decreased due to some disturbing organisms. *Spodoptera exigua* is one of the pests on shallot farming and causes a yield reduction of up to 70%. Farmers use excessive synthetic pesticides to overcome the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Mutation Analysis of PRD-domain of ROR2 and Ig-domain of FLNA in Breast Cancer Development: A Case Study in Malang

Rizqi Layli Khusufi, Delia Wahyu Pangesti, Dwita Novitasari, Vina Rizkiana, Reni Krisdayana, Salsabila Kasta Hygiea Iswara, Dwi Listyorini

ROR2 is a WNT receptor involved in non-canonical Wnt signaling pathways. PRD and IgFLNA domains are functional domains for binding to downstream proteins that lead to cancer development. This study aimed to analyze mutations in the PRD-IgFLNA-ROR2 domain also the role of the WNT5A-ROR2 pathway involved...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

LC50 and Effect of Sublethal Concentration of K₂Cr₂O₇ on Different Developmental Stages of *Osteochilus vittatus*

Gratiana E. Wijayanti, Sharon Hillary, Ani Septiani, Anisa Aulia, Anastasia E. Sintanora

Chromium in form of hexavalent had been reported to be genotoxic and carcinogenic, affects some physiological, and reproductive features. The ability of aquatic animals including fish to tolerate chromium may vary according to different developmental stages. Therefore, this research was conducted to...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Phenotypic Characters Stability of Melon (*Cucumis melo* L. 'Kinaya')

Amir Muhammadi, Budi Setiadi Daryono

Melon is an important commodity in Indonesia and still dominated by imported melons such as Action 434 and Sun Lady. The 'Kinaya' cultivar is a crossbreed between 'Kinanti' and 'Sonya' cultivars. The development of

'Kinaya' cultivar as indonesian superior local melon is needed to improve the quality...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Computational Model of Trisindoline 1 Conjugate to Protein P53 and P53R2: Targets For Breast Cancer Therapy

Shabrina Syifa Ghaissani, Awik Puji Dyah Nurhayati, Vencka Azzahra Putri

In 2020, there were 2.3 million women diagnosed with breast cancer with average, 7 percent to 11 percent of women with early breast cancer experience a local recurrence during this time. Resistance mechanisms in breast cancer include DNA repair mechanisms that protect cancer cells from endogenous or...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Sexual Dimorphism of *Leptocorisa oratorius* Fabricius (Hemiptera: Alydidae) from Special Region of Yogyakarta

Fanuel Triaswanto, Erik Lawijaya, Ananto Puradi Nainggolan, IGM Raka Alpin Aditya, RCH Soesilohadi

Sexual dimorphism (SD) is a common phenomenon in animals, especially in invertebrates and poikilothermic vertebrates, which can cause body size differences of males and females in a species. Rice ear bugs [*Leptocorisa oratorius* Fabricius (Hemiptera: Alydidae)] are a significant rice pest in Indonesia,...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Potency of Inulin to Increase Curcumin Aqueous Solubility as a Co-Processing Material with *Curcuma longa* in Solid Dispersion Approach

Dewi Setyaningsih, Elizabeth Vianita Kurniawan, Yustina Sri Hartini

Curcumin, an identical compound of *Curcuma longa* extract, shows limited bioavailability due to its lipophilicity. Inulin of degree of polymerization (DP) of 11 is a carbohydrate member that can potentially increase aqueous solubility and dissolve lipophilic compounds. This study aimed to investigate...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

CO2 Mitochondrial Gene Identification of *Nisaetus cirrhatus*

as a Part of Indonesian Elang Brontok Genetic Conservation

Reni Krisdayana, Rizqi Layli Khusufi, Dwita Novitasari, Annasa Sabatia, Delia Wahyu Pangesti, Salsabila Kasta Hygiea Iswara, Dwi Listyorini

The Elang Brontok, scientifically named as *Nisaetus cirrhatus* is included in the least Concern in appendix II. The use of the COI gene for taxonomic in our previous study put our samples in a complex species taxonomical position. Further research needs to be done using other mitochondrial genes such...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Pollen Diversity and Propolis's Bioactive Compounds of Stingless Bees (*Tetragonula laeviceps*, Smith 1857) From Kedungpoh Meliponiculture, Gunungkidul, Yogyakarta.

Fiola Oktaweni, Sutikno Sutikno, Ignatius Sudaryadi

The progression of many diseases due to viruses and bacteria makes for an increase in natural alternative medicine. One source of natural medicine is honey and propolis of stingless bees. Efficacy physical and chemical properties of honey are influenced by the type of pollen and the environment, while...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Effects of Water Availability on Physiological Factors of Cayenne Pepper Plant *Capsicum frutescens* L.

Fiana Lathifah, Dwi Umi Siswanti

Chili pepper is a vegetable plant that is consumed by many people and has

high economic value. Chili peppers contain secondary metabolite compounds, capsaicin which is synthesized in the epidermis cells. Capsaicin acts as a spicy taste in chili peppers, so chili peppers are favored by the public. Environmental...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Miocene Giraffids (Giraffidae; Mammalia) from the Lower Siwalik of Pakistan

Kiran Aftab, Muhammad Akbar Khan, Sumera Afsheen, Mubashar Hussain, Areej Arif

New dental material of Giraffidae is recorded from the Lower Siwalik, Punjab, Pakistan. The specimens are assigned to two genera, Giraffokeryx and Giraffa. The new material comprises isolated teeth, maxilla, and mandible fragments. The material was recovered from Lower Siwalik localities, Chabbar Sayadan...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Application of Growth Regulatory Substances CPPU and GA3 on the Growth of Porang Plants from Bulbil

Tutik Nurhidayati, Kristanti Indah Purwani, Zulfan Febriawan, Firda Fortuna Nasich

Porang is known as an alternative food source. The main obstacle for porang production is the long harvest period with growth time needed ranging between 4-5 months and 7-8 months of dormancy. The application of plant growth regulators (PGR) such as CPPU and GA3 could be used to break porang's dormancy....

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Importance of Purification and Activity Analysis of the Purified Product of Thrombolytic Protease from *Bacillus* sp. HSFI-12– A Review

Nurul Islamiyah, Stalis Norma Ethica, Muhammad Ardi Afriansyah, Ana Hidayati Mukaromah, Dewi Seswita Zilda

Mortality and morbidity of Cardio-vascular diseases (CVDs) have been the major issue in the group of non-communicable diseases worldwide. However, the existing antithrombotic drugs to combat CVD still have many shortcomings in terms of price and safety. *Bacillus* sp. HSFI-12 had been previously reported...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Introduction to Plant Metabolism, Secondary Metabolites Biosynthetic Pathway, and In-Silico Molecular Docking for Determination of Plant Medicinal Compounds: An Overview

Risanti Dhaniaputri, Hadi Suwono, Mohamad Amin, Betty Lukiati

Natural ingredient produced by plants are widely used for therapeutic treatment, because they are believed to have fewer side effects and are cheaper than synthetic drugs. Plants used as treatment media contain natural secondary metabolites compounds derived from primary and secondary metabolism. Primary...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Effectiveness of Bio-catharanthine on Peanut (*Arachis hypogea* L.) Lurik Cultivar

Dwi Indahning Rohmah, Melza Mulyani, Laras Nur Janah, Adi Pancoro, Miftahudin Miftahudin, Anjar Tri Wibowo, Budi Setiadi Daryono

The productivity of peanut (*Arachis hypogea* L.) in Indonesia from 2017 until 2021 is estimated to have average production growth minus 11,82% per year and cannot meet the domestic need of the peanut. According to that statistic, Indonesia is the world's second largest importer of peanuts. Looking for...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Detection of Terpenoid and Flavonoid of Five Species of Mistletoes at *Stelechocarpus burahol* (Bl.) Hook.f. & Th and *Lagerstroemia speciosa* (L.) Pers. by using Thin Layer Chromatography Method

Djoko Santosa, SM. Widyastuti, Umami Rosyidah, Betha Silmia

Mistletoe is a parasitic plant that is well known for causing significant damage to forestry plants; nevertheless, its value in traditional medicine has yet to be proven. This study aims to investigate terpenoid and flavonoid compounds of *Macrosolen cochinchinensis* (Lour.) Tiegh, *Scurrula atropurpurea*...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Prospective *In Vivo* Assays on the Antithrombotic Potential of Protease Extracted from *Bacillus* sp. HSFI-12

Okta Yosiana Dewi, Stalis Norma Ethica, Andri Sukeksi, Maya Dian Rakhmawatie, Sri Darmawati

CVD (cardiovascular disease) is a group of non-communicable diseases and a global cause of cardiovascular death. The search for agents inhibiting blood clot formation (thrombus) and enhancing antithrombotic activity are important for the prevention and treatment of CVD. Previous study reported that bacteria...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Prospective Purification and Assay of Thrombolytic Protease from *Bacillus* sp. HSFI-10 Isolated from Sand Sea Cucumber for Antithrombotic Agent Development

Bio Putri Ayanti, Stalis Norma Ethica, Ayu Rahmawati Sulisytaningtyas, Sri Sinto Dewi, Dewi Seswita Zilda

Thrombosis is a cardiovascular disorder due to the formation of a blood clot (thrombus), which can cause a blockage in a blood vessel. Such abnormality is responsible for more than millions of deaths per year in the world. In addition to cardiovascular disease, thrombosis can also occur in patients with...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

A Literature Review on the Potential of the Biodiversity of Thrombolytic Protease-Producing Bacteria Isolated from

Brown Seaweeds *Chnoospora* sp.

Nurhilaliyah Nurhilaliyah, Stalis Norma Ethica, Wikanastri

Hersoelistyorini, Aditya Rahman Ernanto, Wijanarka Wijanarka

Bacterial proteases with antithrombotic activities can be used to treat cardiovascular disease (CVD), one of the leading causes of death in the world. Indonesia is known for its high marine biodiversity, including its brown seaweed varieties and their symbiotic bacteria. The high biodiversity of proteolytic...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Potential of *Dendrobium* spp. Secondary Metabolites as Medicinal Source for SARS-CoV-2

Muhamad Rafli, Tri Rohmiati, Anggiresti Kinasih, Alim El Hakim, Endang Semiarti

Coronaviruses have long been a severe issue due to their capacity to mutate and infect the respiratory tract. *Dendrobium* orchids are the most diverse orchid species in Indonesia. Polysaccharides, bibenzyl, phenanthrene, coumarin, sesquiterpenoids, alkaloids, and steroids were among the phytochemical...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Behavior and Food of Reintroduced Bornean Orangutan (*Pongo pygmaeus wurmbii*) at Feeding Site and Forest Area in Lamandau Wildlife Sanctuary

Yulia Raudhatul Balaqis Zahro, Ani Mardiasuti, Dede Aulia Rahman

The Bornean orangutan (*Pongo pygmaeus wurmbii*) is a Critically Endangered large Asian primate. Reintroduction had been implemented to recover their population in the wild. The research was conducted in February-April 2021 at Lamandau Wildlife Sanctuary, aimed to identify behavior differences at the feeding...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Lipolytic and Proteolytic Activities of Fibrolytic Bacteria from Buffalo (*Bubalus bubalis*) Rumen

Siti Lusi Arum Sari, Triyanto Triyanto, Zuprizal Zuprizal, Irfan Dwidya Prijambada

The use of plant-based feed ingredients in aquaculture is expected to reduce feed costs. However, plant ingredients generally have a low protein content and are difficult to be digested by monogastric animals. The fermentation process is proven to increase the digestibility and nutritional value of plant...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Comparison of the 12S rRNA and COI Regions of Mitochondrial DNA for eDNA Detection of Alligator Gar (*Atractosteus spatula*)

Patricia Agustin, Andi Eko Maryanto, Noviar Andayani

Alligator gar (*Atractosteus spatula*) has been introduced to many places outside its native range. The fish is considered as an invasive alien species that needs to be eradicated upon encounter in Indonesia due to its aggressive behavior. Despite existing regulations in Indonesia, the fish is

still bred...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Comparison of the Diversity of Nepenthaceae, Orchidaceae and Zingiberaceae in Disturbed and Undisturbed Forests in Johor, Malaysia

Vinod Kumar Sivarajah, Alona Cuevas Linatoc

This study was carried out to compare the diversity of Nepenthaceae, Orchidaceae, and Zingiberaceae families in central Johor. The chosen locations are the Belumut Forest-Eco Park, Mo'akil Permanent Forest Reserve, and Soga-Perdana Permanent Forest Reserve, which have forest types categorized as, undisturbed...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Effects Leaf Ethanol Extract of *Graptophyllum pictum* L. Griff. to Inhibit Vaginal Atrophy of Menopausal Mouse

Listijani Suhargo, Sri Puji Astuti W., Alfiah Hayati

Menopausal condition with low estrogen level caused vaginal atrophy that it is a condition where vaginal lining become dryer and thinner and it would be easier to get inflammation. So, it was important to find out the treatment to inhibit it. This research was aimed to determine the effects of *Graptophyllum*...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Construction, Cloning, and Overexpression of Staphylococcal Enterotoxin B Gene Synthetic (SEBSyn) in pET-28a(+): Pre-development Bacterial-Toxin Therapy for Cancer

Achmad Rodiansyah, Marselina Irasonia Tan, Husna Nugrahapraja

Staphylococcal enterotoxin B (SEB) from *Staphylococcus aureus* could be considered as a therapeutic agent to eliminate cancer cells. SEB can activate immune response, which furthermore could induce apoptosis of various cancer cells. This study was proposed to design a SEBSyn coding sequence suitable for...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Antibacterial Effectiveness of Synthesized Copper Nanoparticles by Ultrasonication Assisted Method

Ainul Fitria Mahmudah, Yuni Kusumastuti, Himawan Tri Bayu Murti Petrus, Yekti Asih Purwestri

Antibiotic resistance among pathogenic bacteria has become a problem in the medical community. Copper nanoparticles (CuNPs) have become one method of combating antibiotic resistance in bacteria. The antibacterial activity of Synthesized Copper nanoparticles (CuNPs) from sigma 774103-5G against various...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Weight Growth of the Hybrid Chicken (*Gallus gallus domesticus*, Linnaeus 1758) Crossing Result of Female Pelung with Male F₃ Golden Kamper

Salma Dewi Pratita, Nareta Defiani, Afifah Nur Hidayah, Nurul Hidayah, Putri Shafira Setyowati, Budi Setiadi Daryono

Backcross is a cross between an offspring and its parents, or an individual that is genetically similar to its parents, to minimize the genetic variation of the resulting offspring. A female Pelung cross with a male F₃ Golden Kamper is a type of backcross. Golden Kamper chicken is a breed of chicken...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Immunostimulant Effects of Alang-Alang (*Imperata cylindrica*) Roots Extract on BALB/c Male Mice (*Mus musculus*)

Rosnizar Rosnizar, Fajar Muliani, Iskandar Muda Ramli, Kartini Eriani

Alang-alang (*Imperata cylindrica* L.) is an annual rhizomatous grass and traditionally well known for its therapeutic values, especially in treating fever, muscle soreness, nosebleed and respiratory asphyxia. It has been reported to contain different classes of secondary metabolites and attracted scientists'...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Study of Digestive Tract Diseases in Cats

Soedarmanto Indarjulianto, Yanuartono Yanuartono, Alfarisa Nururrozi,

Slamet Raharjo, Hary Purnamaningsih, Irkham Widiyono, Sri Hartati, Juni Claudia Dami, Luh Putu Eka Damayanti

The cat's digestive tract is one of the vital systems that often facing the deadly diseases. This study aims to identify digestive system disorders/diseases in cats. A total of 100 cats of various breeds and ages with one or more symptoms of digestive disorders was used in this study. All cats were physically...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Pyridinoline and Deoxypyridinoline in Oral Fluids of Menopause Women as Predictor Alveolar Bone Resorption

Agustin Wulan Suci Dharmayanti, Hendy Hendarto

Menopause presents menstrual cessation that occurs physiologically and is associated with osteoporosis and periodontitis. Osteoporosis and periodontitis are silent diseases that present clinically in the late stage, which alveolar bone resorption causing tooth loss represents both of these disorders....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Growth and Productivity of Lurik Peanuts (*Arachis hypogaea* L. var. Lurikensis) after Biofertilizer-Sludge Biogas Application

Dwi Umi Siswanti, Nur Hidayah Pangestuti, Niken Wulansari

One of the most important crop commodities in Indonesia is peanut. Ipeanutse peanut production can be done in various ways, including plant

breeding, and improving land quality through organic fertilization. The effect of the use of biofertilizer combined with sludge on crop productivity of peanuts,...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Potential High Conservation Value of Mount Ungaran as a Step-stone for Essential Ecosystem Area Plan

Margareta Rahayuningsih, Nana Kariada Tri Martuti, Dyah Kartikasari, Lutfian Nazar

High conservation value (HCV) helps stakeholders conduct sustainable forest management by considering social and environmental aspects. It requires a high level of protection to ensure the high conservation value of an area persists in the long term. Mount Ungaran is one of the essential areas in Central...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

A Meta-Analysis Study on *Spodoptera exigua* and *Spodoptera litura* Control: Biopesticides vs. Synthetic Pesticides

Nadya Sofia Siti Sa'adah, Hipny Alwandri, Laurentius Hartanto Nugroho, Sukirno Sukirno, Tri Rini Nuringtyas

Polyphagous lepidopterans like *Spodoptera exigua* (Hübner) and *S. litura* (F.) (Lepidoptera: Noctuidae) are well-known economic importance insect pests that defoliate a variety of economically important crops. Currently, the most widely used control for both is the use of insecticides. Applying large amounts...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Effectiveness of Bio-Catharantin Induction to Increase Red Spinach (*Alternanthera amoena* Voss.) Production

Nabila Shafura, Laras Nur Janah, Muhammad Syafi'atol Huda, Budi Setiadi Daryono

Red spinach (*Alternanthera amoena* Voss.) is a plant with high nutritional value and contains anthocyanin compounds that act as antioxidant compounds to prevent the formation of free radicals in the body, besides that anthocyanin can also treat anemia. The increase in demand for red spinach in the market...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Effects of High-Fat Diet and CCl₄ Administration on Liver Function and Lipid Profile in Non-Alcoholic Fatty Liver Disease Rat Model

Marwan Rosada, Widya Wasityastuti, Yanasta Yudo Pratama, Ken Siwi, Dewiyaning Indah Widasari, Tutik Sri Wahyuni

Non-alcoholic fatty liver disease (NAFLD) is often encountered in the field of hepatology. The disease has a broad spectrum ranging from non-inflammatory fat-accumulating macrovesicles (simple steatosis) and develops to fibrous and cirrhosis. The rate of NAFLD is 15-30% in western countries and around...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Reproductive Aspect and Embryonic Development of Wader pari Fish (*Rasbora lateristriata* Bleeker 1854) from Malang East Java

Hilyatuz Zahro, Khoirudin Anshori, Sandi Fransisco, Amalia Audina Rosa, Bambang Retnoaji

Indonesia is well known for its high freshwater fish diversity. Wader pari (*Rasbora lateristriata*) is one of the endemic fish, which is very popular, but experienced massive exploitation in wild, due to high market demand. High demand and high economic value of this fish in the market induced a massive...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Cytological Analysis of *Aerides odorata* Lour. from Sleman, Special Region of Yogyakarta

Kireida Asta Nugraheni, Febri Yuda Kurniawan, Anindita Della Rosa Riyadi, Anggiresti Kinasih, Hadyan Pratama Lutfi Ilmam, Endang Semiarti

Aerides odorata is a species that belongs to the Orchidaceae family and lives as epiphytic orchids. This orchid is distributed across Southeast Asia, India, and China. This orchid occupies one type of habitat. Morphological characters of flowers in this species vary, depending on the habitat occupied....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Reproductive Aspect and Embryonic Development of Wader

Fish (*Rasbora lateristriata* Bleeker, 1854) from Purworejo, Central Java

Inayah Rizkia Lailiati, Devi Annisa Suci, Amalia Audina Rosa, Veronica Aurelia Fernanda, Bambang Retnoaji

Indonesian native fish "Wader pari" (*Rasbora lateristriata*) from Purworejo is one of the fish has high economic value, which is related to the community enthusiasts and high requests for the fish stock. The demand for fish availability and supply is increasing drastically, which caused fish population...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

In Vitro Selection and Total Phenol Analysis of Moon Orchid [*Phalaenopsis amabilis* (L.) Bl.] Results of Induced With Fusaric Acid

Endang Nurcahyani, Risma Rasmani, Hardoko Insan Qudus

Phalaenopsis amabilis (L.) Bl. is an orchid that is included in the list of endangered plant species. The Fusarium wilt disease on *P. amabilis* is a vital disease caused by *Fusarium oxysporum*, which of one constraint and up to now days is not well yet managed. Disease control that does not cause negative...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Anti-Infective Properties of a Sea Cucumber Associated Actinobacteria *Kocuria* sp. HL 55

Joko Tri Wibowo

“Antibiotics golden era” can end when resistant strains spread immensely. To find new anti-infective agents, we examined sea cucumber associated bacteria isolated from the intestinal part of the echinoderm *Holothuria leucospilota*. Partial identification using 16S rRNA gene Sanger sequencing revealed...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Study of N, P, K, and C on Degradation of Indigosol Batik Dye Effluent by *Aspergillus* sp. GPN

Ratna Stia Dewi, Mardiyah Kurniasih

Indigosol dye batik effluent is toxic since it excesses contain decreasing agent sodium hydrosulfite that is oxidized to alkali and alkaline earth metal sulfate (SO_4^{2-}), sulfite (SO_3^{2-}), and thiosulfate ($\text{S}_2\text{O}_3^{2-}$) that very corrosive. Therefore, the role of environmental biotechnology in the technology...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Availability and Information Needs of Traditional Medicine in Urban Community, Surabaya, Indonesia

Oeke Yunita, Fernanda Rizky Putri Heriwana, Erlin Theterissa, Jimmy Jimmy

Indonesian traditional medicine has been widely used in the urban community to treat several symptoms and prevent diseases before accessing the hospital for professional help. Information is critical in modern culture, since it aids in the development of people's health knowledge. The purpose of this...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Effect of Bokashi and Rabbit Urine Addition on The Tubber of Shallots (*Allium ascalonicum* L.)

Muh. Alwi Husen, Sugiyarto Sugiyarto, Esna Dilli Novianto

Shallots (*Allium ascalonicum* L.) have been cultivated intensively in Indonesia for a long time. These horticultural crops have high potential and economic value. However, the domestic production itself is not able to meet national demands. Continuous use of chemical agents will negatively impact the...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

The Effect of Variations in Sugar Types and Fermentation Time on Enzyme Activity and Total Titrated Acid on Eco-Enzyme Results of Fermentation

Imam Abu Hanifah, Ni Putu Vidya Primarista, Sasangka Prasetyawan, Anna Safitri, Tri Adyati, Arie Srihadyastutie

Eco-enzyme (EE) production is an alternative for reducing organic waste such as fruit peels and vegetables. EE is widely used as natural organic fertilizer, purification of contaminated groundwater, pesticides, herbicides, and natural insecticides. EE is made through a fermentation process for 3 months...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Ethnobotanical Study of Hindu Society in Tabanan Bali and The Conservation Efforts

Putri Sri Andila, Tri Warseno, Winda Syafitri, I Gede Tirta

Balinese have traditionally preserved plants for the ethnobotany used hereditary, such as the utilization of plants for traditional medicine and Hindu ceremonies. Most of the Tabanan people are still live in the villages traditionally. They have used plants for the traditional purpose for hundreds of...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Diversity and Estimated Above Ground Biomass of Shade Trees in Some Coffee-based Agroforestries, Banyuwangi Regency

Jehan Ramdani Hariyati, Dian Siswanto, Endang Arisoesilaningsih, Luchman Hakim

This study was aimed to assess the diversity and estimate the above-ground biomass (AGB) of the shade tree in several coffee-based agroforestries in Banyuwangi, East Java. The data was collected from three villages, i.e. Gombengsari, Papring, and Kependukuh. Vegetation analysis and biomass measurement...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Initial Survey of Tadpole Species Richness on the Upstream

of Pelus River, Lompakuwus, Sumbang, Banyumas

I Gusti Agung Ayu Ratna Puspitasari, Hafizh Aulia Khairy Rakananda, Nugroho Dwi Septianto, Meyta Pratiwi, Eko Setio Wibowo

The southern slopes of Mount Slamet are the upstream areas of many rivers, which become the breeding habitat of Anuran. Anuran tadpole species richness can indicate the diversity of Anuran in the area and should be considered as important as the adult Anuran species richness. Moreover, the sampling of...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Dhole's Ecology and Legal Status: Conservation and Management Implication in Human-Dominated Landscape of Java, Indonesia

Sandy Nurvianto, Muhammad Ali Imron, Sven Herzog

The success of conservation and management of endangered species is highly dependent on the manager's understanding on the biology and ecology of the target species. However, most wildlife management and conservation activities in developing countries are not based on scientific data, but more on the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Case Report: Therapy of Streptococcal Pneumonia in a Cat

Riskha Nurmi Nataria, Soedarmanto Indarjulianto, Yanuartono, Alfarisa Nururrozi, Slamet Raharjo, Hary Purnamaningsih, Heldiar Soedarmanto, Puveanthan Nagappan Govendan

Pneumonia is a lung infection that can be caused by bacteria, including Streptococcus sp. Identification of the cause of pneumonia in cats is very necessary so that diagnosis and therapy will be more precise. This study reports on the diagnosis and therapy of streptococcal pneumonia on a cat. A 10-month-old...

Atlantis Press

+ Article details

Atlantis Press – now part of Springer Nature – is a professional publisher

+ Download article (PDF)

of scientific, technical & medical (STM) proceedings, journals and books.

We offer world-class services, fast turnaround times and personalised

communication. The proceedings and journals on our platform are Open

Proceedings Article millions of downloads every month.

Comparison of Cycle Threshold Value of The Orf 1 ab SARS-CoV-2 Gene from Three Different PCR Reagents

Anita L. Susanti, Fusvita Merdekawati, Rohayati Rohayati

► PROCEEDINGS

► ABOUT

The availability of PCR reagents is an obstacle that still needs to be resolved

► JOURNALS

► NEWS

in the early days of the SARS-CoV-2 pandemic. During the early days of the

► BOOKS

► CONTACT

pandemics, identical PCR reagents supply in the laboratory could not be

guaranteed. The laboratory needs to investigate the reliability of the

di-

► MANAGE COOKIES/DO NOT SELL MY

+ INFO

+ Download article (PDF)

Home Privacy Policy Terms of use



Proceedings Article

Copyright © 2006-2022 Atlantis Press – now part of Springer Nature

Antioxidant Activity, Total Phenolic Content, and Chemical Composition of Liquid Smoke Derived from Wood Sawdust

Ria Suryani, Rafelinta Daradwinta, Syahida Az-Zahra, Syifa' Aulia Rahmah, Wahyu Anggo Rizal, Tri Hadi Jatmiko, Wuri Apriyana, Yekti Asih Purwestri

Liquid smokes result from the process of pyrolysis of biomass as a raw material which generally contains high content of lignin, cellulose, and hemicellulose. The phenolic compounds, acid compounds, and carbonyl compounds are the main components in liquid smoke and show antioxidant activity. One of the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

The Potency of Red Betel (*Piper crocatum* Ruiz & Pav.) Methanolic Extract as α -Amylase and α -Glucosidase Inhibitor

Yustina Sri Hartini^{1,*}, Dewi Setyaningsih¹

¹ Faculty of Pharmacy Sanata Dharma University, Paingan Maguwoharjo Depok Sleman Yogyakarta, Indonesia

*Corresponding author. Email: yustinahartini@usd.ac.id

ABSTRACT

Methanol is an effective menstruum for attracting compounds with various pharmacological activities from *Piper crocatum* Ruiz & Pav. Several studies reported that laboratory tests for reducing blood sugar content of red betel extract such as aqueous, ethanolic, and ethyl acetate extracts. This study aimed to investigate the actions of red betel leaves methanolic extract in reducing α -amylase and α -glucosidase in an *in-vitro* study. The test for decreasing the actions of the enzymes was done using the ultraviolet-visible spectrophotometric method. The test was run by observing carbohydrates left by knowing the effect of iodine-iodide resulting in a blue complex. The red betel leaves extract demonstrated an adequate α -amylase and α -glucosidase inhibitory activity as shown by the IC₅₀ of 8.463 ± 0.318 mg/mL and 10.013 ± 0.070 mg/mL, respectively.

Keywords: α -amylase, α -glucosidase, Methanolic extract, *Piper crocatum*.

1. INTRODUCTION

It has been reported that some medicinal plants species have hypoglycaemic activity. Most plants contain bioactive components, such as polyphenols, alkaloids, terpenoids, flavonoids, coumarins, and other constituents [1]. Variations of phytochemical classes and bioactive compounds' chemical structure may differ in action mechanisms on reducing blood glucose [2]. The imbalance between blood sugar absorption and insulin production resulted in diabetes mellitus type 2.

Controlling blood sugar levels could be done by inhibiting the activity of hydrolytic enzymes that digest dietary starch, such as α -amylase and α -glucosidase enzymes. Methanol, ethanol, or a mixture of alcohol and water are common solvents used to extract bioactive compounds from the plant, also known as the menstruum [3,4]. The literature reported that methanol is an effective menstruum for attracting compounds with various pharmacological activities from red betel (*Piper crocatum* Ruiz & Pav.). Furthermore, Pandithurai *et al.* [7] found a methanolic extract of marine brown algae showed the reduction of α -amylase and α -glucosidase

actions [5,6,7]. Several studies reported that the laboratory test of diabetic medicines from *P. crocatum* compounds such as aqueous, ethanolic, and ethyl acetate extracts [8,9,10,11].

The study aimed to evaluate the antidiabetic potency of red betel leaves methanolic extract activity through the α -amylase and α -glucosidase inhibition and identify the extracted compound. Methanol is expected to be able to extract antidiabetic compounds from red betel. The success of the extraction of these antidiabetic compounds was measured by the ability of *P. crocatum* methanol extract to reduce the actions of the α -amylase and α -glucosidase enzymes.

2. METHODOLOGY

The maceration method was used in the compound extraction of red betel leaves. The red betel leaves methanolic extract compounds profile was checked using Thin Layer Chromatography (TLC). Compound separation of the extracts has been detected by three types of mobile phase composition; n-hexane:ethyl acetate (3:1), chloroform:ethyl acetate (1:1), and toluene:ethyl

acetate (2:1). Spots detection of the separated compounds was carried out by spraying Lieberman Burchard's reagent, cerium sulfate, iodine, FeCl_3 , and observing under UV-Visible spectrophotometer (254 nm and 366 nm). Assay of α -amylase enzyme inhibitory activity was done following Ononamadu *et al.* [12]. The potato powder (1% w/v), 1.0 ml extract/1 ml acarbose, 1.0 ml of the α -amylase enzyme (1% w/v), and 2.0 ml of acetate buffer (0.1M, 7.2 pH) were blended together. Inhibitory effect of the sample blank solution measurement was done by pouring 1.0 ml of 0.5% (w/v) potato starch solution into a test tube. After one hour incubation of the mixture, a 0.1 ml iodine-iodide was then poured.

The α -amylase enzyme inhibitory activity test was done following Pandhithurai *et al.* [7] with slightly modified. Extract and standard solutions of 100.0 μL from each series of test solutions were added with 400.0 μL of phosphate buffer and 250.0 μL of maltose substrate solution into a test tube and then pre-incubated at 37 °C for 5 minutes. After pre-incubation completion, added 250.0 μL of phosphate buffer solution pH 7.0 and then homogenized. Then the solution was incubated for 30 minutes at 37 °C. 0.3 mL of the solution was then taken, and 0.3 mL of DNS reagent was added to the test tube. The solution was homogenized and heated in boiling water for 5 minutes. The solution is mixed by adding 3.0 mL of distilled water. The absorbance was detected using a UV-vis spectrophotometer at 536 nm. The percent activity reduction was counted using Equation (1), where Ac means the absorbance provided by the control experiment, while As is the tested sample absorbance

$$\% \text{ inhibition} = (\text{As}-\text{Ac}/\text{As}) \times 100 \quad (1)$$

The results were statistically evaluated for ANOVA ($p < 0.05$) and continued by the Tukey's Post-Hoc Test ($\alpha=0.05$) using an IBM SPSS version.

3. RESULTS AND DISCUSSION

Red betel leaves methanolic extract inhibited hydrolytic enzymes, both α -amylase and α -glucosidase. The red betel leaves extract significantly increased enzyme inhibition activity against the α -amylase enzyme ($P < 0.05$). The red betel percent inhibition of α -amylase was found to range from 19.44 to 55.56%, while the percent inhibition of α -glucosidase was found to range from 12.24 to 49.02% (Figure 1). The values of their IC_{50} for both α -amylase and α -glucosidase reduction due to the extract of *P. crocatum* leaves were 8.463 ± 0.318 mg/mL and 10.013 ± 0.070 mg/mL, respectively. As the positive control used in this study, acarbose showed IC_{50} value was 0.837 ± 0.076 mg/mL for α -amylase inhibitory and 0.690 ± 0.124 mg/mL for α -glucosidase inhibitory activities. It means that the acarbose elicited greater percentage inhibition value than *P. crocatum* leaves methanolic extract for inhibiting the α -amylase and α -glucosidase. The IC_{50} value of acarbose was. In acarbose,

the concentration to reduce α -amylase was greater than those of α -glucosidase, but the opposite occurs in *P. crocatum* leaves methanolic extract.




Figure 1. Percent enzymes inhibition activity of *P. crocatum* methanolic extract and acarbose: Pc-amylase: *P. crocatum* methanolic extract against α -amylase enzyme; Pc-glucosidase: *P. crocatum* methanolic extract against α -glucosidase enzyme; Ac-amylase: Acarbose against the α -amylase enzyme, and Ac-glucosidase: Acarbose against the α -glucosidase enzyme.

Several researchers tested the antidiabetic activity of plants against α -amylase or α -glucosidase separately or both [9, 11, 13, 7]. It has been observed that the ethanolic extract of 14 medicinal plants reduced the activities of α -amylase and α -glucosidase, and α -amylase was not affected by plant extract concentration [13]. The same compound did not reduce α -amylase action as occurred in α -glucosidase. The compound mixture from plant extracts could act synergist, additive, or antagonistic. If the mixture compounds interact synergistically, it seems that they are more effective than those of pure compounds [14].




Figure 2. Thin Layer Chromatogram of *P. crocatum* methanolic extract observed under 254 nm (1), extract soluble in n-hexane (2), extract insoluble in n-hexane (3) in mobile phase composition of n-hexane:ethyl acetate (3:1) [A], chloroform:ethyl acetate (1:1) [B], and toluene:ethyl acetate (2:1)[C].

The thin layer chromatography mobile phase composition in separating process of compounds in red betel leaves methanolic extract were n-hexane:ethyl acetate (3:1); chloroform:ethyl acetate (1:1) and toluene:ethyl acetate (2:1). Each mobile phase type used in this study separated the compounds in the extract (Figure 2). Further fractionation of the methanol extract was carried out using n-hexane. Some of the red betels leaves methanolic extract compounds were extracted in the soluble and insoluble fractions of n-hexane. However, in general, the n-hexane solvent was quite effective in separating these compounds.

In addition to methanol, compounds extracted with other polar solvents such as water showed antidiabetic activity from the extract. Some phenolic compounds, namely cyanidin 3-O-glucoside, caffeic acid, tannin, p-coumaric acid, and gallic acid are identified for antidiabetic activity [10]. Hartini *et al.* [5] fractionated methanol extract using vacuum liquid chromatography with various compositions of n-hexane and ethyl acetate, detected the presence of essential oils, tannins, alkaloids, terpenoids, and flavonoids. Tannin has been reported as an α -amylase and α -glucosidase enzyme inhibitor [13, 15]. Tropical almonds, petroleum ether, and methanolic extracts are rich in tannins with antidiabetic activity [16]. Chemically tannins are complex substances, and they usually occur in mixtures of polyphenol that are difficult to separate because they do not crystallize.

According to Barret *et al.* [15], the effectiveness of tannins in inhibiting the enzyme's activity was affected by molecule size and structural complexity. In addition to tannins, other groups of compounds reduced enzyme activity. The ethyl acetate fraction of red betel leaves extract contains the terpenoid (Columbine), which was stronger in inhibiting the activity of the α -glucosidase enzyme than other compounds [9]. Many factors seem to affect extract activity in reducing α -glucosidase, i.e., harvesting time, storage conditions, cultivation, climate, processing, and genetic background [10]. Red betel leaves methanolic extract can be fractionated with non-polar solvents such as n-hexane to obtain the compound responsible for the reduction of α -amylase and α -glucosidase enzymes activities. In view of tannins' reported solubility and activity, further research is needed on the activity of compounds in the insoluble n-hexane fraction and the soluble n-hexane fraction in inhibiting the activity of α -amylase and α -glucosidase enzymes.

The antidiabetic ingredients of red betel leaves can be extracted using methanol. Red betel leaves methanolic extract demonstrated inhibitory activities for the α -amylase or α -glucosidase enzyme. Fractionation using n-hexane as solvent effectively separated the compounds in red betel leaves methanolic extract. It needs further investigation to confirm the activity of the soluble and insoluble fractions of n-hexane to ascertain which compound or group of compounds are responsible for the

reduction of α -amylase or α -glucosidase enzymes activity from red betel leaves methanolic extract.

AUTHORS' CONTRIBUTIONS

Yustina Sri Hartini was responsible for the whole experiment, including expressing the idea and designing the experiments, preparing material, reagents, tools, performing the experiments, and writing the paper. Dewi Setyaningsih contributed to reagents provision, performing the experiment, analyzing, and interpreting the data.

ACKNOWLEDGMENTS

Our gratitude to LPPM Universitas Sanata Dharma for funds supporting this research.

REFERENCES

- [1] D.K. Patel, S.K. Prasad, R. Kumar, S. Hemalatha, An overview on antidiabetic medicinal plants having insulin mimetic property, Asian Pac. J. Trop. Biomed vol. 2, 2012, pp. 320-330. DOI: [https://doi.org/10.1016/S2221-1691\(12\)60032-X](https://doi.org/10.1016/S2221-1691(12)60032-X)
- [2] O. Patience, Osadebe, U. Estella, Odoh, F. Philip, Uzor, The search for new hypoglycemic agents from plant, African Journal of Pharmacy and Pharmacology vol 8, 2014, pp. 292-303. DOI: <https://doi.org/10.5897/AJPP2014.3933>
- [3] S. Sasidharan, Y. Chen, D. Saravanan, K.M. Sundram, L.Y. and Latha, Extraction, Isolation and Characterization of Bioactive Compound from Plants' Extracts, Afr. J Tradit Complement Altern Med vol. 8, 2011, pp. 1-10. DOI: [10.4314/ajtcam.v8i1.60483](https://doi.org/10.4314/ajtcam.v8i1.60483)
- [4] A.R. Abubakar and M. Haque, Preparation of medicinal plants: Basic extraction and fractionation procedures for experimental purposes, Journal of Pharmacy & Bioallied Sciences vol. 12, 2020, pp. 1-10. DOI: https://doi.org/10.4103/jpbs.JPBS_175_19
- [5] Y.S. Hartini, S. Wahyuono, S. Widayarni, A. Yuswanto, Phagocytic Macrophage Activity of Fractions from Methanolic Leaf Extract of Red Betel (*Piper crocatum* Ruiz & Pav.) In Vitro, Jurnal Ilmu Kefarmasian Indonesia vol. 11, 2013, pp. 108-115. DOI: <https://doi.org/10.4314/tjpr.v13i10.6>
- [6] B.D. Wicaksono, Y.A. Handoko, E.T. Arung, I.W. Kusuma, D. Yulia, A.N. Pancaputra, Antiproliferative effect of metanol extract of *Piper crocatum* Ruiz & Pav leaves on human breast (T47D) cells *in vitro*, Tropical Journal of Pharmaceutical Research vol. 8, 2009, pp. 345-352. DOI: <https://doi.org/10.4314/tjpr.v8i4.45229>
- [7] M. Pandithurai, S. Murugesan, S. Bhuvaneswari, S. and Thennarasan, In vitro α -amylase and α -

- glucosidase inhibition activity of methanolic extract of marine brown alga *Spatoglossum asperum*. International Journal of Advances in Pharmaceutics vol. 4, 2015, pp. 83-87. DOI: <https://doi.org/10.7439/ijap>.
- [8] M. Safitri and F. Fahma, 2008 Potency of *Piper crocatum* Decoction as an Antihyperglycemia in Rat Strain Sprague dawley, Hayati Journal of Biosciences vol. 15, 2008, pp. 45-48. DOI: <https://doi.org/10.4308/hjb.15.1.45>
- [9] M. Weni, M. Safithri, and D.S.H. Seno, 2020 Molecular Docking of Active Compounds *Piper crocatum* on The Alpha-Glucosidase Enzyme as Antidiabetic, Indonesian Journal of Pharmaceutical Science and Technology, vol. 7, No. 2, 2020, pp. 64-72. DOI: <https://doi.org/10.24198/ijpst.v7i2.21120>
- [10] S.R.S. Kamaruzaman, K.F. Kasim, and M.N. Jaafar, The Effect of Harvesting Time on the Antioxidant and Antidiabetic Activity of *Piper crocatum* (Sirih Merah) Extract. IOP. Series: Material Science and Engineering vol. 864, IOP Publishing, 2020. DOI: [10.1088/1757-899X/864/1/012211](https://doi.org/10.1088/1757-899X/864/1/012211).
- [11] Y.S. Hartini, D. Setyaningsih, M.J.V. Chang, and M.C.I.A. Nugrahanti, Sambiloto (*Andrograpis paniculata* Nees) leaf extract activity as an α -mylase enzyme inhibitor. Pharmacy Education vol. 21, 2021, pp. 305-308. DOI: <https://doi.org/10.46542/pr.2021.212.305308>
- [12] C.J. Ononamadu, O.C. Ezeigwe, T.A. Owolarafe, G.O. Ihegboro, T.A. Lawal, K. Salawu, U.F. Umeoguaju, and I. Aminu, Starch-iodine assay method underestimates -amylase inhibitory potential of antioxidative compound and extracts. BioTechnologia vol. 101, 2020, pp. 45-54. DOI: [http://doi.org/10.5114/bta.2020.93103](https://doi.org/10.5114/bta.2020.93103)
- [13] B. Elya, R. Handayani, R. Sauriasari, U.S. Hasyati, I.T. Permana, and Y.I. Permatasari, Antidiabetic Activity and Phytochemical Screening of Extracts from Indonesian Plants by Inhibition of Alpha Amylase, Alpha Glukcosidase and Dipertidyl Peptidase IV, Pakistan Journal of Biological Sciences vol. 18, 2015, pp. 279-284. DOI: <http://doi.org/10.3923/pjbs.2015.279.284>
- [14] L.K. Caesar and N.B. Cech, Synergi and Antagonism in Natural Product Extracts: When 1+1 does not equal 2, Natural Product Reports vol. 36, 2019, pp. 845-936. DOI: <http://doi.org/10.1039/c9np0011a>
- [15] A. Barret, T. Ndou, C.A. Hughey, C. Straut, A. Howell, Z. Dai, and G. Kaletunc, Inhibition of α -amylase and glucoamylase by tannins extracted from cocoa, pomegranates, cranberries and grapes. J. Agric. Food Chem. vol. 61, 2013, pp. 1477-1486. DOI: <http://doi.org/10.1021/jf304876g>
- [16] A.N. Nagappa, P.A. Thakurdesai, R.N. Venkat, S. Jiwan, Antidiabetic activity of *Terminalia catappa* Linn fruits, J Ethnopharmacol vol. 88, 2003, pp. 45-50. DOI: [http://doi.org/10.1016/s0378-8741\(03\)00208-3](http://doi.org/10.1016/s0378-8741(03)00208-3)