

## Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Enny Murwaningtyas

Assignment title: Periksa similarity

Submission title: European option pricing by using a mixed fractional Browni...

File name: n\_option\_pricing\_by\_using\_a\_mixed\_fractional\_Brownian\_mo...

File size: 898.46K

Page count: 9

Word count: 3,335

Character count: 15,360

Submission date: 10-Mar-2023 02:34PM (UTC+0700)

Submission ID: 2033743074



# European option pricing by using a mixed fractional Brownian motion

by Murwaningtyas Enny

**Submission date:** 10-Mar-2023 02:34PM (UTC+0700)

**Submission ID: 2033743074** 

**File name:** n\_option\_pricing\_by\_using\_a\_mixed\_fractional\_Brownian\_motion.pdf (898.46K)

Word count: 3335

Character count: 15360

#### **PAPER · OPEN ACCESS**

#### European option pricing by using a mixed fractional Brownian motion

To cite this article: C E Murwaningtyas et al 2018 J. Phys.: Conf. Ser. 1097 012081

View the article online for updates and enhancements.

- You may also like
  20
   Inhibitory effect induced by fractional
  Gaussian noise in neuronal system
  Zhi-Kun Li , , Dong-Xi Li et al.
  10
  - Spectral content of fractional Brownian
    10 on with stochastic reset
    Satva N Majumdar and Gleb Oshanin
  - THE DEPENDENCE OF PRESTELLAR CORE MASS DISTRIBUTIONS ON THE STRUCTURE OF THE PARENTAL CLOUD

    Antonio Parravano, Nestor Sánchez and

Emilio J. Alfaro



## European option pricing by using a mixed fractional Brownian motion

#### C E Murwaningtyas 12), S H Kartiko 1), Gunardi 1) and H P Suryawan 3)

- 1) Department of Mathematics, Universitas Gadjah Mada, Indonesia,
- 2) Department of Mathematics Education, Universitas Sanata Dharma, Indonesia,
- 3) Department of Mathematics, Universitas Sanata Dharma, Indonesia.

Corresponding author: enny@usd.ac.id

Abstract. Financial modeling is conventionally based on a Brownian motion (Bm). A Bm is a semimartingale process with independent and stationary increments. However, some financial data do not support this assumption. One of the models that can overcome this problem is a fractional Brownian motion (fBm). In fact, the main problem in option pricing by implementing an fBm is not arbitrage-free. This problem can be handled by using a mixed fBm (mfBm) to model stock prices. The mfBm is a linear combination of an fBm and an independent Bm. The aim of this paper is to find European option pricing by using the mfBm based on Fourier transform method and quasi-conditional expectations. The main result of this research is a closed form formula for calculating the price of European call options.

#### 1. Introduction

The Black-Scholes formula is a formula for calculating option prices based on geometric m. A Bm is a centered and continuous Gaussian process with independent and stationary increments. The existence of long-range dependence in essential topic of both empirical and theoretical research. If stock returns is show long-range dependence, the time series is said to depend on time to time for a long lag. This is the case of an fBm. Long-range dependence in stock returns has been tested in a number of studies, for example [1–6].

Kolmogorov introduced an fBm in 1940. Mandelbrot and Van Ness gave a representation theorem for Kolmogorov's process and introduced the name of fBm in [7]. The fBm has further been developed by Hurst in [8]. Currently, an fBm has an important part in assorted fields of study such as hydrology [8,9], insurance [10,11] and finance [12–14].

The stochastic integral in an fBm is different from the classical Itô integral because the fBm is not a martingale. Duncan et al [15] introduced a Wick product for the fractional Itô's formula. They also introduce  $\frac{1}{26}$  Girsanov's theorem under the fBm. The option model under the fBm is arbitrage-free [16,17], where  $\frac{1}{26}$  e Wick product is applied on the definition of stochastic integration. Hu and Oksendal [16] obtain a pricing formula for a European call option at t = 0. Necula [18] extended the formula in [16] to  $t \in [0,T]$ . Moreover, Necula proved some results regarding quasi-conditional expectations by using Fourier transform.

The European call option pricing formula obtained in [16] is an arbitrage-free and complete market. However, Bender and Elliott [19] and Bjork and Hult [20] still saw a possibility of arbitrage

content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

opportunities in the resulting model [36] [16]. Cheridito [21] and Bender et al. [22] proposed an mfBm to reduce arbitrage opportunities. An mfBm is a linear combination of an fBm and an independent BM. Cheridito [23] has proven that an mfBm is equivalent to a Bm for  $H \in (\frac{3}{4}, 1)$ , therefore it can be said that the option model under an mfBm is an arbitrage-free. The aim of this paper is to obtain the pricing formula for European call options where a stock return is modeled an mfBm.

Mixed fractional Brownian motions
Let H be a constant belonging to (0, 1). An fBm  $B^H = (B_t^H; t \ge 0)$  of Hurst index H is a continuous and centered Gaussian process with covariance function,

$$\mathbb{E}\left[B_{t}^{H}B_{s}^{H}\right] = \frac{1}{2}\left(\left|t\right|^{2H} + \left|s\right|^{2H} - \left|t - s\right|^{2H}\right),\tag{1}$$

for all  $t,s \ge 0$ , see [24]. Here  $\mathbb{E}[\cdot]$  denotes an expectation with respect to a probability measure  $\mathbb{P}^H$ . Properties of fBm, see [24], are

- mean of an fBm is 0;
- variance of an fBm is  $t^{2H}$  for  $t \ge 0$ ;
- an fBm has stationary increments, i.e.,  $B_{t+s}^H B_s^H \stackrel{d}{=} B_t^H$  for all  $t, s \ge 0$ ;
- an fBm is *H*-self similar, i.e.,  $B_{ct}^H = \alpha^H B_t^H$  for  $t \ge 0$ ;
- an fBm has continuous trajectories.

If  $H = \frac{1}{2}$ , then an fBm coincides with a standard Bm. The Hurst index H determines the sign of the covariance of the future and past increments. This covariance is negative when  $H \in (0, \frac{1}{2})$ , zero when  $H = \frac{1}{2}$ , and positive when  $H \in (\frac{1}{2}, 1)$ . As a consequence, for  $H \in (0, \frac{1}{2})$  it has short-range dependence and for  $H \in (\frac{1}{8}, 1)$  it has long-range dependence.

An fBm is neither a semimartingale nor a Markov process unless  $H = \frac{1}{2}$ . When H is not equal to  $\frac{1}{2}$ , the option model has arbites opportunities. An mfBm is introduced by Cheridito [23], to avoid arbitrage opportunities. An mfBm of parameter H, a, and b is a stochastic process  $M^H = (M_t^H; t \ge 0)$ defined in [25] as follows

$$M_{t}^{H} = aB_{t}^{H} + bB_{t}$$
where  $B_{t}^{H}$  is an fBm with Hurst index H and  $B_{t}$  is an independent BM.

#### 3. Quasi-conditional expectations

We will present some results regarding a quasi-conditional expectation in this section which is needed for the rest of this paper. These results were introduced by Necula [18] and then developed by Sun [14] and Xiao et 🔠 [13] for an mfBm. The proofs of theorems in this section can be seen in [13]. Let  $(\Omega, \mathcal{F}^H, \mathbb{P}^H)$  be a probability space such that  $B_i^H$  is an fBm with respect to  $\mathbb{P}^H$  and  $B_i$  is an independent BM.

**Theorem 1** [13]. For every  $t \in (0,T)$  and  $\lambda, \varepsilon \in \mathbb{C}$  we have

$$\tilde{\mathbb{E}}\left[\exp\left(\lambda\varepsilon B_{T}^{H}+\lambda B_{T}\right)\middle|\mathcal{F}_{t}^{H}\right]=\exp\left(\lambda\varepsilon B_{T}^{H}+\lambda B_{T}+\frac{1}{2}\lambda^{2}\varepsilon^{2}\left(T^{2H}-t^{2H}\right)+\frac{1}{2}\lambda^{2}\left(T-t\right)\right),$$

where  $\mathcal{F}_{t}^{H}$  is a  $\sigma$ -algebra generated by  $\left(B_{s}^{H};0\leq s\leq t\right)$  and  $\tilde{\mathbb{E}}\left[\cdot\middle|\mathcal{F}_{t}^{H}\right]$  denotes a quasi-conditional expectation with respect to  $\mathcal{F}_t^H$  under a probability measure  $\mathbb{P}^H$  .

Using Theorem 1, one can determine a quasi-conditional expectation of a function of an mfBm as shown in the theorem below.

**Theorem 2** [13]. Let f be a function such that  $\tilde{\mathbb{E}}\Big[f\Big(B_T^H, B_T\Big)\Big] < \infty$ . Then, for every  $t \in (0,T)$  and  $\lambda, \varepsilon \in \mathbb{R}$  we have

$$\mathbb{\tilde{E}}\left[f\left(\lambda\varepsilon B_{T}^{H} + \lambda B_{T}\right)\middle|\mathcal{F}_{t}^{H}\right] = \int_{\mathbb{R}} \frac{\exp\left(\frac{-\left(y - \lambda\varepsilon B_{t}^{H} - \lambda B_{t}\right)^{2}}{2\left(\lambda^{2}\varepsilon^{2}\left(T^{2H} - t^{2H}\right) + \lambda^{2}\left(T - t\right)\right)}\right)}{\sqrt{2\pi\left(\lambda^{2}\varepsilon^{2}\left(T^{2H} - t^{2H}\right) + \lambda^{2}\left(T - t\right)\right)}}f(y)dy.$$

If f is an indicator function,  $f(y) = 1_A(y)$ , then we can easily obtain a corollary below.

**Corollary** 3 [13]. Let  $A \in \mathcal{B}(\mathbb{R})$ . Then,

$$\widetilde{\mathbb{E}}\left[1_{A}\left(\lambda\varepsilon B_{T}^{H}+\lambda B_{T}\right)\middle|\mathcal{F}_{t}^{H}\right]=\int_{A}\frac{\exp\left(\frac{-\left(y-\lambda\varepsilon B_{t}^{H}-\lambda B_{t}\right)^{2}}{2\left(\lambda^{2}\varepsilon^{2}\left(T^{2H}-t^{2H}\right)+\lambda^{2}\left(T-t\right)\right)}\right)}{\sqrt{2\pi\left(\lambda^{2}\varepsilon^{2}\left(T^{2H}-t^{2H}\right)+\lambda^{2}\left(T-t\right)\right)}}dy.$$

Let  $\theta, \theta \in \mathbb{R}$  and  $t \in [0,T]$ , consider the process,

$$\theta B_t^{H^*} + \vartheta B_t^* = \theta B_t^H + \theta^2 t^{2H} + \vartheta B_t + \vartheta^2 t. \tag{2}$$

From a fractional Girsanov theorem in [24], there exists a probability measure  $\mathbb{P}^{H^*}$  such that  $\theta B_t^{H^*} + \theta B_t^*$  is a new MFBM. We will denote  $\tilde{\mathbb{E}}^* \left[ \cdot \middle| \mathcal{F}_t^H \right]$  as a quasi-conditional expectation under the probability measure  $\mathbb{P}^{H^*}$ . Now, we have defined the process

$$Z(t) = \exp\left(-\theta B_t^H - \frac{1}{2}\theta^2 t^{2H} - \theta B_t - \frac{1}{2}\theta^2 t\right),\tag{3}$$

where  $t \in [0,T]$ .

**Theorem 4** [13]. Let f be a function such that  $\mathbb{E}\left[f\left(B_T^H, B_T\right)\right] < \infty$ . Then, for every  $t \in [0,T]$  we have

$$\tilde{\mathbb{E}}^* \left[ f \left( \theta B_T^H + \mathcal{G} B_T \right) \middle| \mathcal{F}_t^H \right] = \frac{1}{Z(t)} \tilde{\mathbb{E}} \left[ f \left( \theta B_T^H + \mathcal{G} B_T \right) Z(T) \middle| \mathcal{F}_t^H \right]. \tag{4}$$

Theorem 4 illustrates a relationship between a quasi-conditional expectation  $\tilde{\mathbb{E}}[\cdot|\mathcal{F}_t^H]$  with respect to  $\mathbb{P}^H$  and a quasi-conditional expectation  $\tilde{\mathbb{E}}^*[\cdot|\mathcal{F}_t^H]$  with respect to  $\mathbb{P}^{H^*}$ . Based on Theorem 4, a discounted expectation of a function of an mfBm is calculated in the theorem below.

**Theorem 5** [13]. The price at time  $t \in [0,T]$  of a bounded  $\mathcal{F}_t^H$ -measurable claim  $V \in L^2(\mathbb{P}^H)$  is given by

$$V_{t} = e^{-r(T-t)} \widetilde{\mathbb{E}} \left[ V_{T} \middle| \mathcal{F}_{t}^{H} \right], \tag{5}$$

where r is a constant riskless interest rate

#### 4. Results and discussion

aim of this section is to determine a formula for calculating European call option prices. Now let us consider a mixed fractional Black-Scholes market with two investment possibilities:

A bank account which satisfies a differential egsation below

$$dA_{t} = rA_{t}dt, A_{0} = 1, t \in [0,T],$$
 (6)

where r is a constant riskless interest rate.

A stock which satisfies a stochastic differential equation below

$$dS_{c} = \sigma S_{c} d\hat{B}_{c}^{H} + \sigma S_{c} d\hat{B}_{c} + \mu S_{c} dt, \qquad S_{0} > 0, \qquad t \in [0, T], \tag{7}$$

where  $B_i^H$  is an fBm and  $B_i$  is a Bm with respect to  $\hat{\mathbb{P}}^H$ ,  $\mu$  is an appreciation rate, and  $\sigma$  is a volatility coefficient.

By using change of variable  $\sigma B_i + \sigma B_i^H = \mu - r + \sigma \hat{B}_i + \sigma \hat{B}_i^H$ , then under a risk-neutral measure, we have

$$dS_{t} = \sigma S_{t} dB_{t}^{H} + \sigma S_{t} dB_{t} + rS_{t} dt, \qquad S_{0} > 0, \qquad t \in [0, T].$$
(8)

Furthermore by using a Itô formula in [24], we obtain a solution of (8) as

$$S_{t} = S_{0} \exp\left(\sigma B_{t}^{H} + \sigma B_{t} - \frac{1}{2}\sigma^{2}t^{2H} - \frac{1}{2}\sigma^{2}t + rt\right). \tag{9}$$

The price of a European option at time t with an expire date T and a strike price K is denoted  $C(t,S_t)$ . We present a formula for a call option pricing under MFBM in the theorem below.

**Theorem 6.** Suppose a stock price  $S_t$ , defined by then the price at time  $t \in [0,T]$  of a European call option with an expire date T and a strike price K is given by

$$C(t, S_t) = S_t N(d_1) - Ke^{-r(T-t)} N(d_2),$$
(10)

where

$$d_{1} = \frac{\frac{1}{2}\sigma^{2}(T^{2H} - t^{2H}) + \frac{1}{2}\sigma^{2}(T - t) + \ln\left(\frac{S_{t}}{K}\right) + r(T - t)}{\sqrt{\sigma^{2}(T^{2H} - t^{2H}) + \sigma^{2}(T - t)}},$$
(11)

$$d_{2} = \frac{-\frac{1}{2}\sigma^{2}\left(T^{2H} - t^{2H}\right) - \frac{1}{2}\sigma^{2}\left(T - t\right) + \ln\left(\frac{S_{t}}{K}\right) + r(T - t)}{\sqrt{\sigma^{2}\left(T^{2H} - t^{2H}\right) + \sigma^{2}\left(T - t\right)}},$$
(12)

 $N(\cdot)$  is a cumulative probability function f a standard normal distribution.

Proof: Motivated from Theorem 5, the call option with an expire date T and a strike price K is theoretically equivalent to

$$C(t, S_t) = \widetilde{\mathbb{E}} \left[ e^{-r(T-t)} \max\{S_T - K, 0\} \middle| \mathcal{F}_t^H \right]$$

$$= e^{-r(T-t)} \widetilde{\mathbb{E}} \left[ S_T \mathbf{1}_{\{S_T > K\}} \middle| \mathcal{F}_t^H \right] - K e^{-r(T-t)} \widetilde{\mathbb{E}} \left[ \mathbf{1}_{\{S_T > K\}} \middle| \mathcal{F}_t^H \right].$$
(13)

Meanwhile option holders would exercise the option only when  $S_T > K$ . Solving (9) on this boundary, we have

$$\sigma B_T^H + \sigma B_T > \frac{1}{2}\sigma^2 T^{2H} + \frac{1}{2}\sigma^2 T + \ln\left(\frac{K}{S_0}\right) - rT.$$

Let

$$d_2^* = \frac{1}{2}\sigma^2 T^{2H} + \frac{1}{2}\sigma^2 T + \ln\left(\frac{K}{S_0}\right) - rT.$$
 (14)

Using Corollary 3 and applying (14) on the second of the RHS in (13), we have

$$\begin{split} \tilde{\mathbb{E}}\Big[\mathbf{1}_{\left\{S_{T}>K\right\}}\Big|\mathcal{F}_{t}^{H}\Big] &= \tilde{\mathbb{E}}\Big[\mathbf{1}_{\left\{x>d_{2}^{*}\right\}}\Big(\sigma B_{T}^{H} + \sigma B_{T}\Big)\Big|\mathcal{F}_{t}^{H}\Big] \\ &= \int_{d_{2}^{*}}^{\infty} \frac{\mathbf{8}}{\sqrt{2\pi\Big(\sigma^{2}\Big(T^{2H} - t^{2H}\Big) + \sigma^{2}\Big(T - t\Big)\Big)}} \exp\left(-\frac{\Big(y - \sigma B_{t}^{H} - \sigma B_{t}\Big)^{2}}{2\Big(\sigma^{2}\Big(T^{2H} - t^{2H}\Big) + \sigma^{2}\Big(T - t\Big)\Big)}\right) dy \end{split}$$

ICRIEMS 5 IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1097 (2018) 012081 doi:10.1088/1742-6596/1097/1/012081

$$\widetilde{\mathbb{E}}\left[1_{\{S_{T}>K\}} \middle| \mathcal{F}_{t}^{H}\right] = \int_{-\infty}^{\frac{\sigma B_{t}^{H} + \sigma B_{t} - d_{x}^{2}}{\sqrt{\left(\sigma^{2}(T^{2H} - t^{2H})\right) + \sigma^{2}(T - t)\right)}}} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^{2}}{2}\right) dz$$

$$= N(d_{2}), \tag{15}$$

where  $d_2 = \frac{\sigma B_T^H + \sigma B_T - d_2^*}{\sqrt{\left(\sigma^2 \left(T^{2H} - t^{2H}\right) + \sigma^2 \left(T - t\right)\right)}}$ . Furthermore, (9) can be written as

$$\sigma B_t^H + \sigma B_t = \frac{1}{2}\sigma^2 t^{2H} + \frac{1}{2}\sigma^2 t + \ln\left(\frac{S_t}{S_0}\right) - rt.$$
 (16)

Hence, using (14) and (16) on  $d_2$ , we have

$$d_{2} = \frac{-\frac{1}{2}\sigma^{2}(T^{2H} - t^{2H}) - \frac{1}{2}\sigma^{2}(T - t) + \ln\left(\frac{S_{t}}{K}\right) + r(T - t)}{\sqrt{\sigma^{2}(T^{2H} - t^{2H}) + \sigma^{2}(T - t)}}.$$

Let us consider a process

$$\sigma B_t^{H^*} + \sigma B_t^* = \sigma B_t^H - \sigma^2 t^{2H} + \sigma B_t - \sigma^2 t, \tag{17}$$

for  $t \in [0,T]$ . The fractional Girsanov theorem assures us that there is a probability measure  $\mathbb{P}^{H^*}$  such that  $\sigma B_t^{H^*} + \sigma B_t^*$  is an new mfBm under  $\mathbb{P}^{H^*}$ . We will denote

$$Z_{t} = \exp\left(\sigma B_{t}^{H} - \frac{1}{2}\sigma^{2}t^{2H} + \sigma B_{t} - \frac{1}{2}\sigma^{2}t\right)$$

$$\tag{18}$$

By using Theorem 4 and (18) on the first of the RHS in (13), we have

$$\widetilde{\mathbb{E}}\left[S_{T}1_{\{S_{T}>K\}}\middle|\mathcal{F}_{t}^{H}\right] = \widetilde{\mathbb{E}}\left[S_{0} e^{rT} e^{\left(\sigma B_{T}^{H} - \frac{1}{2}\sigma^{2}T^{2H} + \sigma B_{T} - \frac{1}{2}\sigma^{2}T\right)}1_{\{S_{T}>K\}}\middle|\mathcal{F}_{t}^{H}\right] \\
= S_{0} e^{rT}\widetilde{\mathbb{E}}\left[Z_{T}1_{\{S_{T}>K\}}\middle|\mathcal{F}_{t}^{H}\right] \\
= S_{0} e^{rT}\widetilde{\mathbb{E}}\left[Z_{T}1_{\{y>d_{2}^{*}\}}\left(\sigma B_{T}^{H} + \sigma B_{T}\right)\middle|\mathcal{F}_{t}^{H}\right] \\
= S_{0} e^{rT}Z_{t}\widetilde{\mathbb{E}}^{*}\left[1_{\{y>d_{2}^{*}\}}\left(\sigma B_{T}^{H} + \sigma B_{T}\right)\middle|\mathcal{F}_{t}^{H}\right] \\
= S_{0} e^{rT}Z_{t}\widetilde{\mathbb{E}}^{*}\left[1_{\{S_{T}>K\}}\middle|\mathcal{F}_{t}^{H}\right] \tag{19}$$

By substituting (17) into (9), we obtain

$$S_t = S_0 \exp(\sigma B_t^{H^*} + \sigma B_t^* + \frac{1}{2}\sigma^2 t^{2H} + \frac{1}{2}\sigma^2 t + rt).$$
 (20)

Solving (20) in time T for the boundary  $S_T > K$ , we have

$$\sigma B_T^{H^*} + \sigma B_T^* > -\frac{1}{2}\sigma^2 T^{2H} - \frac{1}{2}\sigma^2 T + \ln\left(\frac{K}{S_0}\right) - rT.$$

If we denote

$$d_1^* = -\frac{1}{2}\sigma^2 T^{2H} - \frac{1}{2}\sigma^2 T + \ln\left(\frac{K}{S_0}\right) - rT,$$
(21)

we get

$$\widetilde{\mathbb{E}}^* \left[ \mathbf{1}_{\left\{ S_T > K \right\}} \left| \mathcal{F}_t^H \right. \right] = \widetilde{\mathbb{E}}^* \left[ \mathbf{1}_{\left\{ y > d_1^* \right\}} \left( \sigma B_T^{H^*} + \sigma B_T^* \right) \middle| \mathcal{F}_t^H \right. \right]$$

$$\tilde{\mathbb{E}}^{*} \left[ 1_{\{S_{\tau} > K\}} \left| \mathcal{F}_{t}^{H} \right. \right] = \int_{d_{t}^{*}}^{\infty} \frac{1}{\sqrt{2\pi \left(\sigma^{2} \left(T^{2H} - t^{2H}\right) + \sigma^{2} \left(T - t\right)\right)}} \exp \left( -\frac{\left(y - \sigma B_{t}^{H*} - \sigma B_{t}^{*}\right)^{2}}{2\left(\sigma^{2} \left(T^{2H} - t^{2H}\right) + \sigma^{2} \left(T - t\right)\right)} \right) dy$$

$$= \int_{-\infty}^{\sigma B_t^{H^+} + \sigma B_t^x - d_1^x} \frac{1}{\sqrt{2\pi}} \exp\left(-\frac{z^2}{2}\right) dz$$

$$= N(d_1), \tag{22}$$

where  $d_1 = \frac{\sigma B_t^{H^*} + \sigma B_t^* - d_1^*}{\sqrt{\sigma^2 (T^{2H} - t^{2H}) + \sigma^2 (T - t)}}$ . Subsequently, (20) can be written as

$$\sigma B_{t}^{H^{*}} + \sigma B_{t}^{*} = -\frac{1}{2}\sigma^{2}t^{2H} - \frac{1}{2}\sigma^{2}t + \ln\left(\frac{S_{t}}{S_{0}}\right) - rt.$$
 (23)

Substituting (21) and (23) on  $d_1$ , we get

$$d_{1} = \frac{\frac{1}{2}\sigma^{2}\left(T^{2H} - t^{2H}\right) + \frac{1}{2}\sigma^{2}\left(T - t\right) + \ln\left(\frac{S_{t}}{K}\right) + r(T - t)}{\sqrt{\sigma^{2}\left(T^{2H} - t^{2H}\right) + \sigma^{2}\left(T - t\right)}}.$$


Substitution of (22) into (19) yields


$$\tilde{\mathbb{E}}\left[S_{T}1_{\{S_{T}>K\}} \middle| \mathcal{F}_{t}^{H}\right] = S_{0}e^{rT}Z_{t}N(d_{1})$$

$$= S_{0}e^{rT}\exp\left(\sigma B_{t}^{H} + \sigma B_{t} - \frac{1}{2}\sigma^{2}t^{2H} - \frac{1}{2}\sigma^{2}t\right)N(d_{1})$$

$$= e^{r(T-t)}S_{t}N(d_{1}).$$
Finally, from (13), (15) and (24) we obtain
$$C(t,S_{t}) = e^{-r(T-t)}e^{r(T-t)}S_{t}N(d_{1}) - Ke^{-r(T-t)}N(d_{2})$$
(24)

$$C(t,S_t) = e^{-r(T-t)}e^{r(T-t)}S_t N(d_1) - Ke^{-r(T-t)} N(d_2)$$
  
=  $S_t N(d_1) - Ke^{-r(T-t)} N(d_2)$ .










FIGURE 1. Price of option on Hurst vs strike price at T=0.25.


FIGURE 2. Price of option on Hurst vs strike price at T=1.

FIGURE 3. Price of option on Hurst vs strike price at T=5.

The forgula in Theorem 6 allows us to determine a fair price for a European call option in terms of an expire date T, a strike price K, an initial stock price  $S_0$ , a risk-free interest rest r, and a stock volatility  $\sigma$ . Let S = 100,  $K \in (0,200)$ ,  $H \in (0,1)$ , r = 0.05, and  $\sigma = 0.3$ . When T = 0.25, T = 1, and T = 5, we get Figure 1, 2 and 3 respectively. We see that when  $K \to 200$  and  $H \to 0$  the price decreases significantly in Figure 3.







**FIGURE 4.** Price of option on Hurst vs volatility at T=0.25.

**FIGURE 5.** Price of option on Hurst vs volatility at T=1.

**FIGURE 6.** Price of option on Hurst vs volatility at *T*=5.

Let S = 100; K = 100; r = 0.05,  $\sigma \in (0, 1)$  and  $H \in (0, 1)$ . If T = 0.25 we obtain Figure 4 which is concave upward. The price increases significantly when  $\sigma \to 1$  and  $H \to 0$ . If T = 1 we obtain Figure 5 which is more linear and as the volatility increases for all Hurst parameters the price increases. If T = 5 we obtain Figure 6 which is concave down and we see that the prices increase significantly with high Hurst index and high volatility. Overall, as T,  $\sigma$  and H increase, the price increase in rate and magnitude.

#### 5. Conclusion

In this paper, to exclude arbitrage opportunities in an fBm model and to capture long-range dependence, stock returns are modeled with an mfBm. By using Fourier transformation method and quasi-conditional expectation theory, we get a formula for calculating a price of European call options. This formula can be used by investors to predict option prices for stocks that have long-range dependence.

#### Acknowledg 77 nts

The authors are gra 441 to the Directorate of Research and Community Service, the Republic of Indonesia, due to the funding of Doctoral Dissertation Research Grant 2018 funds with contract number 109 / SP2H / LT / DRPM / 2018 for this work.

#### References

- [1] Mills T C 1993 Is there long-term memory in UK stock returns? Appl. Financ. Econ. 3 303-6
- [2] Cheung Y-W and Lai K S 1995 A search for long memory in international stock market returns J. Int. Money Financ. 14 597–615
- [3] Necula C and Radu A-N 2012 Long memory in Eastern European financial markets returns Econ. Res. - Ekon. Istraživanja 25 316–77
- [4] Goddard J and Onali E 2012 Short and long memory in stock returns data *Econ. Lett.* 117 253–5
- [5] Cajueiro D O and Tabak B M 2008 Testing for long-range dependence in world stock markets Chaos, Solitons and Fractals 37 918–27
- [6] Gyamfi E N, Kyei K and Gill R 2016 Long-memory persistence in African Stock Markets EuroEconomica 35 83–91
- [7] Mandelbrot B B and Van Ness J W 1968 Fractional Brownian motions, fractional noises and applications SIAM Rev. 10 422–37
- [8] Hurst H E 1956 Methods of using long-term storage in reservoirs. Proc. Inst. Civ. Eng. 5 519–43
- [9] Molz F J, Liu H H and Szulga J 1997 Fractional Brownian motion and fractional Gaussian noise in subsurface hydrology: A review, presentation of fundamental properties, and extensions Water Resour. Res. 33 2273–86
- [10] Zhang H Y, Bai L H and Zhou A M 2009 Insurance control for classical risk model with fractional Brownian motion perturbation Stat. Probab. Lett. 79 473–80

- [11] Shokrollahi F and Kılıçman A 2015 Actuarial approach in a mixed fractional Brownian motion with jumps environment for pricing currency option Adv. Differ. Equations 2015 257
- [12] Rostek S 2009 Option Pricing in Fractional Brownian Markets (Springer-Verlag Berlin)
- [13] Xiao W-L, Zhang W-G, Zhang X X and Zhang X X 2012 Pricing model for equity warrants in a mixed fractional Brownian environment and its algorithm *Phys. A Stat. Mech. its Appl.* 391 6418–31
- [14] Sun L 2013 Pricing currency options in the mixed fractional Brownian motion Phys. A Stat. Mech. its Appl. 392 3441–58
- [15] Duncan T E, Hu Y and Pasik-Duncan B 2000 Stochastic calculus for fractional Brownian motion I. Theory SIAM J. Control Optim. 38 582–612
- [16] Hu Y and Øksendal B 2003 Fractional white noise calculus and applications to finance Infin. Dimens. Anal. Quantum Probab. Relat. Top. 06 1–32
- [17] Elliott R J and Van der Hoek J 2003 A general fractional white noise theory and applications to finance Math. Financ. 13 301–30
- [18] Necula C 2002 Option pricing in a fractional Brownian motion environment Adv. Econ. Financ. Res. - DOFIN Work. Pap. Ser. 1–18
- [19] Bender C and Elliott R J 2004 Arbitrage in a discrete version of the Wick-fractional Black-Scholes market Math. Oper. Res. 29 935–45
- [20] Björk T and Hult H 2005 A note on Wick products and the fractional Black-Scholes model Financ. Stochastics 9 197–209
- [21] Cheridito P 2003 Arbitrage in fractional Brownian motion models Financ. Stochastics 7 533–53
- [22] Bender C, Sottinen T and Valkeila E 2007 Arbitrage with fractional Brownian motion? Theory Stoch. Process. 13 23–34
- [23] Cheridito P 2001 Mixed fractional Brownian motion Bernoulli 7 913-34
- [24] Biagini F, Hu Y, Øksendal B and Zhang T 2008 Stochastic Calculus for Fractional Brownian Motion and Applications (Springer)
- [25] Zili M 2006 On the mixed fractional Brownian motion *J. AppliedMathematics Stoch. Anal.* **2006** 1–9

## European option pricing by using a mixed fractional Brownian motion

#### **ORIGINALITY REPORT**

23% SIMILARITY INDEX

16%
INTERNET SOURCES

19%
PUBLICATIONS

9%

STUDENT PAPERS

#### **PRIMARY SOURCES**

Submitted to University of South Florida
Student Paper

2%

Submitted to The Technical University of Kenya

1 %

Student Paper

Wensheng Wang. "Asymptotic analysis for hedging errors in models with respect to geometric fractional Brownian motion", Stochastics, 2018

1 %

Publication

biopen.bi.no
Internet Source

1%

Vasile Dragan, Toader Morozan, Adrian-Mihail Stoica. "Mathematical Methods in Robust Control of Discrete-Time Linear Stochastic Systems", Springer Science and Business Media LLC, 2010

%

Publication

6

www.aimspress.com

Internet Source

1 %

Tollege Student Paper  8 Xili Zhang, Weilin Xiao. "Arbitrage with fractional Gaussian processes", Physica A: Statistical Mechanics and its Applications, 2017 Publication  9 Www.tandfonline.com Internet Source  10 Satya N Majumdar, Gleb Oshanin. "Spectral content of fractional Brownian motion with stochastic reset", Journal of Physics A: Mathematical and Theoretical, 2018 Publication  11 Submitted to University of Hong Kong Student Paper  12 bphm.knu.ua Internet Source  13 Submitted to Peking University Shenzhen Graduate School Student Paper  14 pdfs.semanticscholar.org 15 Thomas L. Toulias, Christos P. Kitsos. "On the Generalized Lognormal Distribution", Journal			
fractional Gaussian processes", Physica A: Statistical Mechanics and its Applications, 2017 Publication  www.tandfonline.com Internet Source  1%  Satya N Majumdar, Gleb Oshanin. "Spectral content of fractional Brownian motion with stochastic reset", Journal of Physics A: Mathematical and Theoretical, 2018 Publication  Submitted to University of Hong Kong Student Paper  Submitted to Peking University Shenzhen Graduate School Student Paper  Mathematics of Hong Kong Student Paper  1%  Thomas L. Toulias, Christos P. Kitsos. "On the	7	College	1 %
Satya N Majumdar, Gleb Oshanin. "Spectral content of fractional Brownian motion with stochastic reset", Journal of Physics A: Mathematical and Theoretical, 2018 Publication  Submitted to University of Hong Kong 1%  bphm.knu.ua Internet Source 1 %  Submitted to Peking University Shenzhen Graduate School Student Paper 1 %  pdfs.semanticscholar.org 1 %  Thomas L. Toulias, Christos P. Kitsos. "On the 1 %	8	fractional Gaussian processes", Physica A: Statistical Mechanics and its Applications, 2017	1 %
content of fractional Brownian motion with stochastic reset", Journal of Physics A: Mathematical and Theoretical, 2018 Publication  11 Submitted to University of Hong Kong Student Paper  12 bphm.knu.ua Internet Source  1 %  13 Submitted to Peking University Shenzhen Graduate School Student Paper  14 pdfs.semanticscholar.org Internet Source  1 %  15 Thomas L. Toulias, Christos P. Kitsos. "On the	9		1 %
bphm.knu.ua Internet Source  Submitted to Peking University Shenzhen Graduate School Student Paper  1 %  pdfs.semanticscholar.org Internet Source  1 %  Thomas L. Toulias, Christos P. Kitsos. "On the	10	content of fractional Brownian motion with stochastic reset", Journal of Physics A: Mathematical and Theoretical, 2018	1 %
Submitted to Peking University Shenzhen Graduate School Student Paper  1 96  1 96  1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	11		1 %
Graduate School Student Paper  14 pdfs.semanticscholar.org Internet Source  1 %  Thomas L. Toulias, Christos P. Kitsos. "On the	12		1 %
15 Thomas L. Toulias, Christos P. Kitsos. "On the	13	Graduate School	1 %
0%	14		1 %
	15		1 %


# of Probability and Statistics, 2013 Publication

16	Submitted to University of Bradford  Student Paper	1 %
17	Submitted to University of Warwick Student Paper	1 %
18	www.fma.org Internet Source	1 %
19	Qinyu Liu, Ting Jin, Min Zhu, Chenlei Tian, Fuzhen Li, Depeng Jiang. "Uncertain Currency Option Pricing Based on the Fractional Differential Equation in the Caputo Sense", Fractal and Fractional, 2022	<1%
20	cpb.iphy.ac.cn Internet Source	<1%
21	Submitted to University of Surrey  Student Paper	<1%
22	theory.ic.ac.uk Internet Source	<1%
23	Cheridito, P "Gaussian moving averages, semimartingales and option pricing", Stochastic Processes and their Applications, 200401 Publication	<1%

24	Panhong Cheng, Zhihong Xu. "Pricing Vulnerable Options in a Mixed Fractional Brownian Motion with Jumps", Discrete Dynamics in Nature and Society, 2021 Publication	<1%
25	binus.ac.id Internet Source	<1%
26	Robert J Elliott, Leunglung Chan. "Perpetual American options with fractional Brownian motion", Quantitative Finance, 2004 Publication	<1%
27	fmipa.unsrat.ac.id Internet Source	<1%
28	www.hrpub.org Internet Source	<1%
29	Karim M. Abadir, Michael Rockinger. "DENSITY FUNCTIONALS, WITH AN OPTION-PRICING APPLICATION", Econometric Theory, 2003 Publication	<1%
30	Li Meng. "Comparison of Black–Scholes Formula with Fractional Black–Scholes Formula in the Foreign Exchange Option Market with Changing Volatility", Asia-Pacific Financial Markets, 10/27/2009	<1%
	oprints soton as uk	4

eprints.soton.ac.uk
Internet Source

32	personal.lut.fi Internet Source	<1%
33	pubs.sciepub.com Internet Source	<1%
34	s3.amazonaws.com Internet Source	<1%
35	www.degruyter.com Internet Source	<1%
36	www.semanticscholar.org Internet Source	<1%
37	Foad Shokrollahi. "The evaluation of geometric Asian power options under time changed mixed fractional Brownian motion", Journal of Computational and Applied Mathematics, 2018 Publication	<1%
38	W. Grecksch, C. Roth, V. V. Anh. "-Fractional Brownian Motion in Infinite Dimensions with Application to Fractional Black–Scholes Market ", Stochastic Analysis and Applications, 2009 Publication	<1%
39	virgo.unive.it Internet Source	<1%
40	www.fm.mathematik.uni-muenchen.de Internet Source	<1%





## Parravano, Antonio, Néstor Sánchez, and Emilio J. Alfaro. "THE DEPENDENCE OF PRESTELLAR CORE MASS DISTRIBUTIONS ON THE STRUCTURE OF THE PARENTAL CLOUD", The Astrophysical Journal, 2012.

<1%

Publication

Exclude quotes On Exclude bibliography On

Exclude matches

< 5 words