# ISCAIE 2018 2018 IEEE Symposium on Computer Applications & Industrial Electronics

28<sup>th</sup> - 29<sup>th</sup> April 2018, Penang Island, MALAYSIA

Organizer:

IEEE Malaysia Industrial Electronics / Industrial Applications Joint Chapter Secretariat:



Centre of Computer Engineering Studies, Universiti Teknologi Mara, Malaysia

### FOREWORD

I am happy to welcome and wish "SELAMAT DATANG" to all delegates attending the 2018 IEEE Symposium on Computer Applications & Industrial Electronics (ISCAIE 2018). We have been organizing ISCAIE for six years now, and we continue our tradition of maintaining its high standard and quality. ISCAIE2018 has received full support and technical assistance from the IEEE Malaysia Industrial Electronics & Industrial Applications Joint Chapter.

We uphold the established four quality assurance features, inclusive of employing web-based paper submission, plagiarism check, double blind review and submission of the ISCAIE2018 proceedings to the IEEE Conference Publications Program (which is accessible via IEEE Xplore worldwide and selected papers will be indexed by SCOPUS). These procedures ensure that ISCAIE2018 proceedings will be of excellent quality, at par with that of previous ISCAIEs, which have been indexed in IEEE Xplore and SCOPUS.

ISCAIE2018 is a conference focusing on a broad domain of industrial electronics and computer applications, enabling meeting opportunities for participants to discuss and share their research, experiences and future endeavors. These fields of interest are highly relevant and together with the progress in Engineering, Science and Technology. We believe that apart from the scheduled presentations, the interaction among the participants will be valuable, enriching research networking as well as promoting the engineering, science and technological developments in this part of the world.

I would like to encourage our foreign guests, to take the opportunity sometime after the conference, to venture around and see our wonderful Penang Island with its white sandy beaches, heritage sites and historical marvels.

I would like to express much gratitude to all ISCAIE2018 organizing committee members for their strong commitment and support. My appreciations also to all reviewers of ISCAIE2018 for assisting us in attaining a high standard for papers submitted to the conference proceeding.

To the IEEE Malaysia Industrial Electronics & Industrial Applications Joint Chapter, I would like to express my special thanks for sponsoring the conference. Many thanks and appreciations also to the Faculty of Electrical Engineering, Universiti Teknologi MARA (UiTM) and all other organizations that have greatly contributed to the success of ISCAIE2018.

Finally, to all the participants of the conference, please have a fruitful and enjoyable time, and thank you very much.

Nooritawati Md Tahir General Chair ISCAIE 2018

# **TABLE OF CONTENTS**

FOREWORD	i
TABLE OF CON	TENTSii
ORGANIZING (	COMMITTEEiii
REVIEWERS	iv
TECHNICAL PR	OGRAM OVERVIEWxii
TECHNICAL PR	OGRAM / ABSTRACTxiii
a.	Technical Session B1
b.	Technical Session A1
с.	Technical Session B2
d.	Technical Session A2
e.	Technical Session B3
f.	Technical Session A3
g.	Technical Session B4
h.	Technical Session A4
i.	Technical Session B5
j.	Technical Session A5
k.	Technical Session B6
1.	Technical Session A6
AUTHOR INDEX	Xxxxvi
MANUSCRIPTS	

## **ORGANIZING COMMITTEE**

**General Chair** Nooritawati Md Tahir

**Secretary** Rozita Jailani

**Finance Chair** Rozita Jailani

**Publication Chair** Syed Farid Syed Adnan

**Technical Chair I** Habibah Hashim

Technical Chair II Ihsan Mohd Yassin

Website & IT Systems Chair Ihsan Mohd Yassin

**Publicity Chair & Admin Representative** Roslina Mohamad

**Local Arrangement Chair** Nooritawati Md Tahir

### **TECHNICAL PROGRAM OVERVIEW**

2018 IEEE Symposium on Computer Applications & Industrial Electronics (ISCAIE2018					
2018-04-27					
17:30 - 19:00	: Registration @ Kayu Manis Room				
2018-04-28					
09:00 - 13:00	: Registration @ Kayu Manis Room				
	Technical Session A1 @ Pandan Room				
	Technical Session B1 @ Halia Room				
10:30 - 10:45	: Morning Break @ Foyer				
10:45 - 12:30	: Technical Session A2 @ Pandan Room				
	Technical Session B2 @ Halia Room				
12:30 - 14:00	: Lunch @ Cinnamon Asian Kitchen				
14:00 - 16:00	: Technical Session A3 @ Pandan Room				
	Technical Session B3 @ Halia Room				
16:00 - 16:15	: Afternoon Break @ Foyer				
16:15 - 17:45	: Technical Session A4 @ Pandan Room				
	Technical Session B4 @ Halia Room				
2018-04-29					
09:00 - 10:30	: Technical Session A5 @ Pandan Room				
	Technical Session B5 @ Halia Room				
10:30 - 10:45	: Morning Break @ Foyer				
10:45 - 12:45	: Technical Session A6 @ Pandan Room				
	Technical Session B6 @ Halia Room				
12:45 - 13:15	: Closing Ceremony, Best Paper Award, Photography Session @ Pandan Room				
13:15 - 14:00	: Lunch @ Cinnamon Asian Kitchen				

# A Method of Mood Classification on Keroncong Music

Iwan Binanto Informatics Department, Faculty of Science and Technology Sanata Dharma University Yogyakarta, Indonesia iwan@usd.ac.id

*Abstract*— Some researchers in Indonesia use keroncong music for depression reduction therapy in elderly and others. It is believed that keroncong music affects someone's mood. This research will classify mood of keroncong music in Thayer model, computationally, by median. MIRToolbox in Matlab used to feature extraction of keroncong music. Median values used to classify mood of keroncong music in Thayer model. The results of this research are expected to support other researchers related to the success of music therapy to reduce levels of depression in elderly.

Keywords—mood classification; mood music; median; music retrieval; mirtoolbox; thayer model mood; keroncong music

#### I. INTRODUCTION

Music therapy can effectively overcome various depressive disorders including severe depression, as well as depression in elderly even with neurological disorders (stroke, dementia) [1]–[3]. Music therapy has been widely used to heal depression [4] with a common method is listening to music [3]. This widespread use is due to the excellence of safe and manageable music therapy [5] [6]. Various types of music that are considered qualified to be used for therapy [3], [4], [7], while [8] in his research uses a kind of music that is considered soothing without detailing the type of music. According to [9], acceptance of music is affectively determined by the cultural tradition rather than the quality of the music itself. This finding gives an idea to [6] to do a research on music therapy to reduce depression and stress levels by using local music, namely Raga's, classical music from India. Similar research also has been done with different research object [10]. Previous literature focused on music therapy to reduce symptoms of mild depression and a seasonal mood disorder in the elderly with Chinese local music [1].

In Indonesia there are several kinds of research on music therapy to reduce depression score with local music, which are utilizing keroncong music [11]–[13] and traditional Javanese music [14], [15].

Mood perception and its relation to local music were investigated by [16]. The method they used is collecting the opinion of the respondents, the listeners who have been selected. This is very subjective because the results depend on the culture, education, personality, and personal experience of the respondents [17], [18]. Therefore, an objective method for classification is required in mood perception. Objective classification, in this case, can only be done computationally without involving human feelings. Related research has been conducted [17]–[22]. Some researchers are based on the classification of mood perceptions using Thayer model [17], [18], [21], [22]. It is because the Thayer model already covers all the basic moods that respond to music [17].

So far, there is no research on classifying mood perception on local Indonesian music, especially keroncong music. Keroncong music is chosen because it is Indonesian original music and also used for music therapy, especially for depression therapy as done by [12], [13], [23]. This paper will discuss classifying mood perception Thayer model based on Keroncong Music, computationally.

#### II. RELATED WORKS

The relationship between music and mood has been studied for a long time. [24] began his research in 1935 and a year later [25] found 8 (eight) groups consist of 66 adjectives describing the mood of the music and creating the cycle. These eight groups of adjectives were corrected and grouped into nine groups of 50 adjectives [26]. The adjectives used to describe the mood are very free and numerous. Fortunately, as explained by [17], there is a study of the fundamental emotional dimensions done by [27] coupled with research conducted by [28] which provides the base of mood taxonomy as well as some important computational modeling cues for the classification of mood perceptions. However, there is no acceptable standard of mood taxonomy [17].

The identification of the mood of the music is important for human especially for music search and organizing [29]. This is because humans want to listen to music in accordance with the mood at the time [30]. This can underlie the music used for therapy.

Music therapy successfully reduced levels of depression associated with mood [4]. There is a mood change in elderly while listening to Javanese Langgam music that has a slow and regular tempo (about 60 beats per minute). Therefore, based on a statistical evaluation, it reduced the level of anxiety in the elderly [14], [15]. Similarly, [3] and [7] said that listening to music could affect and improve the mood that [3] asserts as a healing process. It is because tempo, loudness (sound level), spectrum (timbre), and articulation affect the expression of one's emotions [19].

Research related to the influence of music including local music to the psychological condition of the individual has been widely practiced. The results of these researchers are summarized by [13] as the emergence of physical and mental reactions in response to music. These reactions can be a calm, relaxation or a change in the rhythm of breathing, blood pressure and blood flow to the heart and hormone production which lead to better mood changes. These reactions are also presented by [11], [14], [15].

Another research was conducted by [20] who successfully automatically extracted the various expressions of emotion inherent in a work of music embodied in the form of human facial expressions in a 3D model. Previously, [17] stated that it is possible to establish a mood perception classification system in certain contexts, in this case: classical music in western culture. He provided a method to automatically classify mood perceptions from music data based on Thayer model. Similar research with different contexts has been done by [18] who provided a method for automatically classifying mood perception on Indian Bollywood music based on Thayer model. This model has also been used by [21] as a model to track mood changes in a musical composition, especially western classical music. [22] also used the Thayer model to perceive the mood of western music and Indian music.

There are many researchers on music related to human mood, but those which use Indonesian music such as keroncong music haven't been found. There has been a lot of researchers on keroncong music for therapy, one of them was conducted by [31] who found that keroncong music positively influences performances of data input operators because it can reduce stress and monotony of work. While other researchers more on elderly as done by [12] and [13] who found that keroncong music reduced the level of depression in the elderly. In addition, keroncong music can also reduce the incidence of insomnia in the elderly as proposed by [32]. Decreasing levels of anxiety in the elderly were also successfully performed with keroncong music therapy by [23].

#### III. MUSIC AND MOOD

Characteristics of music such as rhythm/tempo, melody, harmony, pitch, and timbre have an important role in the physiological and psychological functions of humans, so as to create mood changes [33]. They can be used to communicate the mood of the music to the listener. Several kinds of research have been done, especially the computing mood classification using acoustic characteristics [19].

At the psychological level, music can involve multiple social functions, improve communication and social cohesion, and can promote empathetic relationships, especially in an active music therapy approach. While from a rehabilitative point of view, music can affect motor function and regulation. This effect is associated with pleasure so that positively affects the mood and causes the healing process to occur [3]. Thayer model is used as the base for the classification of the human mood to music or to create a mood classification of music to facilitate the search for music and organizing. This is because the four clusters already cover all the basic mood that responds to music [17].

Fig.1 shows the Thayer model rewritten by [17] and drawn by [22]. There are two basic stimuli affect the mood response, namely Stress, and Energy. From left to right represent pleasant to unpleasant, and from bottom to top show Calm to Energetic. Four quadrants represent happy-energetic (Exuberance), happy-quiet (Contentment), tense-energetic (Anxious / Frantic), and tense-quiet (Depression). It should be noted that the names of each dimension and each cluster are not the same (varies) in different literature. For example, the Energy dimension is also referred to as Arousal or Activity Level, while the Stress dimension is also called Pleasure or Valence. In Fig. 1, all names are attempted to be written.



Fig. 1. Thayer model for classification mood [22]

For classification music mood computationally, [22] extracted 4 (four) features based on the mood that represents the basic moods, which are intensity, timbre, pitch, and rhythm. While [17] and [18] extracted 3 (three) features only, which are intensity, timbre, and rhythm. They are very important because affect the particular mood.

The mood grouped into two major groups, namely Group 1: Contentment and Depression and Group 2: Exuberance and Anxious / Frantic [17]. For mood classification of a music, [17] provided the basic rules, if the intensity is low, then the music will be grouped into Group 1 (Contentment and Depression). If the intensity is HIGH, then the music is grouped into Group 2 (Exuberance and Anxious / Frantic). While [18] provided other which are if music has a rhythm with LOW/SLOW tempo, it will be grouped into Group 1. If music has a rhythm with a FAST/HIGH tempo, it will be grouped into Group 2. Combining basic rules from [17] and [18] will be used in this research.

#### IV. METHOD

86 songs of Keroncong music are collected from various sources, ranging from CD to downloaded music files from youtube. This collection then cleaned from the duplicate, so only get 1 version of the song per title, because, at the time of collecting song, there are some songs with same titles with different versions, especially at tempo. Therefore, must choose one of the songs only. The selection method is random, not based on the version of the songs.

Finally obtained 78 songs of keroncong music which then extracted their features: intensity and tempo. Intensity and tempo are gained by computing RMS (Root-Mean-Square) as well as did by [22] and [18]. This work done by MIRToolbox in MATLAB [34].

The data of each feature extraction are saved to Microsoft Excel that will be classified. Method to classified the data is by Median as [35] did, which were divided the data into two sub-data. The LOW if it is less than the Median value, and the HIGH if it is the opposite. This is to determine the group mood of a song, whether grouped to Group 1 (Contentment and Depression) or Group 2 (Exuberance and Anxious / Frantic). This illustrated in Fig. 2.



Fig. 2. The Median value for grouping mood

With this classification, there are songs that are contradictory, which have LOW Intensity, but HIGH tempo, whereas the song must meet LOW on both Intensity and Tempo as well as for HIGH. This is to accommodate the basic rules given by [17] about the intensity and [18] about the tempo, which must have the same category for both Intensity and Tempo, to be able to group in one of the mood groups.

To do this, author developed an algorithm that implemented in Microsoft Excel's Visual Basic Macro. The algorithm will match one by one song title which had the same intensity and tempo. If the intensity and tempo both were the same, then the song title will be saved. The song title then stored in a specific column in the worksheet. With this algorithm, many songs are not grouped, either in Group 1 nor Group 2.

Once the songs grouped into the appropriate group, then calculating Median in each group. This will put the songs into their according to mood. Fig. 3 and 4 show the median as the limit value of LOW or HIGH from the intensity and tempo, which will be classified into Group 1 (Contenment & Depression) or Group 2 (Exuerance & Anxious/Frantic).

#### V. RESULTS AND DISCUSSIONS

That algorithm generate 22 songs grouped to Group 1, which have both LOW Intensity and Tempo, while 23 songs grouped to Group 2, which have both HIGH Intensity and Tempo.

From this point, each group be calculated its Median again to find LOW and HIGH groups, which would be categorized by definite mood. Fig. 5., Fig. 6., Fig. 7., Fig. 8. show this respectively. This categorization is done with the same algorithm as before.

From Fig. 2, it seen that in Group 1, the LOW both Intensity and Tempo will placed into the Contentment mood, otherwise, it will placed into the mood of Depression. In Group 2, the LOW both Intensity and Tempo will placed into the Exuberance mood, otherwise, it will placed to the Anxious / Frantic mood.

In the final result, only 23 songs title are classified properly, while the rest cannot fit into either of 4 categories of Thayer mood models. There are 5 songs in Group 1 belong to Contentment and 5 songs belong to Depression as in Table I. While 6 songs in Group 2 belong to Exuberance and 7 songs belong to Anxious/Frantic as in Table II. This is due to the basic rule the intensity [17] and [18] about the tempo as stated above.

TABLE I. MOOD GROUP 1

CONTENTMENT		DEPRESSION	
No.	Song Title	No.	Song Title
1	Sampul Surat	1	Kicir-kicir
2	Rujak Uleg	2	Nyiur Hijau
3	Kau Selalu Dihatiku	3	Kalung Mutiara
4	Lambaian sang merah putih	4	Senyuman Chandra
5	Dunia Berdamai	5	Wuyung

TABLE II. MOOD GROUP 2

EXUBERANCE		ANXIOUS/FRANTIC	
No.	Song Title	No.	Song Title
1	Jenang Gulo	1	Hasrat Menyala
2	Sebelum Aku Mati	2	Dinda bestari
3	Gethuk	3	Jembatan Merah
4	Juwita Malam	4	Solo Diwaktu Malam
5	Rindu Malam	5	Mahameru
6	Kota Solo	6	Bengawan Solo
		7	Sejak Kita Berpisah



Fig. 3. The Median of Intensity



Fig. 4. The Median of Tempo



Fig. 5. The Median of Intensity in Group 1



Fig. 6. The Median of Tempo in Group 1



Fig. 7. The Median of Intensity in Group 2



Fig. 8. The Median of Tempo in Group 2

#### VI. CONCLUSIONS

This research is an early stage to be able to classify keroncong music completely and perfectly so that it can be used as a reference for the use of keroncong music as depression therapy for elderly. Nevertheless, this research objectively produced a list of keroncong music songs that belong to the four mood classifications in the Thayer model because there is no human feelings intervention.

Although this research is very early, it can be concluded that not all keroncong music songs can be classified and included in one category of Thayer mood models. Therefore, not all keroncong music songs can be used for depression therapy in elderly. However, these conclusions and opinions still have to be verified. It need further research, especially in the relationship between mood perceptions of keroncong music with depression therapy in elderly. It should include keroncong music experts and/or psychologists, especially in the field of depression.

#### ACKNOWLEDGMENT

Thanks to LPPM Sanata Dharma University which has funded this research in the grant year 2017 and also to Prof. Riyanarto Sarno from Institut Teknologi Sepuluh Nopember Surabaya for the idea.

#### REFERENCES

- X. Liu, X. Niu, Q. Feng, and Y. Liu, "Effects of five-element music therapy on elderly people with seasonal affective disorder in a Chinese nursing home," *J. Tradit. Chinese Med.*, vol. 34, no. 2, pp. 159–161, 2014.
- [2] S. Nizamie and S. Tikka, "Psychiatry and music," *Indian J. Psychiatry*, vol. 56, no. 2, pp. 128–140, 2014.
- [3] A. Raglio, "Effects of music and music therapy on mood in neurological patients," *World J. Psychiatry*, vol. 5, no. 1, p. 68, 2015.
- [4] A. Maratos, C. Gold, X. Wang, and M. Crawford, "Music therapy for depression," *Cochrane Database Syst. Rev.*, vol. 1, no. 1, pp. 1–20, 2009.
- [5] K. V. Jespersen, J. Koenig, P. Jennum, and P. Vuust, "Music for insomnia in adults," *Cochrane database Syst. Rev.*, vol. 8, no. November, 2015.
- [6] S. Sumathy, D. Pavitra, and P. Parin N., "Indian Classical Music As Receptive Music Therapy Improves Tridoshic Balance and Major Depression in a Pregnant Woman," *Int. J. Ayurveda Pharma Res.*, vol. 4, no. 9, pp. 8–11, 2016.
- [7] D. Evans, "The effectiveness of music as an intervention for hospital patients: A systematic review," J. Adv. Nurs., vol. 37, no. 1, pp. 8–18, 2002.
- [8] R. Mottaghi, A. Kamkar, and A. Maredpoor, "Effectiveness of Targeted Musical Therapy on Sleep Quality and Overcoming Insomnia in Seniors," *Iran. J. Ageing*, vol. 11, no. 2, pp. 348–357, 2016.
- [9] A. H. Gregory and N. Varney, "Cross-cultural comparisons in the affective response to music," *Psychol. Music*, vol. 24, no. 1, pp. 47–52, 1996.
- [10] K. K. Alphonsa and C. Bhattacharya, "Music Effect on Distressed Relatives Waiting Outside ICUs," *Int. J. Heal. Scences Res.*, vol. 6, no. 4, pp. 298–305, 2016.
- [11] S. E. Purbowinoto and Kartinah, "Pengaruh Terapi Musik Terhadap Perubahan Tingkat Depresi Pada Lansia Di PSTW (Panti Sosial Tresna Wredha) Unit Budi Luhur, Kasongan, Bantul Yogyakarta," *Ber. Ilmu Keperawatan*, vol. 4, no. 1, pp. 44–49, 2011.
- [12] F. Firdawati and S. Riyadi, "Hubungan Terapi Musik Keroncong Dengan Tingkat Depresi Pada Lansia Di Panti Wredha Budhi Dharma

Yogyakarta 2014," J. Kesehat. Samodra Ilmu, vol. 5, no. 2, pp. 155–162, 2014.

- [13] L. A. Soeraya and S. Sarifah, "PENGARUH TERAPI MUSIK KERONCONG TERHADAP TINGKAT DEPRESI PADA LANSIA," *J. Profesi*, vol. 13, no. 1, pp. 52–55, 2015.
- [14] R. Lestari, Setyohadi, Kumboyono, and R. Purwo S., "Pengaruh Irama Musik Tradisional Jawa Terhadap Penurunan Skor Depresi pada Lanjut Usia," *J. Ilmu Keperawatan UGM*, vol. 4, no. 3, pp. 141–145, 2009.
- [15] Z. Noor and Junaidi, "Penurunan Tingkat Kecemasan Pada Lansia Melalui Terapi Musik Langgam Jawa," J. Keperawatan Indones., vol. 13, no. 3, pp. 195–201, 2010.
- [16] L. L. Balkwill, W. F. Thompson, and R. Matsunaga, "Recognition of emotion in Japanese, Western, and Hindustani music by Japanese listeners," *Jpn. Psychol. Res.*, vol. 46, no. 4, pp. 337–349, 2004.
- [17] L. Lu, D. Liu, and H. Zhang, "Automatic Mood Detection and Tracking of Music Audio Signals," *IEEE Trans. Audio. Speech. Lang. Processing*, vol. 14, no. 1, pp. 5–18, 2006.
- [18] V. Hampiholi, "A method for Music Classification based on Perceived Mood Detection for Indian Bollywood Music," *Int. J. Comput. Electr. Autom. Control Inf. Eng.*, vol. 6, no. 12, pp. 1636–1643, 2012.
- [19] P. N. Juslin, "Cue utilization in communication of emotion in music performance: Relating performance to perception.," J. Exp. Psychol. Hum. Percept. Perform., vol. 26, no. 6, pp. 1797–1813, 2000.
- [20] M. Mancini, R. Bresin, and C. Pelachaud, "A virtual head driven by music expressivity," *IEEE Trans. Audio, Speech Lang. Process.*, vol. 15, no. 6, pp. 1833–1841, 2007.
- [21] J. Grekow, "Mood Tracking of Musical Compositions," in 20th International Conference on Foundations of Intelligent Systems, 2012, pp. 228 – 233.
- [22] A. S. Bhat, "An Efficient Classification Algorithm For Music Mood Detection In Western and Hindi Music Using Audio Feature Extraction," in 2014 Fifth Int rna ona Conference on Signa s and Im ge Proc ssing 2014 Fifth Inteernattiionall Conferencee on Signall and Imaage Proceessing An, 2014, pp. 359–364.

- [23] S. Maryati and Suyami, "Dampak terapi musik keroncong terhadap tingkat kecemasan lansia di desa pasung wedi klaten," *J. Mot.*, vol. 10, no. 21, pp. 44–54, 2015.
- [24] K. Hevner, "The Affective Character of the Major and Minor Modes in Music," vol. 47, no. 1, pp. 103–118, 1935.
- [25] K. Hevner, "Experimental Studies of the Elements of Expression in Music," Am. J. Psychol., vol. 48, no. 2, pp. 246–268, 1936.
- [26] P. R. Farnsworth, "A Study Of The Hevner Adjective List," J. Aesthet. Art Crit., vol. 13, no. 1, pp. 97–103, 1954.
- [27] J. A. Russell, "A Circumplex Model of Affect," J. Pers. Soc. Psychol., vol. 39, no. 6, pp. 1162–1178, 1980.
- [28] R. E. Thayer, *The biopsychology of mood and arousal*. Oxford University Press, 1990.
- [29] X. Hu and J. S. Downie, "Exploring Mood Metadata: Relationships with Genre, Artist and Usage Metadata," *Proc. 8th Int. Conf. Music Inf. Retr. ISMIR*'07, pp. 67–72, 2007.
- [30] T. T. Dang and K. Shirai, "Machine learning approaches for mood classification of songs toward music search engine," *KSE 2009 - 1st Int. Conf. Knowl. Syst. Eng.*, pp. 144–149, 2009.
- [31] R. Dwi Astuti, B. Suhardi, and Jwalitasari, "Analisis Pengaruh Jenis Musik Terhadap Performansi Kerja Operator Input Data," *Semin. Nas. Ergon.*, no. November, pp. 1–13, 2006.
- [32] A. A. K. Ardi and Y. Isnaeni, "Pengaruh Terapi Musik Keroncong Terhadap Kejadian Insomnia Pada Lanjut Usia Di UPT Panti Wredha Budhi Dharma Ponggalan Yogyakarta," 2015.
- [33] C. J. Murrock, "Music and Mood," in Psychology of Moods, 2005.
- [34] O. Lartillot, P. Toiviainen, and T. Eerola, "A Matlab Toolbox for Music Information Retrieval," *Data Anal. Mach. Learn. Appl.*, pp. 261– 268, 2007.
- [35] W. K. Tsao, A. J. T. Lee, Y. H. Liu, T. W. Chang, and H. H. Lin, "A data mining approach to face detection," *Pattern Recognit.*, vol. 43, no. 3, pp. 1039–1049, 2010.