

# ANTICANCER ACTIVITY OF MANGOSTEEN PERICARP DRY EXTRACT AGAINST MCF-7 BREAST CANCER CELL LINE THROUGH ESTROGEN RECEPTOR - $\alpha$

*by Agustina Setiawati*

---

**Submission date:** 05-Feb-2023 10:10PM (UTC+0700)

**Submission ID:** 2006620198

**File name:** 212-163-1-PB.pdf (823.68K)

**Word count:** 2523

**Character count:** 14160

## Research Article

# ANTICANCER ACTIVITY OF MANGOSTEEN PERICARP DRY EXTRACT AGAINST MCF-7 BREAST CANCER CELL LINE THROUGH ESTROGEN RECEPTOR - $\alpha$

Agustina Setiawati<sup>1</sup>, Florentinus Octa Dika Riswanto<sup>1</sup>, Sri Hartati Yuliani<sup>1</sup>,  
Enade Perdana Istyastono<sup>2</sup>

<sup>1</sup>Faculty of Pharmacy,  
Sanata Dharma University,  
Yogyakarta 55281,  
Indonesia

<sup>2</sup>Center for Environmental  
Studies Sanata Dharma  
University (CESSDU),  
Soropadan, Condongcatur,  
Yogyakarta 55283,  
Indonesia

Submitted: 12-03-2014

Revised: 24-04-2014

Accepted: 16-06-2014

\*Corresponding author  
Agustina Setiawati

Email :  
nina@usd.ac.id

## ABSTRACT

Breast cancer has very complex morphological and molecular characteristic. Estrogen receptor is one of biomarker in breast cancer progression, more than 60% breast cancer overexpress estrogen receptor  $\alpha$  (ER $\alpha$ ). Xanthone in *Garcinia mangostana* was investigated whether to have anticancer activity on colorectal, prostate, lung, blood and breast cancer. This research was focused on molecular mechanism of anticancer activity of mangosteen pericarp extract (MPE) on ER- $\alpha$ . This study used MCF-7 cells as a model of ER- $\alpha$  overexpressed breast cancer cells. Cytotoxic study towards MCF-7 cells was designed by using MTT test, further apoptotic induction assay was determined by double staining method using acridine orange and ethidium bromide. Extract molecular mechanism against breast cancer was assayed by immunocytochemistry. The MTT data was analyze using probit analysis to get IC<sub>50</sub> then apoptosis and immunocytochemistry data were analysis qualitative analysis. MPE had strong cytotoxic activity on MCF-7 cells with IC<sub>50</sub> of 45 $\mu$ g/mL and morphological changes passed through apoptosis induction. The expression of ER- $\alpha$  in MPE treated cells was same as untreated cells. MPE did not suppress ER- $\alpha$  in both nucleus and cytoplasm. Anticancer activity of MPE might be mediated by other gene involved in ER- $\alpha$  signaling pathway in breast cancer cells.

**Key words:** Breast cancer, *Garcinia mangostana* peel extract, estrogen receptor- $\alpha$

## INTRODUCTION

The most commonly diagnosed types of cancer among women in 2012 was breast cancer and breast cancer is expected to account for 29% (226,870) of all new cancer cases among women (Siegel *et al.*, 2012). In 2005, 186,467 women suffered from breast cancer and 41,116 of them passed away because of these disease (CDC, 2009). Breast cancer was the major caused death cancer after cervical cancer in Indonesia (Idamardi, 2007).

Breast cancer has very complex morphological and molecular characteristic. One important marker of breast cancer for prognosis is estrogen reseptor (Giacinti *et al.*, 2006). Approximately 60% of breast cancer over expressed estrogen receptor- $\alpha$  (ER- $\alpha$ ) (Hanstein *et al.*, 2004). The development of molecular targeted drug has great advanced to cancer therapy. ER- $\alpha$  is a molecular target to stop the progression of breast cancer.

Mangosteen (*Garcinia mangostana* Linn) pericarp contains various phytochemicals, primary xanthenes. Xanthone isolated from mangosteen pericarps were:  $\alpha$ -mangostin,  $\beta$ -mangostin,  $\gamma$ -mangostin, mangostinone and 2-isoprenyl-1,7 dihydroxy-3-methoxy xanthone inhibited human leukemia HL60 cell line. Among them,  $\alpha$ -mangostin suppressed proliferation and induced apoptosis of HL60 cells at 10 $\mu$ M (Nishimoto *et al.*, 2003). Xanthone extract (81%  $\alpha$ -mangostin and  $\gamma$ -mangostin) suppressed HT116 colon cancer cell (Aisha *et al.*, 2012; Abandith *et al.*, 2004; Nakagawa *et al.*, 2007). Xanthone:  $\alpha$ -mangostin,  $\beta$ -mangostin,  $\gamma$ -mangostin had anticancer activity on DL16 (Akao *et al.*, 2008). Alpha mangostin had antiproliferative activity against BR3 human breast adenocarcinoma cell line (Moongkarndi *et al.*, 2004) and MDA-MB231 (Kurosc *et al.*, 2012). This compound were also proof to reduce lymph node

metastasis<sup>5</sup> in xenograft model in Balb/c mice (Shibata *et al.*, 2011). *Garcinia mangostana* Linn pericarp extract has potential property to be chemopreventive agent against breast cancer.

This study investigated *Garcinia mangostana* pericarp extract (MPE) activity against breast cancer. Furthermore, it focused on ER $\alpha$  as molecular target to suppress proliferation of MCF-7 cell, in vivo model of ER- $\alpha$  over expressed breast cancer cell.

## MATERIAL AND METHODS

MPE was purchased from PT. Borobudur had 28.10%  $\alpha$ -mangosteen. Fructus cortex of *Garcinia mangostana* Linn was extracted use 70% ethanol with ratio botanical extract 10:1 and used maltodextrine as its excipient. MCF-7 breast cancer cell line was cultured in Parasitology laboratory, Faculty of Medicine, Gadjah Mada University. Apoptosis staining using ethidium bromide and acridine orange from SIGMA. This research used US BIO estrogen receptor  $\alpha$  primary antibody.

### MTT Cytotoxic Assay<sup>10</sup>

MCF-7 cells breast cancer cell line were subcultured<sup>15</sup> until confluent. Approximately  $5 \times 10^3$  cells were seeded into 96-well microplate and incubated at 37°C and 5% CO<sub>2</sub> for 24h. Medium was fride treatment control, were prepared from Tamofen® tablet. MPE and tamoxifen were dissolved in DMSO as stock solutions. Various concentration of MPE and tamoxifen in medium were added into 96-well plate 100 $\mu$ L each and incubated in the same as previous condition for 24h. Each concentration was assayed in triplicates (n=3). The medium culture was removed and rinsed by PBS 10%, one hundreds  $\mu$ L into 100 $\mu$ L contain 5mg/mL MTT was added into each well and incubated feather for 4h. Event, medium containing MTT was removed and 100 $\mu$ L SDS was added each well to dissolve the optical density of the resulting solution. The 96-well microplate was incubated for 24h and avoid from light contact. Formazan crystals were detected by ELISA reader using 595nm wavelength.

### Apoptosis Assay

MCF-7 were seeded  $5 \times 10^4$ /100 $\mu$ L into coverslip in 24-well plate, then adapted in 37°C

and 5% CO<sub>2</sub> for 24h. Further, the medium was removed and rinsed by PBS. Seventeen and 34 $\mu$ g/mL of MPE in Dulbecco's Modified Eayle's Medium were added into the well plate and incubated for 24h. These concentrations were determined failure calculation IC<sub>50</sub>, as estimated concentration closed to IC<sub>50</sub>. The medium was removed from MCF-7 cells and rinsed by PBS. The coverslip was removed from well plate and transferred to object glass, and acridine orange-ethidium bromide (AE) was dropped at the coverslip. This slide was soon observed under flouresence microscope.

### Immunocytostaining of ER- $\alpha$

MCF-7 cells in DMEM was seeded 10<sup>5</sup>/100 $\mu$ L on 6-well plate and incubated under 5% CO<sub>2</sub> for 24h. MPE in concentration of 17 and 34 $\mu$ g/mL were applied on cells and incubated for 24 hours. After incubation, cells was collected and washed in PBS. They were resuspended in DMEM and placed in object glass, fixated for 5min, incubated in H<sub>2</sub>O<sub>2</sub> for 10-15min and then washed again using PBS. Monoclonal antibody anti ER- $\alpha$  was and incubated at least for an hour and washed three time using PBS. Secondary antibody of biotinylated goat anti-polyvalent was added and incubated at room temperature for 10min and washed four times in PBS. DAB as chromogen the mistune was dropped and incubated for and 3-8min, washed using aquadest. Finally, hematoxylin solution was added and incubated for 3-4 minutes. This slides were dried and observed under light microscope. Expression of ER- $\alpha$  was showed by brown cell colour.

### Analysis

The cytotoxic data of MPE and tamoxifen were analyzed by excel. IC<sub>50</sub> was then calculated using probit analysis (Finney and Stevens, Any new references?

## RESULT AND DISCUSSION

This study investigated another cytotoxic proper<sup>17</sup> of *Garcinia mangostana* pericarp against breast cancer cell line. MCF-7 was used as in vivo model of over expressed ER- $\alpha$  cells. MPE and tamoxifen we<sup>9</sup> dissolved into excess DMSO as inert solvent in cell culture. Cavalli *et al.* (2011) and Licciardi *et al.* (2010) showed that DMSO 2 $\mu$ L in volume (2% v/v) have no




Figure 1. Effect of MPE on MCF-7 viability after 24 hour incubation




Figure 2. Effect of MPE and tamoxifen on MCF-7 morphology after 24h incubation (A) Untreated cells (B) 40µg/mL MPE treated cells (C) 10-6µg/mL tamoxifen treated cells.

cytotoxic activity against MCF-7 cells. This research used 0.02% v/v DMSO for MPE and oxifen. MTT was absorbed into living cells and converted into purple coloured-formazan complex by succinate dehydrogenase in mitochondria (Doyle and Griffith, 2000). MPE exhibit cytotoxic parabolic profile (Figure 1), probit analysis yielded MPE  $IC_{50}$  of 45µg/mL.

The usage of tamoxifen as positive control is a estrogen receptor  $\alpha$  partial agonis, was regarded as first line drug in ER- $\alpha$  expressed breast cancer treatment (Ao *et al.*, 2011; Fowler *et al.* 2004). Tamoxifen, was also tested against MCF-7 cells by the same procedure. Tamoxifen had parabolic cytotoxic profile and had  $IC_{50}$  47µM or equal to 12,65.10- $\mu$ g/mL by probit analysis. Tamoxifen had strong cytotoxic activity making MCF-7 cells broken into debris but MPE (Figure 2). This

could be explained that tamoxifen is pure chemical compound having specific target on the cells. MPE also driven morphological changes in MCF-7 cells even it did not make the cells broken into debris. MPE contained various natural compounds so it did not have spesific target compared to THAT OFtamoxifen.

Assay then n<sup>o</sup> 19, be designed in other study to confirm molecular mechanism of MCF-7 cells. This study used double staining method using acridine orange and ethidium bromide. Viable cells were green and apoptotic cells were red. MPE showed apoptosis induction towards MCF-7 cells were dose-dependent.

This research used apoptotic cells 17 and 34µg/mL MPE to investigate apoptotic staining. MCF-7 cells were orange and those






Figure 3. Apoptotic cells observation using double staining method under fluorescence microscope using 400x magnification. (A) Untreated cells, (B) Cells treated 17 µg/mL MPE (C) point C: MPE concentration should be "cells treated 34 µg/mL MPE" (as described on the text) not 'cells treated 13 µg/mL'; (D) Cells treated 12,65.10<sup>-7</sup> µg/mL tamoxifen ⇔ living cells; → apoptotic cells.




Figure 4. The observation of ER-α expression using immunocytochemistry method under light microscope using 400x magnification. (A) Untreated cells, (B) Untreated cells without antibody (C) MPE 17 µg/mL treated cells (D) tamoxifen → expressed ER-α

membrane integrity was disturbed (Figure 3). MPE has lower cytotoxic and apoptosis induction activity compare to that of tamoxifen, it might be due to complexity of MPE natural compound. These novel findings underline the benefit of in vitro study to elucidate molecular mechanisms of chemopreventive agent.

The molecular mechanism of apoptosis induction of MPE on ER-α was confirmed using immunocytochemistry method. Tamoxifen, a Selective Estrogen Receptor Modulators (SERMs), suppress ER-α expression. It down regulated ER-α by changing conformation of ER-α thus AF-2 domain is hide so AF-2 dependent co-activator hardly bind to the receptor (Brufsky, 2011). Untreated cells showed high expression of ER-α both in nucleus and cytoplasm. ER-α is located in nucleus and cytoplasm near cell

membrane in the absense and presence of estrogen (Cheskis *et al.*, 2007). MCF-7 cells treated with 34 µg/mL made cells being fragmented into debris there fore this study MPE with concentratum of 17 µg/ml, was used is detect ER-α expression. Even MPE had strong cytotoxic and apoptosis induction activity, it hardly suppressed ER-α expression. Is inhibited ER-α signaling pathway and might have other protein targets on MCF-7. There were 83 genes involved in ER-α signaling pathway in positive ER-α cell lines (Chisamore *et al.*, 2009). Therefore, other genes involved in ER-α signaling pathway must be investigated.

#### ACKNOWLEDGEMENT

This work was funding by Internal Research Grant, Center for Research and Public Services Sanata Dharma University, Yogyakarta

## REFERENCES

- Ao A., Morrison BJ., Wang H., Lopez A., Reynolds BA., Lu J., 2011, *Response of Estrogen Receptor-Positive Breast Cancer Tumorspheres to Antiestrogen Treatments*, 6(4): e18810.
- Aisha AF., Abu-Salah KM., Ismail Z., Majid AMSA. 2012. In vivo and in vivo anti-colon cancer effect of Garcinia mangostana xanthenes extract. *BMC Complementary and Alternative Medicine*, 12: 104- 113
- Akao Y., Nakagawa Y., Linuma M., Nozawa Y. 2008. Anti-Cancer Effects of Xanthenes from pericarps of Mangosteen, *Int. J. Mol. Sci.* 9, 355 – 70.
- Brufsky. 2011. Understanding the estrogen receptor signaling pathway: focus on current endocrine agent for breast cancer in postmenopausal women., *Community Oncology*, 8(8): 343- 352.
- Cavalli A., Bisazza R., Bussano, 2011, Poly(amidoamine)-cholesterol conjugate nanoparticles obtained by electrospraying as novel tamoxifen delivery system, *Journal of Drug Delivery*, Article ID 587604, 9 pages.
- Center for Disease Control and Prevention (CDC). 2009. *Breast Cancer*, available online at <http://www.cdc.gov/cancer/breast/>, accessed on March 3, 2014.
- Cheskis BJ., Greger JG., NagpalS., Freedman LP. 2007. Signaling by estrogens. *J.Cell Physiol.* 213: 610-617.
- Chisamore MJ., Wilkisonm HA., Flores O., Chen JD. 2009. *Estrogen-related receptor- $\alpha$  antagonist inhibits both estrogen receptor-positive and estrogen receptor-negative breast tumor growth in mouse xenograft*, 8: 672- 681.
- Doyle A. and Griffiths JB. 2000. *Cell and Tissue Culture for Medical Research*, John Wiley & Sons LTD, England.
- Finney DJ., Stevens WL. 1948. A Table for Calculation of Working Probits and Weights in Probit Analysis, *Biometrika*, 35(1-2): 191- 201.
- Flowler AM., Solodin N., Mara T., Preisler M., Zhang P., Lee AV, Alarid ET. 2004. Increases in estrogen receptor- $\alpha$  concentration in breast cancer cells promote serine 118/104/106-independent AF-1 transactivation and growth in the absence of estrogen, *FASEB Journal*, 18: 81-93
- Hanstein B., Djahansouzi S., Dall P., Beckmann MW., Bender HG. 2004. *Insight into molecular biology of the estrogen receptor define novel therapeutic targets for breast cancer*, 150: 243- 255.
- Giacinti L., Claudio PP., Lopez M., Giordano A. 2006. Epigenetic Information and Estrogen Receptor Alpha Expression in Breast Cancer. *The Oncologist*. 11(1): 1-8
- Idamardi, 2007. *Kanker Payudara*, Yayasan Kanker Payudara Indonesia, Jakarta.
- Kurose H., Shibata MA., linuma M., Otsuki Y. 2012. Alteration in Cell Cycle and Induction of Apoptotic Cell Death in Breast Cancer Cells Treated with  $\alpha$ -Mangostin Extracted in Mangosteen Pericarp, *J Biomee and Biotec.* 1-9
- Licciardi G., Cavallaro M., Di Stefano G., Pitarresi, C., Fiorica G., Giammona. 2010. New self-assembling polyaspartylhydrazide copolymer micelles for anticancer drug delivery, *Int Journal Pharm*, 396(1-2):219–228.
- Matsumoto K., Akao Y., Kobayashi E., Ohguchi K., Ito T., Tanaka T., Linuma M., Nozawa Y. 2003. Induction of Apoptosis by Xanthenes from Mangosteen in Human Leukemia Cell Lines, *J.Nat. Prod.* 66: 1124- 1127.
- Moongkarndi P., Koscm N., Kaslungka S., Luanratana O., Pongon N., Neungton N. 2004. Antiproliferation, Antioxidation and Induction of Apoptosis by Garcinia mangostana (mangosteen) on SKBR3 human breast cancer cell line. *J.Ethnopharmacol.* 90(1): 161-6.
- Nabandith V., Suzui M., Kaneshiro T., Kinjo T., Matsumoto K., Akao Y., Linuma M., Yoshimi N., Inhibitory Effect of Crude Alpha-Mangostin. 2004. a Xanthone derivative, on two different categories of colon preneoplastic lesion induced by 1,2-dimethylhydrazine in rat. *Asian Pac J Cancer Prev.* 5(4): 433-8.
- Nakagawa Y. Linuma M., Nozawa Y., Akao Y. 2007. Characterized mechanism of  $\alpha$ -mangostin induced cell death: Caspase-independent apoptosis with release of endonuclease-G from mitochondria and

- increased miRNA-143 expression in human colorectal cancer DLD-1 cells. *Bioorg.Med. Chem.* 15: 5620- 5628.
- Siegel R., Naishadham D., Jemal A. Cancer Statistics. 2012. *CA Cancer J. Clin.* 2012. 62:10 – 29.
- Shibata MA., Linuma M., Morimoto J., Kurose H., Akamatsu A., Okuno Y., Akao Y., Otsuki. 2011. A-mangostin extracted from the pericarp of the mangosteen (*Garcinia mangostana* Linn) reduces tumor growth and lymph node metastasis in an immunocompetent xenograft model of metastatic mammary cancer carrying a p53 mutation, *BMC Medicine*, (9): 69.
- Ueda J., Tezuka Y., Baskota AH., Tran QL., Tran QK., Haniyama Y., Saiki I., Kadota S., 2002. Antiproliferative Activity of Vietnamese Medicinal Plants. *Biol.Pharm. Bull.*, 25(6): 753- 760.

# ANTICANCER ACTIVITY OF MANGOSTEEN PERICARP DRY EXTRACT AGAINST MCF-7 BREAST CANCER CELL LINE THROUGH ESTROGEN RECEPTOR - $\alpha$

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

8%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |   |    |
|---|---|----|
| 1 | Submitted to Kuala Lumpur Infrastructure University College<br>Student Paper  | 3% |
| 2 | Submitted to School of Business and Management ITB<br>Student Paper   | 2% |
| 3 | Margarita González-Vallinas, Susana Molina, Gonzalo Vicente, Ruth Sánchez-Martínez et al. "Modulation of estrogen and epidermal growth factor receptors by rosemary extract in breast cancer cells", ELECTROPHORESIS, 2014<br>Publication | 2% |
| 4 | <a href="https://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a><br>Internet Source  | 2% |
| 5 | Mark L. Failla, Fabiola Gutiérrez-Orozco. "Mangosteen Xanthones: Bioavailability and Bioactivities", Wiley, 2017<br>Publication   | 1% |



6	Richa Mardianingrum, Muhammad Yusuf, Maywan Hariono, Amira Mohd Gazzali, Muchtaridi Muchtaridi. "α-Mangostin and its derivatives against estrogen receptor alpha", Journal of Biomolecular Structure and Dynamics, 2020 Publication	1 %
7	fjps.springeropen.com Internet Source	1 %
8	flex.flinders.edu.au Internet Source	1 %
9	cyberleninka.org Internet Source	1 %
10	www.scisoc.or.th Internet Source	1 %
11	Lindsay S. Kalick, Hamaad A. Khan, Erica Maung, Yasmany Baez et al. "Mangosteen for malignancy prevention and intervention: Current evidence, molecular mechanisms, and future perspectives", Pharmacological Research, 2022 Publication	1 %
12	Yun Mi Lee, Eunjung Son, Seung-Hyung Kim, Ohn Soon Kim, Dong-Seon Kim. " Anti-inflammatory and anti-osteoarthritis effect of extract ", Pharmaceutical Biology, 2019 Publication	1 %

13	hdl.handle.net Internet Source	1 %
14	V.P Santhanakrishnan, E. Varun, Javid A. Banday, S. Rajesh, K. Rajamani. "Studies on the invitro anticancer activity of mangostin and acetylated mangostin against MCF-7 cell lines", Chemical Data Collections, 2020 Publication	1 %
15	journals.plos.org Internet Source	1 %
16	Dmitriy Obolskiy. " <i>Garcinia mangostana</i> L.: a phytochemical and pharmacological review", Phytotherapy Research, 08/2009 Publication	<1 %
17	www.dovepress.com Internet Source	<1 %
18	Submitted to Mahidol University Student Paper	<1 %
19	Yuan-Bing Lee. "Î±-Mangostin, A Novel Dietary Xanthone, Suppresses TPA-Mediated MMP-2 and MMP-9 Expressions through the ERK Signaling Pathway in MCF-7 Human Breast Adenocarcinoma Cells", Journal of Food Science, 01/2010 Publication	<1 %
20	www.freepatentsonline.com Internet Source	

<1 %

---

21 [www.sphinxesai.com](http://www.sphinxesai.com)  
Internet Source

<1 %

---

---

Exclude quotes Off  
Exclude bibliography On

Exclude matches Off