



1 of 1

[Download](#) [Print](#) [E-mail](#) [Save to PDF](#) [Save to list](#) [More... >](#)
Journal of Tropical Biodiversity and Biotechnology • Open Access • Volume 5, Issue 1, Pages 53 - 58 • 15 April 2020

Document type

Article • Gold Open Access • Green Open Access

Source type

Journal

ISSN

25409573

DOI

10.22146/jtbb.53635

Publisher

Universitas Gadjah Mada, Faculty of Biology

Original language

English

View less

Cited by 0 documents

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

Related documents

Cultivation of chlorella pyrenoidosa as a raw material for the production of biofuels in palm oil mill effluent medium with the addition of urea and triple super phosphate

Elystia, S. , Muria, S.R. , Erlangga, H.F. (2020) *Environmental Health Engineering and Management*

Growth of Spirulina sp. microalgae in the media of fish waste water treatment from Cemara market, Medan, North Sumatera

Yusni, E. , Hartono, E.S. (2021) *IOP Conference Series: Earth and Environmental Science*

Ethanol of ajwa

Rodliyana, M.D. , Rahmatullah, S. , Gojali, D. (2018) *MATEC Web of Conferences*

View all related documents based on references

Find more related documents in Scopus based on:

Authors > Keywords >

Optimization of Spirulina sp. Growth in Walne Media with Variation of Urea and NaHCO₃ Supplements

[Caturwati, Lusiana Novia](#) ; [Setyati, Retno Herrani](#)
[Save all to author list](#)

^a Department of Biology Education, Faculty of Teacher Training and Education, Sanata Dharma University, Paingan, Maguwoharjo, Depok, Sleman, Special Region of Yogyakarta, 55282, Indonesia

5

Views count

[View all metrics >](#)[View PDF](#)

Full text options

Export

Abstract

Author keywords

Sustainable Development Goals 2022

SciVal Topics

Metrics

Funding details

Abstract

One alternative biofuel to substitute fossil fuels is bioethanol. Microalgae Spirulina sp. contains high carbohydrates, which has 17-25% potential to produce bioethanol. Urea and NaHCO₃ can be used as additional nutrients sources of nitrogen and carbon to Spirulina sp. cultivation. Deficiency of nitrogen causing the cell's enzymes change that shown through decreased lipid and chlorophyll synthesis. While deficiency of carbon can affect the growth rate. In this research, the growth rate of Spirulina sp. is analyzed using Optical Density (OD) method. The growth rate calculation is used to measure the growth of microalgae cells shown in the growth curve. This was a laboratory-scale method using CRD with 4 treatments and 5 replications namely treatment A addition of 0.36 g/500 ml urea without addition of NaHCO₃, treatment B addition of 0.043 g/500 ml NaHCO₃ without addition of urea, treatment C addition of 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃, and control without addition of urea or NaHCO₃. The results indicated that addition of urea and NaHCO₃ didn't affect to OD and Spirulina sp. growth rate. The highest growth rate was treatment A with 0.00906/day of growth rate followed by treatment C which has 0.00865/day of growth rate. Treatment B and control treatment (K) showed a low growth rate. The maximum OD value obtained in treatment C was 0.674 cells/ml on the 10th day. This research can be used as the reference to larger scale of Spirulina sp. cultivation in the field of bioethanol production. © 2020, J. Tropical Biodiversity Biotechnology

Author keywords

Cultivation; Growth rate; Optical density (OD); Spirulina sp

Sustainable Development Goals 2022

SciVal Topics

References (14)

[View in search results format >](#) All [CSV export](#) [Print](#) [E-mail](#) [Save to PDF](#) [Create bibliography](#)

- 1 Ambarwati, Diyah Putri, Yudiaty, Ervia, Supriyantini, Endang, Maslukah, Lilik
Pola Pertumbuhan, Biomassa dan Kandungan Protein Kasar pada Kultur Mikroalga Skeletonema costatum Skala Massal dengan Konsentrasi Kalium Nitrat (KNO₃) yang Berbeda
(2018) *Buletin Oseanogafi Marina*, 2 (7), pp. 75-80.
- 2 Christwardana, Marcellinus, Nur, Muhamad Maulana Azmatun, Hadiyanto, H.
Spirulina platensis: Potensinya Sebagai Bahan Pangan Fungsional
(2012) *Jurnal Aplikasi Teknologi Pangan*, 1 (2), pp. 1-4. Cited 7 times.
- 3 Dewi, Sri Suminar
Teknologi Membran dalam Produksi Bioetanol
(2016) *Institut Teknologi Bandung*, pp. 1-6.
- 4 Hadiyanto, Azim, Maulana
(2012) *Mikroalga: Sumber Pangan dan Energi Masa Depan*, pp. 1-18. Cited 9 times.
UPT UNDIP Press, Semarang
- 5 Juneja, A., Ceballos, R.M., Murthy, G.S.
Effects of environmental factors and nutrient availability on the biochemical composition of algae for biofuels production: A review ([Open Access](#))
(2013) *Energies*, 6 (9), pp. 4607-4638. Cited 487 times.
<http://www.mdpi.com/1996-1073/6/9/4607/pdf>
doi: 10.3390/en6094607
[View at Publisher](#)
- 6 Kawaroe, Mujizat, Prartono, Tri, Saefurahman, Ganjar
Kepadatan dan Laju Pertumbuhan Spesifik *Nannochloropsis* sp. pada Kultivasi Heterotropik Menggunakan Media Hidrolisat Singkong
(2015) *Jurnal Omni-Akuatika*, 11 (2), pp. 15-19. Cited 2 times.
- 7 Melanie, Susiana, Fithriani, Diini
Rendemen Minyak dari Mikroalga *Spirulina* sp. dan *Chlorella* sp. dengan Teknik Pemecahan Dinding Sel
(2015) *Jurnal Widyaerset*, 1 (1), pp. 61-70.
- 8 Muyassaroh, Rini Kartika Dewi, Anggorowati, Dwiana
Kultivasi Mikroalga *Spirulina platensis* dengan Variasi Pencahayaan Menggunakan Lampu TL dan Matahari
(2018) *Prosiding Seminar Nasional Aplikasi Sains & Teknologi (SNAST)*, pp. 381-386.
- 9 Prayitno, Joko
Pola Pertumbuhan dan Pemanenan Biomassa dalam Fotobioreaktor Mikroalga untuk Penangkapan Karbon
(2016) *Jurnal Teknologi Lingkungan*, 1 (17), pp. 45-52. Cited 5 times.
- 10 Sari, Fitria Yuli Anggita, Suryajaya, I Made Aditya, Hadiyanto
Kultivasi Mikroalga *Spirulina platensis* dalam Media POME dengan Variasi Konsentrasi POME dan Komposisi Jumlah Nutrien
(2012) *Jurnal Teknologi Kimia dan Industri*, 1 (1), pp. 487-494. Cited 8 times.

- 11 Seftian, Deky, Antonius, Ferdinand, Faizal, M.
Pembuatan Etanol dari Kulit Pisang Menggunakan Metode Hidrolisis Enzimatik dan Fermentasi
(2012) *Jurnal Teknik Kimia*, 1 (18), pp. 10-16. Cited 3 times.

-
- 12 Suyono, Eko Agus, Haryadi, Winarto
Optimasi Media untuk Produksi Biomassa Mikroalga Chaetoceros sp dan Skeletonema sp Isolat Jepara dan Analisis Kandungan Asam Lemaknya
(2006) *Jurnal Sains dan Teknologi*, pp. 6-8.

-
- 13 Syaichurozzi, Iqbal, Jayanudin
Kultivasi Spirulina platensis pada Media Bernutrisi Limbah Tahu dan Sintetik
(2016) *Jurnal Bahan Alam Terbarukan*, 5 (2), pp. 68-73. Cited 3 times.

-
- 14 Widianingsih, Ali Ridho, Hartati, Retno, Harmoko
Kandungan Nutrisi Spirulina platensis yang Dikultur pada Media yang Berbeda
(2008) *Jurnal Ilmu Kelautan*, 3 (13), pp. 167-170.

✉ Caturwati, L.N.; Department of Biology Education, Faculty of Teacher Training and Education, Sanata Dharma University, Paingan, Maguwoharjo, Depok, Sleman, Special Region of Yogyakarta, Indonesia;
email:lusiananc11@gmail.com

© Copyright 2021 Elsevier B.V., All rights reserved.

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗ All rights reserved. Scopus® is a registered trademark of Elsevier B.V.



We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗



Journal of Tropical Biodiversity and Biotechnology

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Indonesia  Universities and research institutions in Indonesia	Agricultural and Biological Sciences └ Agricultural and Biological Sciences (miscellaneous) └ Ecology, Evolution, Behavior and Systematics Biochemistry, Genetics and Molecular Biology └ Biotechnology Immunology and Microbiology └ Applied Microbiology and Biotechnology	Universitas Gadjah Mada, Faculty of Biology	2
PUBLICATION TYPE	ISSN	COVERAGE	
Journals	25409573, 25409581	2020-2021	

 Join the conversation about this journal



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scim...
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of April 2022



what the level international journal Journal of Tropical Biodiversity and Biotechnology

3 months ago

i cant to see level of journal

reply



Melanie Ortiz 3 months ago

SCImago Team

Dear Sir/Madam,

Thank you for contacting us. Could you please expand a little bit on your comment?

Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:




Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Edit Cookie Consent](#)

Get More with
SINTA Insight[Go to Insight](#)

JOURNAL OF TROPICAL BIODIVERSITY AND BIOTECHNOLOGY

[UNIVERSITAS GADJAH MADA](#)
 P-ISSN : 25409573 <> E-ISSN : 25409581



1.72727
Impact Factor



464
Google Citations



Sinta 1
Current
Acreditation

[Google Scholar](#)

[Garuda](#)


[Website](#)

[Editor URL](#)

History Accreditation

2018 2019 2020 2021 2022 2023 2024 2025

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2017
Citation	464	461
h-index	11	11
i10-index	13	13

[Garuda](#)[Google Scholar](#)

[New Record of Arbuscular Mycorrhizal Fungi \(AMF\) Association with Kebar Grass \(*Biophytum petersianum* Klotzsch.\) in the Grassland Area of Kebar, Tambrauw Regency, West Papua, Indonesia](#)

Universitas Gadjah Mada [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 2 \(2022\): August jtbb70021](#)

2022 DOI: 10.22146/jtbb.70021 Accred : Sinta 1

[The Protective Effect of *Gynura procumbens* Adventitious Root against Lead Acetate Toxicity in Mice](#)

Universitas Gadjah Mada [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 2 \(2022\): August jtbb69453](#)

2022 DOI: 10.22146/jtbb.69453 Accred : Sinta 1

[Pharmacological Maneuver of Mangrove Endophytic Fungi in the South China Sea – A review](#)

Universitas Gadjah Mada [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 2 \(2022\): August jtbb69913](#)

2022 DOI: 10.22146/jtbb.69913 Accred : Sinta 1

[How to link: Plasmid Curing and Lead Tolerance Ability of *Pediococcus pentosaceus*](#)

Universitas Gadjah Mada [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 2 \(2022\): August jtbb68615](#)

2022 DOI: 10.22146/jtbb.68615 Accred : Sinta 1

[Biostratigraphy and Paleobathymetry Microfossil Foraminera in the Sentolo Formation on the Jambon Line, Bantul Regency, Special Region of Yogyakarta Province](#)

[Universitas Gadjah Mada](#)
 (2022): April jtbb62239

 [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)

 2022  DOI: 10.22146/jtbb.62239  Accred : Sinta 1

Get More with
SINTA Insight

[Go to Insight](#)

[Herpetofaunal Assemblages in the Lowland Regions of Sumatera Barat](#)

[Universitas Gadjah Mada](#)  [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)
 (2022): April jtbb63820

 2022  DOI: 10.22146/jtbb.63820  Accred : Sinta 1

Citation Per Year By Google Scholar



[Aboveground Forest Carbon Stock in Protected Area: A Case Study of Bukit Tigapuluh National Park, Indonesia](#)

[Universitas Gadjah Mada](#)  [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)
 (2022): April jtbb64827

 2022  DOI: 10.22146/jtbb.64827  Accred : Sinta 1

[The Effect of Accumulation of Leaf Litters and Allelochemicals in the Soil to the Sustainability of the Newly Introduced Crop Plants](#)

[Universitas Gadjah Mada](#)  [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)
 (2022): April jtbb65227

 2022  DOI: 10.22146/jtbb.65227  Accred : Sinta 1

Journal By Google Scholar

	All	Since 2017
Citation	464	461
h-index	11	11
i10-index	13	13

[Essential Oils Composition of Kaffir Lime \(*Citrus hystrix DC.*\) Collection of Bogor Botanic Gardens from Central Java and East Sumba](#)

[Universitas Gadjah Mada](#)  [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)
 (2022): April jtbb66061

 2022  DOI: 10.22146/jtbb.66061  Accred : Sinta 1

[The Potential of *Trichosanthes tricuspidata* Lour. from Bangli, Baturiti, Bali for Free Radicals Scavenging](#)

[Universitas Gadjah Mada](#)  [Journal of Tropical Biodiversity and Biotechnology Vol 7, No 1](#)
 (2022): April jtbb66111

 2022  DOI: 10.22146/jtbb.66111  Accred : Sinta 1

[View more ...](#)



Journal of Tropical Biodiversity and Biotechnology

□ 2540-9573 (PRINT) / 2540-9581 (ONLINE)

Website

ISSN Portal

Feedback

About Articles

PUBLISHING WITH THIS JOURNAL

฿ The journal charges up to:

1500000 IDR

75 USD

as publication fees (article processing charges or APCs).

There is a waiver policy for these charges.

🔍 Look up the journal's:

- Aims & scope
- Instructions for authors
- Editorial Board
- Double blind peer review

→ This journal checks for plagiarism.

⌚ Expect on average **12 weeks** from submission to publication.

BEST PRACTICE

✍ This journal began publishing in **open access** in 2016. ⓘ

This website uses cookies to ensure you get the best experience. Learn more about DOAJ's privacy policy.

HIDE THIS MESSAGE



SEARCH

MENU

→ Look up their [open access statement](#) and their [license terms](#).

© The author **does not retain unrestricted** copyrights and publishing rights.

→ Learn more about their [copyright policy](#).

Permanent article identifier:

- DOI

Feedback

JOURNAL METADATA

① Publisher

[Universitas Gadjah Mada, Indonesia](#)

Society or institution

Fakultas Biologi, Universitas Gadjah Mada, Indonesia

Manuscripts accepted in

English

② LCC subjects

[Agriculture: Agriculture \(General\)](#)

[Agriculture: Plant culture](#)

Keywords

functional biology tropical biodiversity biotechnology conservation

Added 7 April 2018 • Updated 17 January 2021

SEARCH

Journals

Articles

DOCUMENTATION

API

OAI-PMH



SEARCH

MENU

ABOUT

[About DOAJ](#)[DOAJ team](#)[Ambassadors](#)[Advisory Board & Council](#)[Editorial Subcommittee](#)[Volunteers](#)[News](#)

APPLY

[Application Form](#)[Guide to applying](#)[The DOAJ Seal](#)[Transparency & best practice](#)[Why index your journal in DOAJ?](#)[Licensing & copyright](#)[FAQs](#)

SUPPORT

[Support DOAJ](#)[Contributors](#)[Publisher supporters](#)[Supporters](#)

Feedback

STAY UP TO DATE

[Twitter](#)[Facebook](#)[Github](#)[Linkedin](#)[WeChat](#)[Atom feed](#)

© DOAJ 2022 default by all rights reserved unless otherwise specified.

[Accessibility](#) [Privacy](#) [Contact](#) [T&Cs](#) [Media](#)[IS4OA](#) [Cottage Labs](#)

Content on this site is licensed under a Creative Commons Attribution-ShareAlike 4.0 International (CC BY-SA 4.0) license.

Copyrights and related rights for **article metadata** waived via CC0 1.0 Universal (CC0) Public Domain Dedication.

Photos used throughout the site by David Jorre, Jean-Philippe Delberghe, JJ Ying, Luca Bravo, Brandi Redd, & Christian Perner from Unsplash.

This website uses cookies to ensure you get the best experience. Learn more about DOAJ's privacy policy.

[HIDE THIS MESSAGE](#)

Volume 5, Issue 1, April 2020



Published by:

Faculty of Biology
Universitas Gadjah Mada

In collaboration with:





Journal of Tropical Biodiversity and Biotechnology

online <http://jtbb.or.id>

ISSN 2540-9581 (online)

Menu

[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing & Accreditation](#) [Sitemap](#) [Contact](#)

[Home](#) > [Editorial Team](#)

Editorial Team

Editor-in-Chief



Dr. Miftahul Ilmi
Faculty of Biology, Universitas Gadjah Mada, Indonesia
[Link: Google Scholar, Scopus, Academic Profile](#)

[EDITORIAL TEAM](#)

[Focus & Scope](#)

[Submissions](#)

[Author Guidelines](#)

[Author Fees](#)

[Publication Ethics](#)

[Screening For Plagiarism](#)

[JOURNAL RANK](#)

[ScIImago Journal Country Rank](#)

[TEMPLATE](#)



[WRITING TOOLS](#)

[Sciwheel](#)

[MENDELEY](#)

[EndNote](#)
...Bibliographies Made Easy™

[grammarly](#)

[USER](#)

[Username](#)

[Password](#)

[Remember me](#)

[Login](#)

[JOURNAL CONTENT](#)

Associate Editors



Dr. Ardaning Nuriliani
Faculty of Biology, Universitas Gadjah Mada, Indonesia
[Link: Google Scholar, Scopus, Academic Profile](#)



Dr. Furzani binti Pa'ee
Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia, Malaysia
[Link: Google Scholar, Scopus, Academic Profile](#)

Technical Editors



Sri Nopitasari, M.Sc.
Manuscript Editor
[Link: Google Scholar](#)



Liya Audinah, B.Sc.
Manuscript Editor



Annisa Widayarsi, M.Sc.
Manuscript Editor
[Link: Google Scholar, ResearchGate](#)



Salwa Shabria Wafi, A.Md.
Layout and Copyeditor
[Link: LinkedIn](#)

Editorial Board



Prof. Dr. Wibowo Mangunwardoyo
Department of Biology, FMIPA, Universitas Indonesia, Indonesia
[Link: Google Scholar, Scopus, Orcid ID, ResearchGate, Academic Profile, CV](#)



Prof. Dr. Budi Setiadi Daryono, M.Agr.Sc.
Faculty of Biology, Universitas Gadjah Mada, Indonesia
[Link: Google Scholar, Scopus, Orcid ID, ResearchGate, Academic Profile, CV](#)



Prof. Dr. Jonathan A. Anticamara
Institute of Biology, College of Science, University of the Philippines, Philippines
[Link: Google Scholar, Scopus, Orcid ID, ResearchGate, Academic Profile, CV](#)



Prof. Jean W. H. Yong, Ph.D.
 Dept of Biosystems and Technology, Sveriges lantbruksuniversitet (SLU), Sweden
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Dr. Farid Asif Shaheen
 Department of Entomology, PMAS-Arid Agriculture University Rawalpindi, Pakistan
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Ts. Dr. Kamarul Rahim bin Kamarudin
 Department of Technology and Natural Resources, FAST, Universiti Tun Hussein Onn Malaysia, Malaysia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Assoc. Prof. Dr. Wong Wey Lim
 Department of Biological Science, Faculty of Science, Universiti Tunku Abdul Rahman, Malaysia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Dr. Phoon Lee Quen
 Department of Allied Health Sciences, Faculty of Science, Universiti Tunku Abdul Rahman, Malaysia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Sukirno, M.Sc., Ph.D.
 Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Dr. rer. nat. Andhika Puspito Nugroho
 Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Assoc. Prof. Dr. Ruqiah Ganda Putri Panjaitan
 Biology Education Program, Universitas Tanjungpura, Pontianak, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Dr. Abdul Razaq Chasani
 Faculty of Biology, Universitas Gadjah Mada, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)



Dr. Ratna Stia Dewi
 Faculty of Biology, Universitas Jenderal Soedirman, Purwokerto, Indonesia
 Link: [Google Scholar](#), [Scopus](#), [Orcid ID](#), [ResearchGate](#), [Academic Profile](#), [CV](#)

JOURNAL CONTENT

Search

Search Scope

Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

NOTIFICATIONS

- ▶ View
- ▶ Subscribe

ISSN BARCODES



9 772540 95804

KEYWORDS

Alas Purwo National Park
 Bali DNA barcoding
 DPPH Dendrobium East J:
 LAB Malang altitude
 antioxidant bacteriocin
 biodiversity compositic
 conservation divers
 endophytic fungi
 identification lead
 phenetic relationship reef
 fish vanilla

114915

[View My Stats](#)

Editorial address:

Faculty of Biology, UGM

Jl. Teknika Selatan, Sekip Utara, Yogyakarta, 55281, Indonesia

ISSN: 2540-9581 (online)



Journal of Tropical Biodiversity and Biotechnology

online <http://jtbb.or.id>

ISSN 2540-9581 (online)

Menu

[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing & Accreditation](#) [Sitemap](#) [Contact](#)

[Home](#) > [Archives](#) > [Vol 5, No 1 \(2020\)](#)

Vol 5, No 1 (2020)

April

Full Issue

[View or download the full issue](#)

Table of Contents

Short Communications

- Bioethanol Levels of Dragon Fruit (*Hylocereus polyrhizus*) Peel with the Addition of Blend Crude Cellulase Enzyme from *Trichoderma reesei* and *Aspergillus niger* 1-5

Trianik Widyaningrum, Masreza Parahadi

10.22146/jtbb.52189 Abstract views : 2001 | views : 1457

- Diversity and Distribution of Herpetofauna in Banyu Nibo Waterfall Ecotourism Region, Nglangeran, Gunung Kidul, Special Region of Yogyakarta 6-9

Alfonsus Toribio Eko Saputro, Elika Boscha, Ananto Puradi Nainggolan, Donan Satria Yudha, Rury Eprilurahman

10.22146/jtbb.53637 Abstract views : 1810 | views : 1934

Research Articles

- Vegetation Composition of Savanna Ecosystem as a Habitat For The Komodo Dragon (*Varanus komodoensis*) on Padar and Komodo Islands, Flores East Nusa Tenggara Indonesia 10-15

S. Sutomo

10.22146/jtbb.48280 Abstract views : 2723 | views : 3271

- Conservation, Phytoremediation Potential and Invasiveness Status of Bali Botanic Garden Aquatic Plant Collection 16-26

Farid Kuswantoro

10.22146/jtbb.49312 Abstract views : 147239 | views : 1763

- Tree Species Composition and Natural Regeneration Status of Southeast Region of Bangladesh 27-34

Abir Dey, Aklima Akther

10.22146/jtbb.49988 Abstract views : 1981 | views : 1658

- Characteristics of Shell Bone as an Identification Tool for Turtle Species (Reptiles: Testudines) in Java, Borneo, and Sumatra 35-43

Donan Satria Yudha, Fidelis Triki Sadewa Aritona, Rury Eprilurahman

10.22146/jtbb.47227 Abstract views : 2726 | views : 1760

- Diversity analysis of *Rhacophorus margaritifer* (Schlegel, 1837) in Baturraden based on RAPD markers 44-52

Edition Team

Focus & Scope

Submissions

Author Guidelines

Author Fees

Publication Ethics

Screening For Plagiarism

JOURNAL RANK

ScIImago Journal & Country Rank

TEMPLATE



WRITING TOOLS

Sciwheel

MENDELEY

EndNote
...Bibliographies Made Easy™

grammarly

USER

Username

Password

Remember me

JOURNAL CONTENT

Search

Fajar Nur Sulistyahadi, I Gusti Agung Ayu Ratna Puspitasari, Agus Nuryanto

 10.22146/jtbb.49110  Abstract views : 1439 |  views : 1500

Optimization of *Spirulina* sp. Growth in Walne Media with Variation of Urea and NaHCO₃ Supplements

Lusiana Novia Caturwati, Retno Herrani Setyati

 10.22146/jtbb.53635  Abstract views : 2655 |  views : 2417

Effect of *Arthrospira maxima* and *Chlorella vulgaris* to Lipid Profile and Visceral Fat Index Alteration in Streptozotocin-Induced Hyperglycemia Rats

Mulyati Mulyati, Aprilia Rahmawati, Slamet Widiyanto

 10.22146/jtbb.49937  Abstract views : 1588 |  views : 1373

RAPD Analysis for Genetic Variability Detection of Mutant Soybean (*Glycine max*(L.) Merr)

Didik Wahyudi, Lia Hapsari, S. Sundari

 10.22146/jtbb.53653  Abstract views : 2552 |  views : 2058

53-58

59-67

68-77

Search Scope

All 


Browse

- ▶ By Issue
- ▶ By Author
- ▶ By Title
- ▶ Other Journals

NOTIFICATIONS

- ▶ View
- ▶ Subscribe

CURRENT ISSUE



ISSN BARCODES



KEYWORDS

Alas Purwo National Park
 Bali DNA barcoding
 DPPH Dendrobium East Java
 LAB Malang altitude
 antioxidant bacteriocin
 biodiversity composition
 conservation diversity
 endophytic fungi
 identification lead
 phenetic relationship reef
 fish vanilla

114917

Editorial address:

Faculty of Biology, UGM

Jl. Teknika Selatan, Sekip Utara, Yogyakarta, 55281, Indonesia

ISSN: 2540-9581 (online)

Research Article

Optimation of *Spirulina* sp. Growth in Walne Media with Variation of Urea and NaHCO₃ Supplements

Lusiana Novia Caturwati^{1*}, Retno Herrani Setyati¹

1) Department of Biology Education, Faculty of Teacher Training and Education, Sanata Dharma University, Paingan, Maguwoharjo, Depok, Sleman, Special Region of Yogyakarta 55282

Submitted: 11 October 2019; Accepted: 17 February 2020; Published: 15 April 2020

ABSTRACT

One alternative biofuel to substitute fossil fuels is bioethanol. Microalgae *Spirulina* sp. contains high carbohydrates, which has 17-25% potential to produce bioethanol. Urea and NaHCO₃ can be used as additional nutrients sources of nitrogen and carbon to *Spirulina* sp. cultivation. Deficiency of nitrogen causing the cell's enzymes change that shown through decreased lipid and chlorophyll synthesis. While deficiency of carbon can affect the growth rate. In this research, the growth rate of *Spirulina* sp. is analyzed using Optical Density (OD) method. The growth rate calculation is used to measure the growth of microalgae cells shown in the growth curve. This was a laboratory-scale method using CRD with 4 treatments and 5 replications namely treatment A addition of 0.36 g/500 ml urea without addition of NaHCO₃, treatment B addition of 0.043 g/500 ml NaHCO₃ without addition of urea, treatment C addition of 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃, and control without addition of urea or NaHCO₃. The results indicated that addition of urea and NaHCO₃ didn't affect to OD and *Spirulina* sp. growth rate. The highest growth rate was treatment A with 0.00906/day of growth rate followed by treatment C which has 0.00865/day of growth rate. Treatment B and control treatment (K) showed a low growth rate. The maximum OD value obtained in treatment C was 0.674 cells/ml on the 10th day. This research can be used as the reference to larger scale of *Spirulina* sp. cultivation in the field of bioethanol production.

Keywords: Cultivation, *Spirulina* sp., growth rate, optical density (OD).

INTRODUCTION

As the time flies, biofuel is highly needed for transportation and industry. The fuel that is often used to fulfill human needs is fossil fuel that can't be renewed such as coal, natural gas, and petroleum. In fact, its availability in nature is decreasing. Meanwhile, using biofuel as alternative energy can replace the use of fossil fuel.

The alternative energy is produced from other resources that will not run out and sustainable if it is managed properly. One of the biofuels called bioethanol, made from fermented sugar liquid from carbohydrate source by using microorganisms. Bioethanol developed from wastes that still contains carbohydrate (Seftian *et al.* 2012). According to Dewi (2016), bioethanol can be produced from variety of plants such as corn, cassava, potatoes, sugar cane,

sorghum, and algae.

Microalgae is a potential energy source in the future because it contains carbohydrates that can be processed into several types of compounds such as biodiesel, bioethanol, and methane (Melanie and Diini, 2015; Hadiyanto and Maulana, 2012). Microalgae is an unicellular organism. It has chlorophyll and utilizes the process of photosynthesis to produce biomass. Microalgae is widely used to produce food supplements because of its high protein. The advantages of microalgae compared to other organisms are able to produce biomass and energy supplies in a short time, only need a small area, can be grown on non-productive land, and high growth rate so it is easily to cultivate (Hadiyanto and Maulana, 2012).

Spirulina sp. is a type of microalgae with carbohydrates and high protein. According to Christwardana *et al.* (2012), *Spirulina* sp. contains about 17-25% of carbohydrate and 56-62% of

*Corresponding author

Email: lusiananc11@gmail.com

© 2020, J. Tropical Biodiversity Biotechnology (CC BY-SA 4.0)

protein. The high content of carbohydrate from this microalga cause *Spirulina* sp. potentially produce bioethanol as a renewable energy product. The advantages possess by *Spirulina* sp. from other types of microalgae are relatively fast to reproduce and easier to harvest because it has a large biomass size (Syaichurozzi and Jayanudin, 2016). Cultivation of *Spirulina* sp. can use sea water, fresh water, or brackish water. Cultivation requires appropriate culture media and contains nutrients for *Spirulina* sp. *Spirulina* sp. also requires additional nutrients both macro and micro nutrients for their survival. Those macro nutrients are N, P, C, H, O, Ca, Mg, Na, and K, while the micro nutrients are Fe, Mn, Cu, Zn, B, and cyanocobalamin (Sari *et al.*, 2012).

According to Widianingsih (2008), carbon (C) and nitrogen (N) are the most important elements for *Spirulina* sp. to growth. The carbon is used in respiration, an energy source, and raw material for the formation of additional cells. The lack of carbon in the growth media affects the growth rate. Nitrogen play a role in the formation of proteins and nucleic acid. Ambarwati *et al.* (2018), also say that nitrogen play a role to stimulate vegetative growth and increase cell number. The deficiency of nitrogen can cause the enzymes in cells change that shown through decreased lipid synthesis and chlorophyll synthesis (Juneja *et al.*, 2013). Urea and NaHCO₃ can be used as additional nutrients to fulfill the survival of *Spirulina* sp. Urea is a source of nitrogen (N) while NaHCO₃ is a source of carbon (C). Urea and NaHCO₃ are more economical and easier to obtain than other N and C sources.

The maximum density of *Spirulina* sp. can be seen through the measurement of OD (Optical Density) on growth media. OD value is proportional to the population density of microalgae. Density measurement used to analyze the growth rate of microalgae using OD method. According to Prayitno (2006), the calculation of growth rate used as a measure growth speed of microalgae cells. The results of OD measurements during this cultivation period can be shown through a microalgae curve. The growth curve used as a determinant when microalgae enter the highest density peak. The application of urea and NaHCO₃ are mixed in a *Spirulina* sp. medium at certain doses. One of the growth media of *Spirulina* sp. is Walne's media. According to Widianingsih (2008), Walne's media is a good culture media for *Spirulina* sp. growth. This research is to determine the effect of urea and NaHCO₃ supplies on OD and the growth rate of *Spirulina* sp., to find out the value of *Spirulina* sp. growth rate with urea and NaHCO₃ supplements, and to find out the maximum OD value of *Spirulina* sp.

MATERIALS AND METHODS

This was a laboratory scale method using Completely Randomized Design (CRD) with 4 treatments and 5 replications namely; treatment A addition of 0.36 g/500 ml urea without addition of NaHCO₃, treatment B addition of 0.043 g/500 ml NaHCO₃ without addition of urea, treatment C addition of 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃, and control without addition of urea or NaHCO₃. Microalgae used in this study was *Spirulina* sp. from Balai Pengembangan Teknologi Perikanan Budidaya (BPTPB) Cangkringan, Yogyakarta. This study used Walne media as basic media. Cultivation was carried out for 10 days with batch culture method. The standardization of *Spirulina* sp. cells at the beginning of cultivation cannot be uniform, so the standardization in this study is the volume of inoculants and the volume of media. Optical density (OD) of *Spirulina* sp. measured every 24 hours for 10 days cultivation using Spectrophotometer UV-1800 Shimadzu wavelength 680 nm according to the research of Syaichurozzi and Jayanuddin (2016). The volume of sea water as a solvent of Walne media is 300 ml, volume inoculant 200 ml, Walne media, and vitamin B₁₂ addition respectively 0.3 ml, pH 8.5, salinity 27‰, aeration, and lighting every day for 24 hours.

Spirulina sp. cultivation tools used 500 ml culture bottles, aerator, two 40-watt tube light lamps on each rack, and aeration hose arranged as in Figure 1. The cultivation tools were designed in a such way to adjust BPTPB Cangkringan laboratory conditions, which was the place to do the research.

Researchers used the formula presented by Hirata *et al.* as referred by Kawaroe *et al.* (2009); Kawaroe *et al.* (2015); and Syaichurozzi & Jayanudin (2016) which uses the same formula to calculate the growth rate as follows:

$$k = \frac{\log \frac{N_1}{N_0}}{T_1 - T_0} \times 3.22$$

Where:

- k = growth rate(/day)
- N₁ = microalgae density at time t
- N₀ = microalgae density at time 0
- 3.22 = constant
- T₁ = observation time at time t
- T₀ = observation time at time 0

Data analysis was performed by quantitative descriptive methods and simple statistics by calculating the average value of Optical Density (OD) to get the growth rate and curve making. Data analysis technique used to obtain the most optimal nutrient addition for *Spirulina* sp. conducted




Figure 1. Design of cultivation tools.

quantitatively using the ANOVA test on IBM SPSS Statistics version 20. In this study, the X variable is the addition of urea and NaHCO_3 , while the Y variable represents the Optical Density (OD) value. Before the ANOVA test, normality tests and homogeneity tests were carried out to ensure that the data are normal and homogeneous.

RESULTS AND DISCUSSION

The Density and Growth Pattern of *Spirulina* sp.
Density data were calculated starting from the 0th day of cultivation to the 10th day. Based on Figure 2, it's shown that the highest density value on treatment K (control) was on the 6th day which was 0.492 cells/ml, treatment A on the 10th day which was 0.672 cells/ml, treatment B on the 7th day which was 0.494 cells/ml, and treatment C on the 10th day which was 0.674 cells/ml. The density of *Spirulina* sp. at the beginning cannot be formed so that it has a different value in each treatment. However, differences in density are not very different from one another.

Increased density on day 0 to 1st day indicates that *Spirulina* sp. has adapted to new growth media. Suyono and Winarto (2006), mentioned that in the process of microalgae cell adaptation, they have utilized the nutrients in the media, although not yet optimal. This phase is called the lag phase (induction phase). Muyassaroh (2018) also stated that the growth of *Spirulina* sp. marked with a bluish-green

color in the growth media, while a yellowish color in the growth media shows that *Spirulina* sp. has experienced a phase of death. The thicker bluish green color on the media shows that the growth of *Spirulina* sp. increased both in terms of size and number of cells.

Based on the growth curve from Figure 2, it can be seen that the growth of *Spirulina* sp. each treatment has a different pattern. The density of *Spirulina* sp. treatment K (control) continued to increase since the first day of cultivation and experienced a peak on the 6th day, but decreased from the 7th day to the 10th day. The density of *Spirulina* sp. treatment A continued to increase from the 1st day of cultivation to the 5th day, then down to the 9th day, and increased again on the 10th day. The density of *Spirulina* sp. treatment B continued to increase since the 1st day of cultivation and experienced a peak on the 7th day, then decreased from the 8th to the 10th day, while the density of *Spirulina* sp. treatment C has increased since the 1st day of cultivation and continues to increase until the 10th day. In treatments A and C, the highest peak was on the 10th day but the possibility could still increase until an unknown day. *Spirulina* sp. culture will stay alive as long as the nutrients in the growth media are still available. Decreased density of *Spirulina* sp. in treatment A on the 6th day until the 9th day was caused by aeration on the A3 culture bottle died, thereby reducing the average value of density.

Table 1. Growth Rate of *Spirulina* sp. in Control and 3 Treatment.

Sample	Day										\bar{x}
	1	2	3	4	5	6	7	8	9	10	
K	0.02302	0.01650	0.01157	0.00943	0.00618	0.00356	-0.00043	-0.00214	-0.01058	-0.00941	0.00477
A	0.02501	0.02804	0.01045	0.01015	0.00566	-0.00042	-0.00049	-0.00017	0.00060	0.01183	0.00906
B	0.02904	0.00416	0.01852	0.01342	0.00382	0.00133	0.00192	-0.00510	-0.01156	-0.00736	0.00482
C	0.02518	0.02629	0.01328	0.00052	0.00727	0.00218	0.00343	0.00227	0.00422	0.00188	0.00865

Note: A = medium with 0.36 g/500 ml urea; B = medium with 0.043 g/500 ml NaHCO₃; C = medium with 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃; K = control medium without addition of urea and NaHCO₃.




Figure 2. Growth pattern of *Spirulina* sp. based on average density of control and treated medium. A = medium with 0.36 g/500 ml urea ; B = medium with 0.043 g/500 ml NaHCO₃; C = medium with 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃; K = control medium without addition of urea and NaHCO₃.

The data density of *Spirulina* sp. above showed that each treatment has a different standard deviation. The standard deviation on treatment A, B, C and control treatment (K) in a row were 0.161, 0.115, 0.168, and 0.114. The observations showed that treatments A and C were treated with the highest OD values compared to treatments K (control) and B. Treatment A and C contained the addition of urea nutrients as a source of N (nitrogen). The nitrogen content in urea added in treatments A and C has been shown to increase the growth of *Spirulina* sp. when compared with the control treatment (K). According to Ambarwati *et al.* (2018), the addition of N elements in the cultivation of microalgae with the right amount can optimally increase the population of microalgae. Rauf *et al.* (in Ambarwati *et al.*, 2018) state that in the growth of microalgae, N element plays a role in stimulating vegetative growth and increasing the number of microalgae cells. The increase in growth seen in the growth curve in treatments A and C proved an increase in the growth of *Spirulina* sp. until the last day of cultivation.

Based on the results of normality and homogeneity tests, data on *Spirulina* sp. are normal

and homogeneous ($p > 0.05$). However, the results of the analysis using the ANOVA test showed a significant value is 0.132 ($p > 0.05$), so that further tests could not be carried out. These results indicate that the addition of nutrients in the form of urea and NaHCO₃ does not affect the density of *Spirulina* sp.

Growth Rate of *Spirulina* sp.

In this study, calculation of the growth rate (growth rate/day) is used to determine the rate of growth of *Spirulina* sp. per day. The results of the calculation of the growth rate are presented in Table 1.

Based on the *Spirulina* sp. growth rate data above showed that each treatment has a different standard deviation as same as the density. The standard deviation on treatment A, B, C, and control treatment (K) in a row that are 0.01041, 0.01241, 0.00971, and 0.01082. The average growth rates of *Spirulina* sp. in each treatment are presented in Figure 3.




Figure 3. Comparison of the average growth rate of *Spirulina* sp. in control and treatments medium. A = medium with 0.36 g/500 ml urea; B = medium with 0.043 g/500 ml NaHCO₃; C = medium with 0.36 g/500 ml urea and 0.043 g/500 ml NaHCO₃; K = control medium without addition of urea and NaHCO₃.

Observation of growth rate (k) *Spirulina* sp. in each treatment has different results. The highest average growth rate of *Spirulina* sp. was in treatment A which was 0.00906/day followed by treatment C which was 0.00865/day, while the lowest average

growth rate of *Spirulina* sp. was in the control treatment (K) which was 0.00477/day followed by treatment B which was 0.00482/day. Based on the results of normality and homogeneity, growth rate data of *Spirulina* sp. normal and homogeneous ($p > 0.05$). However, the results of the ANOVA test showed a significant value is 0.707 ($p > 0.05$), so that further tests could not be carried out. Based on the ANOVA test results, the addition of nutrients such as urea and NaHCO₃ did not affect the growth rate of *Spirulina* sp.

Growth Phase of *Spirulina* sp.

Based on the growth pattern curve of *Spirulina* sp. in Figure 2, it can be analyzed the growth phases of *Spirulina* sp. which include:

Lag Phase / Induction Phase

According to Suyono and Winarto (2006), the lag phase is also referred to as the resting phase. On the growth pattern curve *Spirulina* sp. in Figure 3.2 the lag phase of each treatment occurred on day 0 and the 1st day. The density of *Spirulina* sp. in the control treatment (K) on day 0 was 0.147 cells/ml, treatment A was 0.142 cells/ml, treatment B was 0.143 cells/ml, and treatment C was 0.153 cells/ml. The density of each treatment which increased on the 1st day of cultivation where the control treatment (K) reached 0.219 cells/ml, treatment A reached 0.218 cells/ml, treatment B reached 0.235 cells/ml and treatment C reached 0.235 cells/ml. Differences density in each treatments are control treatment (K) of 0.072 cells / ml, treatment A of 0.076 cells/ml, treatment B of 0.092 cells/ml, and treatment C of 0.082 cells/ml. The increase in density on day 0 to the 1st day shows that the culture of *Spirulina* sp. has adapted to the media and its cultural environment. As according to Suyono and Winarto (2006), the lag time depends on cell viability, which is the possibility of cells to be able to live adjusting their environmental conditions. At a young inoculant (exponential phase), *Spirulina* sp. the possibility is still viable and adapts more quickly to the environment, whereas at the age of the older inoculants (stationary phase), the lag phase will last longer. *Spirulina* sp. culture which is used as an inoculant in this study is an inoculant with the age of 10 days which is likely to be in an exponential phase so that the lag phase only lasts for 1 day.

Log Phase

In this phase, the number of *Spirulina* sp. has increased rapidly. According to Suyono and Winarto (2006), this phase proves that microalgae cells have successfully adapted to the new growth media and utilize the nutrients contained in the media. Based on the growth pattern of *Spirulina* sp. in figure 3.2,

the exponential phase of control treatment (K) lasts from the 1st day to the 5th day, treatment A takes place from the 1st day to the 5th day, treatment B takes place from the 1st to the 6th day, and treatment C lasts from the 1st day to the unknown day if cultivation time is continued.

Stationary Phase

In this phase, the growth of *Spirulina* sp. stationary or permanent. This shows that the growth rate of *Spirulina* sp. the same as the rate of death. Based on the growth pattern of *Spirulina* sp. in figure 3.2, the stationary phase of the control treatment (K) lasts from the 6th day to the 7th day, treatment A takes place from the 5th to the 9th day, treatment B takes place on the 7th day, and C treatment is thought to have not yet reached the stationary phase.

Death Phase

In this phase, the decreased growth of *Spirulina* sp. higher than the stationary phase, so the growth pattern tends to decrease. Based on the growth pattern of *Spirulina* sp. in figure 3.2, the death phase in the control treatment (K) lasted from the 8th day to the 10th day, in treatment A it was thought that it had not yet experienced a death phase because the cell density increased again on the 10th day, in treatment B the death phase lasted since the 8th day to the 10th day, and treatment C has not yet experienced a phase of death because cell density still continues to increase until the last day of cultivation.

Maximum OD of *Spirulina* sp.

In this study, the density of *Spirulina* sp. measured every 24 hours for 10 days of cultivation. This is done to get the maximum density (maximum OD) of each treatment. In the control treatment (K), the density of *Spirulina* sp. was on the 6th day with a value of 0.492 cells/ml. In treatment A, the density of *Spirulina* sp. the highest was on the 10th day with a value of 0.672 cells/ml. In treatment B, the highest density of *Spirulina* sp. was on the 7th day with a value of 0.494 cells/ml, while in treatment C the highest density of *Spirulina* sp. was on the 10th day with a value of 0.674 cells/ml. The difference in density is due to each treatment. The growth media is different which causes the nutrient content in the growth media is also different. Thus, this is in accordance with the statement of Widianingsih (2008), that differences in cell density are caused by differences in nutrient content in growth media.

CONCLUSION

The results indicated that addition of urea and NaHCO₃ didn't affect to OD and *Spirulina* sp.

growth rate. OD illustrates population density meanwhile, the increasing OD indicates the growth of Spirulina sp. The growth of Spirulina sp. per unit of time is the growth rate. The highest growth rate was treatment A with addition 0.36 g/500 ml of urea without addition of NaHCO₃ supplements which had 0.00906/day, followed by treatment C with addition 0.36 g/500 ml of urea and 0.043 g/500 ml of NaHCO₃ supplements which had 0.00865/day. Treatment B with addition of 0.043 g/500 ml of NaHCO₃ without addition of urea and the control treatment showed a low growth rate, which were 0.00482/day and 0.00477/day. The maximum OD value obtained in treatment C with addition of 0.36 g/500 ml of urea and 0.043 g/500 ml of NaHCO₃ supplements was 0.674 cells/ml on the 10th day.

ACKNOWLEDGMENTS

The authors are grateful to Lembaga Penelitian dan Pengabdian Masyarakat of Sanata Dharma University for financial support and Balai Pengembangan Teknologi Perikanan Budidaya (BPTPB) Cangkringan to provide research facilities.

REFERENCES

- Ambarwati, Diyah Putri, Ervia Yudiaty, Endang Supriyatini, and Lilik Maslukah, 2018, 'Pola Pertumbuhan, Biomassa dan Kandungan Protein Kasar pada Kultur Mikroalga *Skeletonema costatum* Skala Massal dengan Konsentrasi Kalium Nitrat (KNO₃) yang Berbeda', *Buletin Oseanografi Marina*, 2 (7), pp 75 -80.
- Christwardana, Marcellinus, Muhamad Maulana Azimatun Nur, and H. Hadiyanto, 2012, '*Spirulina platensis*: Potensinya Sebagai Bahan Pangan Fungsional', *Jurnal Aplikasi Teknologi Pangan*, 1 (2) pp 1-4.
- Dewi, Sri Suminar, 2016, 'Teknologi Membran dalam Produksi Bioetanol', Institut Teknologi Bandung, pp 1-6.
- Hadiyanto, and Maulana Azim, 2012, 'Mikroalga: Sumber Pangan dan Energi Masa Depan', UPT UNDIP Press, Semarang, pp 1-18.
- Juneja, Ankita, Ruben Michael Ceballos, and Ganti S. Murthy, 2013, 'Effect of Environmental Factors and Nutrien Availability on the Biochemical Composition of Algae for Biofuels Production', *Journal Energies*, (6) pp 4607-4638.

Kawaroe, Mujizat, Tri Prartono, and Ganjar Saefurahman, 2015, 'Kepadatan dan Laju Pertumbuhan Spesifik *Nannochloropsis* sp. pada Kultivasi Heterotropik Menggunakan Media Hidrolisat Singkong', *Jurnal Omni-Akuatika*, 11 (2) pp 15-19.

Melanie, Susiana and Diini Fitriani, 2015, 'Rendemen Minyak dari Mikroalga *Spirulina* sp. dan *Chlorella* sp. dengan Teknik Pemecahan Dinding Sel', *Jurnal Widya Riset*, 1 (1) pp 61-70.

Muyassaroh, Rini Kartika Dewi, and Dwiana Anggorowati, 2018, 'Kultivasi Mikroalga *Spirulina platensis* dengan Variasi Pencahayaan Menggunakan Lampu TL dan Matahari', *Prosiding Seminar Nasional Aplikasi Sains & Teknologi (SNAST)*, Pp 381-386.

Prayitno, Joko, 2016, 'Pola Pertumbuhan dan Pemanenan Biomassa dalam Fotobioreaktor Mikroalga untuk Penangkapan Karbon', *Jurnal Teknologi Lingkungan*, 1 (17) pp 45-52.

Sari, Fitria Yuli Anggita, I Made Aditya Suryajaya, and Hadiyanto, 2012, 'Kultivasi Mikroalga *Spirulina platensis* dalam Media POME dengan Variasi Konsentrasi POME dan Komposisi Jumlah Nutrien', *Jurnal Teknologi Kimia dan Industri*, 1 (1) pp 487-494.

Seftian, Deky, Ferdinand Antonius, and M. Faizal, 2012, 'Pembuatan Etanol dari Kulit Pisang Menggunakan Metode Hidrolisis Enzimatik dan Fermentasi', *Jurnal Teknik Kimia*, 1 (18) pp 10-16.

Suyono, Eko Agus, and Winarto Haryadi, 2006, 'Optimasi Media untuk Produksi Biomassa Mikroalga *Chaetoceros* sp dan *Skeletonema* sp Isolat Jepara dan Analisis Kandungan Asam Lemaknya', *Jurnal Sains dan Teknologi*, Pp 6-8.

Syaichurozzi, Iqbal, and Jayanudin, 2016, 'Kultivasi *Spirulina platensis* pada Media Bernutrisi Limbah Cair Tahu dan Sintetik', *Jurnal Bahan Alam Terbarukan*, 5 (2) pp 68-73.

Widianingsih, Ali Ridho, Retno Hartati, and Harmoko, 2008, 'Kandungan Nutrisi *Spirulina platensis* yang Dikultur pada Media yang Berbeda', *Jurnal Ilmu Kelautan*, 3 (13) pp 167-170.