

Volume 40, Issue 21, 2022

ISSN: 0739-1102

Journal of **Biomolecular Structure and Dynamics**



 Taylor & Francis
Taylor & Francis Group

Journal of Biomolecular Structure and Dynamics

Publish with us

Submit an article ▼

About this journal ▼

Explore

Browse all articles & issues ▼

 Latest issue

Subscribe

Alerts & RSS feed ▼

+ Purchase a subscription




Log in

| Register



Cart



Browse this journal

▶ Latest articles

▶ Current issue

▶ List of issues

▶ Special issues

▶ Open access articles

▶ Most read articles

▶ Most cited articles

Journal of Biomolecular Structure and Dynamics

Sign in here to start your access
to the latest two volumes for 14 days

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

See all volumes and issues

	Volume 40, 2022	Vol 39, 2021	Vol 38 
---	------------------------	---------------------	---

 ssue Issue Issue Issue 
20 19 18 17

 Browse by section (All) | 

 Display order (Default) | 



 Download citations  Download PDFs

Research Articles



Article

Design, synthesis, and molecular dynamics simulation studies of quinoline derivatives as protease inhibitors against SARS-CoV-2 

Vishal K. Singh, Himani Chaurasia, Priyanka Kumari, Anup Som, Richa Mishra, Ritika Srivastava, Farha Naaz, Anuradha Singh & Ramendra K. Singh

Pages: 10519-10542

Published online: 12 Jul 2021

Journal of Biomolecular Structure and Dynamics



Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription

▼	▼	▼
320	7	1
Views	CrossRef citations	Altmetric



Article

***In silico* identification of antidiabetic target for phytochemicals of *A. marmelos* and mechanistic insights by molecular dynamics simulations** >

Priyanka Sharma, Tushar Joshi, Shalini Mathpal, Subhash Chandra & Sushma Tamta

Pages: 10543-10560

Published online: 06 Jul 2021

228	3	1
Views	CrossRef citations	Altmetric



Article

Molecular docking and simulation studies of flavonoid compounds against PBP-2a of methicillin-resistant *Staphylococcus aureus* >

Abhishek Kumar Verma, Sk. Faisal Ahmed, Md. Shahadat Hossain, Ali Asger Bhojiya, Ankita Mathur, Sudhir K. Upadhyay, Abhishek K. Srivastava, Naveen Kumar Vishvakarma, Mayadhar Barik, Md. Mizanur Rahaman & Newaz Mohammed Bahadur

Pages: 10561-10577

Published online: 09 Jul 2021

630	8	1
Views	CrossRef citations	Altmetric

Journal of Biomolecular Structure and Dynamics

[Kinetic epitopes of *Leishmania donovani*: Potential application in vaccine development](#) >

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



139
Views



2
CrossRef citations



1
Altmetric



Article

[An enhanced methodology for predicting protein-protein interactions between human and hepatitis C virus via ensemble learning algorithms](#) >

Xin Liu, Liang Wang, Cheng-Hao Liang, Ya-Ping Lu, Ting Yang & Xiao Zhang

Pages: 10592-10602

Published online: 11 Jul 2021

117

Views

0

CrossRef citations

0

Altmetric



Article

[In silico modeling and empirical study of 4-*n*-Butylresorcinol nanoliposome formulation](#) >

Rini Dwiastuti, Muhammad Radifar, Dina Christin Ayuning Putri, Florentinus Dika Octa Riswanto & Maywan Hariono

Pages: 10603-10613

Published online: 08 Jul 2021

98

Views

0

CrossRef citations

0

Altmetric



Article

Journal of Biomolecular Structure and Dynamics

Pages: 10614-10628

Published online: 24 Jul 2021

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Article

Plant derived active compounds as potential anti SARS-CoV-2 agents: an *in-silico* study >

Dharmendra Kashyap, Shweta Jakhmola, Deeksha Tiwari, Rajesh Kumar, N. S. Hari Narayana Moorthy, Manivannan Elangovan, Natércia F. Brás & Hem Chandra Jha

Pages: 10629-10650

Published online: 06 Jul 2021

514

Views

9

CrossRef citations

2

Altmetric



Article

Spectroscopic, quantum chemical, molecular docking and molecular dynamics investigations of hydroxylic indole-3-pyruvic acid: a potent candidate for nonlinear optical applications and Alzheimer's drug >

Stève-Jonathan Koyambo-Konzapa, Gilbert Yvon Mbesse Kongbonga, Premkumar R, Berthelot Saïd Duvalier Ramlina Vamhindi, Mama Nsangou & A. Milton Franklin Benial

Pages: 10651-10664

Published online: 15 Jul 2021

144

Views

5

CrossRef citations

1

Altmetric



Article

In silico identification of potential Hsp90 inhibitors via ensemble docking, DFT and molecular dynamics simulations >

Journal of Biomolecular Structure and Dynamics



Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription

242 Views	2 CrossRef citations	Altmetric
--------------	-------------------------	-----------



Article

Tuning the structural and catalytic properties of copper(II)-based complexes containing pyridine-2,6-diimines >

Mohamed M. Ibrahim, Shaban Y. Shaban, Ahmed A. Amer, Soha F. Mohamed, Ahmed M. Fathy, Sami A. Al-Harbi & Abd El-Motaleb M. Ramadan

Pages: 10677-10695

Published online: 26 Aug 2021

135 Views	1 CrossRef citations	Altmetric
--------------	-------------------------	-----------



Article

In silico design of epitope-based peptide vaccine against non-typhoidal *Salmonella* through immunoinformatic approaches >

Md. Chayan Ali, Mst. Shanzeda Khatun, Sultana Israt Jahan, Raju Das, Yeasmin Akter Munni, Md Mafizur Rahman & Raju Dash

Pages: 10696-10714

Published online: 11 Jul 2021

224 Views	1 CrossRef citations	Altmetric
--------------	-------------------------	-----------

Journal of Biomolecular Structure and Dynamics

anticancer, BSA and DNA interactions, molecular docking and antioxidant studies >

Publish
with us

Explore


Subscribe

 Latest
issue

 Purchase a
subscription



Published online: 09 Jul 2021



204
Views

2
CrossRef citations

Altmetric



Article

Synthesis, α -amylase and α -glucosidase inhibition and molecular docking studies of indazole derivatives >

Muhammad Nawaz, Muhammad Taha, Faiza Qureshi, Nisar Ullah, Manikandan Selvaraj, Sumaira Shahzad, Sridevi Chigurupati, Samar A. Abubshait, Tauqir Ahmad, Sampath Chinnam & Soleiman Hisaindee

Pages: 10730-10740

Published online: 31 Aug 2021

331
Views

5
CrossRef citations

Altmetric



Article

Identification of potential phytochemicals from *Citrus Limon* against main protease of SARS-CoV-2: molecular docking, molecular dynamic

Journal of Biomolecular Structure and Dynamics

Pages: 10741-10752

Published online: 19 Jul 2021


Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription



434

Views

14

CrossRef citations

Altmetric



Article

Computational studies on phylogeny and drug designing using molecular simulations for COVID-19 >

Faisal Nabi, Owais Ahmad, Yawar Ali Khan, Anas Nabi, Hashmi Md Amiruddin, Faizan Abul Qais, Aiman Masroor, Malik Hisamuddin, Vladimir N. Uversky & Rizwan Hasan Khan

Pages: 10753-10762

Published online: 19 Jul 2021

357

Views

8

CrossRef citations

Altmetric



Article

One microsecond MD simulations of the SARS-CoV-2 main protease and hydroxychloroquine complex reveal the intricate nature of binding >

Prateek Kumar, Taniya Bhardwaj, Ankur Kumar, Neha Garg & Rajanish Giri

Pages: 10763-10770

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



[Mechanism of viral and host proteins by spike protein](#) >

Kottapalli Srividya, Snober S. Mir, Saravanamuthu Thiyagarajan & Aamir Nazir

Pages: 10771-10782

Published online: 13 Jul 2021

163

Views

0

CrossRef citations

Altmetric



Article

[Data mining and network analysis reveals C-X-C chemokine receptor type 5 is involved in the pathophysiology of age-related macular degeneration](#) >

Madhu Sudhana Saddala, Anton Lennikov, Anthony Mukwaya, Xu Yang, Shibo Tang & Hu Huang

Pages: 10783-10792

Published online: 09 Jul 2021

212

Views

1

CrossRef citations

Altmetric



Article

[Non-equilibrium molecular dynamics study of human aquaporin-2 in the static external electric fields](#) >

Hooman Hadidi & Reza Kamali

Pages: 10793-10801

Published online: 09 Jul 2021

83

1

Journal of Biomolecular Structure and Dynamics

Article

Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription

Published online: 21 Jul 2021

254	1	
Views	CrossRef citations	Altmetric

Article

Drug repurposing based novel anti-leishmanial drug screening using *in-silico* and *in-vitro* approaches >

Praveen Rai, Hemant Arya, Satabdi Saha, Diwakar Kumar & Tarun Kumar Bhatt

Pages: 10812-10820

Published online: 11 Jul 2021

244	1	
Views	CrossRef citations	Altmetric

Article

Structural insights into the mechanism of human methyltransferase hPRMT4 >

Amar Pratap Singh, Rakesh Kumar & Dinesh Gupta

Pages: 10821-10834

Published online: 24 Jul 2021

305	0	
Views	CrossRef citations	Altmetric

Journal of Biomolecular Structure and Dynamics

antioxidant activity of some halogenated bisphenols >

Saman Zare Gheshlaghi, Ali Ebrahimi & Zeinab Faghih

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Home | Section Editors | Authors



Article

Understanding the co-evolutionary molecular mechanisms of
resistance in the HIV-1 Gag and protease >

Veronna Marie & Michelle Gordon

Pages: 10852-10861

Published online: 12 Jul 2021

127

Views

2

CrossRef citations

Altmetric



Article

Polymorphisms at site 469 of B-Raf protein associated with skin
melanoma may be correlated with dabrafenib resistance: An *in silico*
study >

Samadrita Chatterjee, Rajkumar Chakraborty & Yasha Hasija

Pages: 10862-10877

Published online: 19 Jul 2021

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription

protein associated with skin melanoma may be correlated with dabrafenib resistance: an in-silico study.

Molecular Dynamic Simulation of variants and drug interaction

binding affinity
BRAF G469E

145

Views

0

CrossRef citations

Altmetric



Article

Unveiling the mechanistic roles of chlorine substituted phthalazinone-based compounds containing chlorophenyl moiety towards the differential inhibition of poly (ADP-ribose) polymerase-1 in the treatment of lung cancer >

Felix O. Okunlola, Fisayo A. Olotu & Mahmoud E. S. Soliman

Pages: 10878-10886

Published online: 31 Aug 2021

162

Views

1

CrossRef citations

Altmetric



Article

Network pharmacological evaluation of *Withania somnifera* bioactive phytochemicals for identifying novel potential inhibitors against neurodegenerative disorder >

Sonu Pahal, Ayushi Gupta, Princy Choudhary, Amit Chaudhary & Sangeeta Singh

Pages: 10887-10898

Published online: 19 Jul 2021

352

5

Journal of Biomolecular Structure and Dynamics

○ Article

Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription

▼ ▼ ▼

Pages: 10899-10904

Published online: 31 Aug 2021

85

Views

1

CrossRef citations

Altmetric


○ Article

Protein degradation: a novel computational approach to design protein degrader probes for main protease of SARS-CoV-2 >

Muhammed Shaheer, Ravi Singh & M. Elizabeth Sobhia

Pages: 10905-10917

Published online: 30 Jul 2021



1125

Views

7

CrossRef citations

Altmetric

○ Article

Journal of Biomolecular Structure and Dynamics

calculation study >

Childhood type 1 diabetes mellitus and risk factor of interactions

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Views | CrossRef citations | Altmetric



Article

Childhood type 1 diabetes mellitus and risk factor of interactions between dietary cow's milk intake and HLA-DR3/DR4 genotype >

Yasmine Benslama, Nouria Dennouni-Medjati, Majda Dali-Sahi, Fatima Zahra Meziane & Yahia Harek

Pages: 10931-10939

Published online: 20 Jul 2021

144

Views

2

CrossRef citations

Altmetric



Article

Molecular insights into *MYO3A* kinase domain variants explain variability in both severity and progression of DFNB30 hearing impairment >

Amal Souissi, Dorra Abdelmalek Driss, Imen Chakchouk, Mariem Ben Said, Ikhlas Ben Ayed, Mohamed Ali Mosrati, Ines Elloumi, Abdelaziz Tlili, Sami Aifa & Saber Masmoudi

Pages: 10940-10951

Published online: 23 Aug 2021

111

Views

1

CrossRef citations

Altmetric



Article

Journal of Biomolecular Structure and Dynamics

Y. Sheena Mary, Y. Shyma Mary, Stevan Armaković, Sanja J. Armaković, Rohitash Yadav, Ismail Celik & Razieh Razavi

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Views | CrossRef citations | Altmetric



Article

In silico approach to target PI3K/Akt/mTOR axis by selected *Olea europaea* phenols in *PIK3CA* mutant colorectal cancer >

Arindam Sain, Thirukumaran Kandasamy & Debdut Naskar

Pages: 10962-10977

Published online: 23 Jul 2021

267

Views

3

CrossRef citations

Altmetric



Article

Functional binding dynamics relevant to the evolution of zoonotic spillovers in endemic and emergent *Betacoronavirus* strains >

Patrick Rynkiewicz, Miranda L. Lynch, Feng Cui, André O. Hudson & Gregory A. Babbitt

Pages: 10978-10996

Published online: 21 Jul 2021

171

Views

4

CrossRef citations

Altmetric



Article

Structure-function correlations and system dynamics in oxygenic photosynthesis: classical perspectives and murburn precepts >

Journal of Biomolecular Structure and Dynamics

Published online: 29 Jul 2021

Publish
with us



Explore



 Latest
issue



Subscribe



 Purchase a
subscription



Article

Validating the predictions of murburn model for oxygenic photosynthesis: Analyses of ligand-binding to protein complexes and cross-system comparisons >

Kelath Murali Manoj, Daniel Andrew Gideon, Abhinav Parashar, Vijay Nirusimhan, Pushparaj Annadurai, Vivian David Jacob & Afsal Manekkathodi

Pages: 11024-11056

Published online: 30 Jul 2021

167

Views

10

CrossRef citations

Altmetric



Article

HSA nanoparticles in drug recognition: mechanistic insights with naproxen, diclofenac and methimazole >

Pooja Prasanthan & Nand Kishore

Pages: 11057-11069

Published online: 23 Jul 2021

285

Views

1

CrossRef citations

Altmetric



Article

Inhibition of multiple SARS-CoV-2 proteins by an antiviral biomolecule, seselin from *Aegle marmelos* deciphered using molecular docking analysis >

Journal of Biomolecular Structure and Dynamics

Published online: 25 Aug 2021

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Article

New imino-methoxy derivatives: design, synthesis, characterization, antimicrobial activity, DNA interaction and molecular docking studies >

Sultan Onur, Mustafa Çeşme, Muhammet Köse & Ferhan Tümer

Pages: 11082-11094

Published online: 06 Aug 2021

273

Views

5

CrossRef citations

Altmetric




Article

Design and *in silico* investigation of novel Maraviroc analogues as dual inhibition of CCR-5/SARS-CoV-2 M^{pro} >

G. Mahaboob Basha, Rishikesh S. Parulekar, Abdullah G. Al-Sehemi, Mehboobali Pannipara, Vidavalur Siddaiah, Sunanda Kumari, Prafulla B. Choudhari & Yasinalli Tamboli

Pages: 11095-11110

Published online: 26 Jul 2021



Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

Sayed Samina Ahmed & Md Ackas Ali

Pages: 11111-11124

Published online: 27 Jul 2021

372

Views

3

CrossRef citations

Altmetric



Article

Identification of the structural features of quinazoline derivatives as EGFR inhibitors using 3D-QSAR modeling, molecular docking, molecular dynamics simulations and free energy calculations >

Fangfang Wang, Wei Yang, Hongping Liu & Bo Zhou

Pages: 11125-11140

Published online: 02 Aug 2021

115

Views

1

CrossRef citations

Altmetric



Article

Salinomycin mediated therapeutic targeting of circulating stem like cell population in oral cancer >

Shanaya Patel, Aditi Patel, Aishwarya Nair, Kavan Shah, Kanisha Shah, Vivek Tanavde & Rakesh Rawal

Pages: 11141-11153

Published online: 26 Jul 2021

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

simulation and multi-spectroscopic perspectives >

Soroush Behjati Hosseini, Maryam Asadzadeh-Lotfabad, Maryam Erfani, Fatemeh Babayan-Mashhadi, Parisa Mokaberi, Zeinab Amiri-Tehranizadeh, Mohammad Reza Saberi & Jamshidkhan Chamani

Pages: 11154-11172

Published online: 30 Jul 2021

115	1	
Views	CrossRef citations	Altmetric



Article

Deleterious single nucleotide polymorphisms (SNPs) of human IFNAR2 gene facilitate COVID-19 severity in patients: a comprehensive *in silico* approach >

Shamima Akter, Arpita Singha Roy, Mahafujul Islam Quadery Tonmoy & Md Sajedul Islam

Pages: 11173-11189

Published online: 06 Aug 2021

370	1	
Views	CrossRef citations	Altmetric



Article

Modeling the structure and reactivity landscapes of a pyrazole-ammonium ionic derivative using wavefunction-dependent characteristics and screening for potential anti-inflammatory activity >

Jamelah S. Al-Otaibi, Y. Sheena Mary, Shiji Fazil, Y. Shyma Mary & S. Sarala

Journal of Biomolecular Structure and Dynamics

Views | CrossRef citations | Altmetric

Publish
with us

Explore

Subscribe

Latest
issue


+ Purchase a
subscription

Andrographis paniculata: An insilico approach >

Gopishankar Thirumoorthy, Sharma Pooja Tarachand, Praveen Nagella & Vasantha Veerappa Lakshmaiah

Pages: 11203-11215

Published online: 28 Jul 2021



230
Views

2

CrossRef citations

Altmetric



Article

Polyols, increasing global stability of cytochrome c, destabilize the thermal unfolding intermediate >

K. Tejaswi Naidu & N. Prakash Prabhu

Journal of Biomolecular Structure and Dynamics

Views | CrossRef citations | Altmetric

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Computational study of molecular docking mechanism of natural compounds against protein–ligand interaction for the *E. coli* MurB enzyme >

Sonam Nirwan, Varun Chahal & Rita Kakkar

Pages: 11229-11238

Published online: 29 Jul 2021

98

Views

0

CrossRef citations

Altmetric



Article

Evolving ribonucleocapsid assembly/packaging signals in the genomes of the human and animal coronaviruses: targeting, transmission and evolution >

Vladimir R. Chechetkin & Vasily V. Lobzin

Pages: 11239-11263

Published online: 02 Aug 2021

122

Views

1

CrossRef citations

Altmetric



Article

Combined molecular docking and dynamics simulations studies of natural compounds as potent inhibitors against SARS-CoV-2 main protease >

Mebarka Ouassaf, Salah Belaidi, Samir Chtita, Touhami Lanez, Faizan Abul Qais & Hashmi Md Amiruddin

Pages: 11264-11273

Published online: 27 Jul 2021

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

Puspita Halder & Pralay Mitra

Pages: 11274-11290

Published online: 02 Aug 2021

177

Views

0

CrossRef citations

Altmetric

 Article

Accuracy and precision of binding free energy prediction for a tacrine related lead inhibitor of acetylcholinesterase with an arsenal of supercomputerized molecular modelling methods: a comparative study >

Rafael Dolezal

Pages: 11291-11319

Published online: 29 Jul 2021

157

Views

0

CrossRef citations

Altmetric

 Article

Discovery of potential inhibitors for stat3: ligand based 3D pharmacophore, virtual screening, molecular docking, dynamic studies and *in vitro* evaluation >

Kaviarasan Lakshmanan, Praveen T. K., Sreedhara Ranganath K. Pai, Kalirajan Rajagopal & Gowramma Byran

Pages: 11320-11338

Published online: 31 Aug 2021

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe


Latest
issue

+ Purchase a
subscription

Anacleto Silva de Souza, Robson Francisco de Souza & Cristiane Rodrigues Guzzo

Pages: 11339-11356

Published online: 09 Aug 2021



384
Views

4
CrossRef citations

Altmetric



Article

Designing Self-Inhibitory fusion peptide analogous to viral spike protein against novel severe acute respiratory syndrome (SARS-CoV-2) >

Indra Singh, Shalini Singh, Krishna Kumar Ojha & Neetu Singh Yadav

Pages: 11357-11372

Published online: 11 Aug 2021

Journal of Biomolecular Structure and Dynamics



Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription



Article

**Antitumor drugs effect on the stability of double-stranded DNA:
steered molecular dynamics analysis** >

Rakesh Kumar Mishra & Lakshmi Maganti

Pages: 11373-11382

Published online: 06 Aug 2021

106

Views

0

CrossRef citations

Altmetric



Article

***In silico* screening of FDA approved drugs against ACE2 receptor:
potential therapeutics to inhibit the entry of SARS-CoV-2 to human
cells** >

Selvaraj Ayyamperumal, Dhananjay Jade, Vyshnavi Tallapaneni, M. J. N. Chandrasekar & M. J. Nanjan

Pages: 11383-11394

Published online: 28 Aug 2021

Journal of Biomolecular Structure and Dynamics



Publish
with us

Explore

Subscribe

Latest
issue

+ Purchase a
subscription



153

Views

0

CrossRef citations

Altmetric



Article

Anticorrelated position fluctuation of lipids in forming membrane water pores: molecular dynamics simulations study with dengue virus capsid protein >

Suneth P. Rajapaksha & Sithara U. Nawagamuwage

Pages: 11395-11404

Published online: 03 Aug 2021

142

Views

0

CrossRef citations

Altmetric



Article

Structure prediction and discovery of inhibitors against phosphopantothenoyl cysteine synthetase of *Acinetobacter baumannii* >

Akshita Gupta, Viswanathan Vijayan, Pradeep Pant, Punit Kaur, Tej P. Singh, Pradeep Sharma & Sujata Sharma

Pages: 11405-11417

Published online: 04 Aug 2021

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

Reyhan Metin & Ebru Demet Akten

Pages: 11418-11433

Published online: 06 Aug 2021

162

Views

1

CrossRef citations

Altmetric



Article

N1 neuraminidase of H5N1 avian influenza A virus complexed with sialic acid and zanamivir – A study by molecular docking and molecular dynamics simulation >

R. A. Jeyaram & C. Anu Radha

Pages: 11434-11447

Published online: 09 Aug 2021

235

Views

2

CrossRef citations

Altmetric

Review Article



Review Article

Computational study of DMPC liposomes loaded with the N-(2-Hydroxyphenyl)-2-propylpentanamide (HO-AAVPA) and determination of its antiproliferative activity *in vitro* in NIH-3T3 cells >

Xelhua Marcos, Yudibeth Sixto-López, Silvia Pérez-Casas & José Correa-Basurto

Pages: 11448-11459

Journal of Biomolecular Structure and Dynamics

Publish
with us

Explore

Subscribe

 Latest
issue


 Purchase a
subscription

In-silico study on viability of MXenes in suppressing the coronavirus infection and distribution >

Ebrahim Ghasemy, Ahmad Miri Jahromi, Mohammad Khedri, Pegah Zandi, Reza Maleki & Lobat Tayebi

Pages: 11460-11466

Published online: 30 Jul 2021



205
Views

2
CrossRef citations

Altmetric

Explore articles

Latest

Open access

Most read

Most cited

Trending

Explore the most recently published articles

Journal of Biomolecular Structure and Dynamics

William M. Neal et al.

Publish
with us

Explore

Subscribe

 Latest
issue

 Purchase a
subscription

Naaila Tamkeen et al.

Article | Published online: 13 Apr 2023

Design and characterization of a binary CT complex of imidazole-oxyresveratrol: exploring its pharmacological and computational aspects >

Nisat Alam et al.

Article | Published online: 13 Apr 2023

Vitamin D analog calcitriol for breast cancer therapy; an integrated drug discovery approach >

Nagaraj B. S. et al.

Article | Published online: 13 Apr 2023

[View all latest articles](#) >

Journal of Biomolecular Structure and Dynamics

Authors

R&D professionals

Publish
with us



Explore



Overview

Open journals

Subscribe



Latest
issue



Purchase a
subscription

Opportunities

Reprints and e-prints

Advertising solutions

Accelerated publication

Corporate access solutions

Help and information

Help and contact

Newsroom

All journals

Books

Keep up to date

Register to receive personalised research and resources
by email



Sign me up



[Log in](#)[Register](#)[Cart](#)

[Home](#) ▶ [All Journals](#) ▶ [Journal of Biomolecular Structure and Dynamics](#) ▶ [Editorial Board](#)

Journal of Biomolecular Structure and Dynamics

Publish with us

[Submit an article](#) ▼

[About this journal](#) ▼

Explore

[Browse all articles & issues](#) ▼



[Latest issue](#)

Subscribe

[Alerts & RSS feed](#) ▼



[Purchase a subscription](#)

Ready to submit?

Start a new manuscript submission or continue a submission in progress

[Go to submission site](#)

Submission information

[▶ Instructions for authors](#)

[▶ Editorial policies](#)

Editing services

[▶ Editing services site](#)

About this journal

[▶ Journal metrics](#)

[▶ Aims & scope](#)

[▶ Journal information](#)

[▶ Editorial board](#)

[▶ News & call for papers](#)

[▶ Advertising information](#)

Editorial board

Editor-in-Chief

Ramaswamy H. Sarma

Institute of Biomolecular Stereodynamics, Department of Chemistry

The University at Albany, Albany NY 12222 USA

Ph: 518-456-9362/518-442-4454; Fx:

518-452-4955

; Email: rhs07@albany.edu

Associate Editors

Mohd Adnan

University of Ha'il, Saudi Arabia

Imtiyaz Hassan

Centre for Interdisciplinary Research in Basic Sciences, India

Vsevolod Makeev

Vavilov Institute of General Genetics, Russia

Freddie Salisbury

Wake Forest University, USA

Board of Editors

Elena Bichenkova

University of Manchester, UK

Thomas E. Cheatham, III

University of Utah, USA

Christo Z. Christov

Northumbria University, Newcastle-upon-Tyne, UK

Miroslav Fojta

Institute of Biophysics, Czech Republic

Yaakov (Koby) Levy

Weizmann Institute of Science, Israel

Luis A. Marky

University of Nebraska Medical Center, USA

Aditya Mittal

IIT-Delhi, India

Remo Rohs

University of Southern California., USA

Wolfram Saenger

Freie Universität Berlin, Germany

Irina M. Russu

Wesleyan University, USA

Zippora Shakked

Weizmann Institute of Science, Israel

R. Sowdhamini

National Center for Biological Sciences, India

Jiri Sponer

Academy of Sciences of the Czech Republic

Vladimir Uversky

Univ. of South Florida, Tampa, USA

Sybren S. Wijmenga

Radboud University Nijmegen, Netherlands

Krystyna Zakrzewska

IBCP, Lyon, France

Dmitry O. Zharkov

SB RAS Institute of Chemical Biology, Russia

Editorial Advisory Board

Manju Bansal

Indian Institute of Science, Bangalore, India

David L. Beveridge

Wesleyan University, USA

Samir K. Brahmachari

CSIR, New Delhi, India

Takashi Gojobori

National Institute of Genetics, Mishima, Japan

Andrew H.-J. Wang

Academia Sinica, Taiwan

David M. J. Lilley FRS

University of Dundee, UK

Martin Karplus

Nobel Laureate,

Harvard University, USA

Olga Kennard FRS

University of Cambridge, UK

Hamilton Smith

Nobel Laureate,

Johns Hopkins University, USA

Edward N. Trifonov

University of Haifa, Israel

Chun-Ting Zhan

Tianjin University, China

Sample our Bioscience Journals



>> **Sign in here** to start your access
to the latest two volumes for 14 days

CHOOSE **open**

ALL LIFE METHODS

SUBMIT NOW



Information for

[Authors](#)

[R&D professionals](#)

[Editors](#)

[Librarians](#)

[Societies](#)

Opportunities

[Reprints and e-prints](#)

[Advertising solutions](#)

[Accelerated publication](#)

[Corporate access solutions](#)

Open access

[Overview](#)

[Open journals](#)

[Open Select](#)

[Dove Medical Press](#)

[F1000Research](#)

Help and information

[Help and contact](#)

[Newsroom](#)

[All journals](#)

[Books](#)

Keep up to date

Register to receive personalised research and resources
by email



Sign me up



Copyright © 2023 Informa UK Limited [Privacy policy](#) [Cookies](#) [Terms & conditions](#)

[Accessibility](#)

Registered in England & Wales No. 3099067
5 Howick Place | London | SW1P 1WG



In silico modeling and empirical study of 4-*n*-Butylresorcinol nanoliposome formulation

Rini Dwiastuti, Muhammad Radifar, Dina Christin Ayuning Putri, Florentinus Dika Octa Riswanto & Maywan Hariono

To cite this article: Rini Dwiastuti, Muhammad Radifar, Dina Christin Ayuning Putri, Florentinus Dika Octa Riswanto & Maywan Hariono (2021): *In silico* modeling and empirical study of 4-*n*-Butylresorcinol nanoliposome formulation, Journal of Biomolecular Structure and Dynamics, DOI: [10.1080/07391102.2021.1946430](https://doi.org/10.1080/07391102.2021.1946430)

To link to this article: <https://doi.org/10.1080/07391102.2021.1946430>



Published online: 08 Jul 2021.



Submit your article to this journal [↗](#)



Article views: 22



View related articles [↗](#)



View Crossmark data [↗](#)



In silico modeling and empirical study of 4-*n*-Butylresorcinol nanoliposome formulation

Rini Dwiastuti^a, Muhammad Radifar^b, Dina Christin Ayuning Putri^a, Florentinus Dika Octa Riswanto^c and Maywan Hariono^c

^aDepartment of Pharmaceutical Technology, Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia; ^bMedical Laboratory Technology, Guna Bangsa Institute of Health Science, Yogyakarta, Indonesia; ^cDepartment of Pharmaceutical Analysis and Medicinal Chemistry, Faculty of Pharmacy, Sanata Dharma University, Yogyakarta, Indonesia

Communicated by Ramaswamy H. Sarma

ABSTRACT

A study to incorporate *in silico* modeling with an empirical experiment has been carried out to formulate nanoliposome containing 4-*n*-butylresorcinol as the active ingredient. The *in silico* modeling was performed using molecular dynamics simulation followed by radius of gyration observation to provide insight into the mechanisms of 4-*n*-butylresorcinol stabilization by liposome due to their nano-size. The empirical experiment was conducted by formulating the nanoliposome using soy lecithin phospholipid formula as suggested by the *in silico* modeling followed by determining its particle size as well as its shape. From their incorporation, it was found that 3200 phospholipid molecules were selected in formulating nanoliposome containing 4-*n*-butylresorcinol. The results of the nanoliposomes size observation in the modeling of 3200 lipid molecules was 87.01 ($\pm$ 0.59) nm, whereas the size from the empirical study was 87.57 ($\pm$ 0.06) nm.

List of abbreviation: LUV: Large Unilamellar Vesicles; SUV: Small Unilamellar Vesicles; AFA: Adaptive Focused AcousticsTM; CGMD: Coarse-Grained Molecular Dynamic; DMPC: dimyristoyl-phosphatidylcholine; DPPC: dipalmitoyl-phosphatidylcholine; TEM: Transmission Electron Microscopy; PIPC: palmitoyl-1-nonyl-phosphatidylcholine; vdW: van der Waals

ARTICLE HISTORY

Received 5 February 2021
Accepted 17 June 2021

KEYWORDS

nanoliposome; 4-*n*-butylresorcinol; formulation; *in silico* modeling; simulation

Introduction

Nanoparticle technology has become one of the trending interests in drug delivery system research in the last decade (Boisseau & Loubaton, 2011; Puri et al., 2009). Applied technology in nanoparticle formulation can be performed using two different methods, namely top-down and bottom-up methods (Singh et al., 2011; Reverchon & Adami, 2006). Drug delivery system improvement using lipid nanoparticle formulation has been developed by arranging phospholipid bilayer fragments as well as liposomes for a diverse administration route in the human body. Lipid nanoparticle formulation provides several advantages such as low toxicity, which is proven by *in vivo* observation and its capability to improve the physical stability of the active ingredient in the dosage form formulation (Akbarzadeh et al., 2013). Nanoparticle technology can be applied in the drug formulation with the encapsulation process by several matrices such as nanospheres, nanoliposomes, and nanoemulsions (Martien et al., 2012).

One of the lipid nanoparticle drug delivery systems that has been commonly developed is nanoliposome since it has been reported to have several advantages such as increasing the efficacy and therapeutic index as well as improving the drug stability through the encapsulation system (Akbarzadeh

et al., 2013). Liposome, a topical dosage form, has been developed in the drug formulation due to its good penetration into the skin (El Maghraby et al., 2010; El Maghraby et al., 2010). Nanoliposome showed good activity of penetrating the cell wall and intercellular space due to their nano-size, i.e. < 200 nm (Reverchon & Adami, 2006). The trend of nanoliposome formulation development increased continuously since the formulation process was reported to play an essential role in drug encapsulation (Wang et al., 2011; Zhao et al., 2009). The physical stability of the dosage form could be maintained by increasing the drug encapsulation and reducing the liposome leakage during the storage (Eloy et al., 2014; Laouini et al., 2012).

Nevertheless, several formulation problems were found during the formulation of nanoliposomes such as the time-consuming process, organic solvent selection consideration, and economical consideration for performing trial and error formulation. Computational methods have been used to predict drug delivery systems before drug formulations. It aims to minimize investment in drug design and development (Li & Hou, 2010). The modeling of liposome formation molecules with phospholipids 1,2- dipalmitoyl-sn-glycero-3-phosphatidylcholine had been carried out by Koshiyama and Wada (2016), showing that the formed liposomes have different

shapes and sizes depending on the number of phospholipids being used. A study related to *in silico* modeling of liposome formation had also been carried out by Dwiaastuti et al. (2018) using 1,2-dilauroyl-sn-glycero-3-phosphatidylethanolamine producing almost the same shape and size of the liposomes in empirical experiments. The difference between the two studies is the software being used in the *in silico* modeling simulation. However, it has not been supported by empirical verification of the liposome manufacture. All of these problems can be overcome by generating an accurate prediction, taking into account the atomic interaction between the drug and its delivery system using appropriate molecular modeling simulation (Hashemzadeh et al., 2020). Furthermore, the number of water and lipid molecules obtained from the *in silico* modeling is correlated with the empirical observations of the liposome formation. The conversion factor should be calculated to obtain a quantitative correlation of phospholipid numbers used both for simulation modeling and empirical observation (Jämbeck et al., 2014). The *in silico* modeling of liposome formation can be applied in a formulation study by employing mathematical prediction to achieve a good nanoliposomes drug delivery system supported by the computational approach (Dogra et al., 2019; Cern et al., 2017; Cern et al., 2014).

One of the liposome formulation methods is a lipid dispersion in an aqueous media with sonication along with controlling its temperature, which is the easiest method (Akbarzadeh et al., 2013). The previous study by Shen et al. (2015) had empirically approached the formulation of Large Unilamellar Vesicles (LUV) or Small Unilamellar Vesicles (SUV) liposomes using focused ultrasonic irradiation with the usage of Adaptive Focused Acoustics™ (AFA) (Shen et al., 2015). This applied method was performed at a low temperature without organic solvent and produced liposomes with a size of < 600 nm. One way to reduce trial and error in experimental study is by using *in silico* modeling to predict the formation of liposomes sub-molecularly (Trisilowati and Mallet, 2012). The lipid dispersion in aqueous media with sonication energy and its controlled temperature can be used as the system in the modeling approach, which is the first step before starting the empirical study. Furthermore, the *in silico* modeling study should be compared with the empirical observations to obtain a correlation between the number of phospholipid molecules resulted from the modeling with the number of phospholipid molecules in the empirical study.

In silico modeling in this study was described as a computational approach to study the interaction between drug molecules with the liposome phospholipid membrane and to present the whole molecular interaction during the liposome formation. Risselada and Marrink (2009) studied the effect of temperature and membrane composition during the simulation process towards the structure and dynamic behavior of the liposome membrane with the diameter of 15-20 nm using Coarse-Grained Molecular Dynamic (CGMD) simulation (Risselada & Marrink, 2009). Another study by Siwko et al. (Siwko et al., 2009) was conducted on *in silico* modeling of dimyristoyl-phosphatidylcholine (DMPC) phospholipid which used CGMD at 323 K to investigate the interaction between resorcinol and the bilayer

lipid membrane. The results showed that the presence of resorcinol interestingly increased the membrane stability of the lipid bilayer systems. Hudiyanti et al. (Hudiyanti et al., 2014) also employed the CGMD for simulating the *in silico* modeling to prepare liposomes from various phospholipids. In this study, three types of liposome (origin, deformed liposome, and planar bilayer) were *in silico* simulated to interact with water, to give an insight understanding on which type of liposome will give the most stable structure during 160 ns of the simulation time. Therefore, the use of CGMD in drug delivery system modeling has been commonly used to provide the best formula in the liposome production.

In this present study, 4-*n*-butylresorcinol was used as the drug instead of resorcinol as the drug model. It functions as tyrosinase inhibitor for the management of pigmentation disorders, such as the topical treatment of hyperpigmentation (Kolbe et al., 2013). This resorcinol derivative is more stable than resorcinol due to the oxidation process (Love et al., 2005). A double-blind study for melasma with 4-*n*-butylresorcinol being applied twice daily within 8 weeks was necessary to achieve the hypopigmenting effect. However, 12 adverse events including mild erythema, dryness, peeling, and desquamation have appeared (Khemis et al., 2007). The 4-*n*-butylresorcinol was encapsulated in liposomes to improve the stability and to reduce skin irritation through hydration of the epidermis (Huh et al., 2010; De Leeuw et al., 2009). The stability should be improved by encapsulating the compound into the complex of nanoliposome structure. Encapsulating phenol-related compounds such as 4-*n*-butylresorcinol aims to preserve the biological activity and to improve the stability of the active compounds, as well as to ensure controlled release of the latter. Encapsulation has the advantage of being a non-thermal stabilization approach and is suitable for temperature-sensitive natural biologically active compounds. The wall material usually improves the stability of the active compounds by protecting them from direct exposure to air and light (Coradini et al., 2014; Love et al., 2005). The *in silico* modeling simulation in the initial step of the formulation was important to predict the physical properties of the lipid nanoparticle complex production. The formulation technique was necessarily developed to produce nanoliposomes with nano size (<100 nm) in a short time usage of sonication. It was reported that the heating and sonication processes were important in the nanoliposomes formulation. Mozafari et al. reported that the production of nanoliposomes with the size of ± 600 nm was successfully carried out by the heating process without any extrusion process in polycarbonate membrane as well as sonication (Mozafari et al., 2007). Another study reported that the particle size reduction (≤ 100 nm) occurred with an increase of sonication time for more than 21 min (Silva et al., 2010). Hence, it is important to consider the correlation between the *in silico* modeling approach and empirical observational results in an integrated research. This study aims to correlate Molecular Dynamic (MD) simulation and an empirical observation in the production of nanoliposomes using soy lecithin phospholipid and 4-*n*-butylresorcinol as the delivery system and active ingredient, respectively.

Table 1. The nanoliposome formula on the empirical observation.

Materials	Amount
Soy lecithin	7.74 g
4- <i>n</i> -butylresorcinol	0.1% w/v
Re-distilled water	100.00 mL

Materials and methods

Materials

Soy lecithin phospholipid (Pharmaceutical Grade, Nacalai Tesque, Japan), redistilled water, 4-*n*-butylresorcinol (Pharmaceutical Grade, SHREEJI Pharma International, India), ethanol (Analytical Grade, Merck, USA).

Gas chromatography-mass spectrometry analysis (GC-MS) of soy lecithin

The GC-MS analysis was performed using a Shimadzu GCMS-QP2010S with Agilent HP-5ms column (30 m x 0.25 mm x 0.25 μ m). The injector temperature was set to 310 °C. Helium was used as carrier gas at a constant flow rate of 0.40 mL/min. The column temperature was kept at 120 °C for 5 min and then increased from 120° to 300 °C at 5° C/min. As for the ion source, the temperature was set to 250° C, and the interface temperature was set to 305° C. The mass scanning was set from m/z 28 to 600.

Molecular dynamics simulation of soy lecithin

Coarse-grained Molecular Dynamics using MARTINI 2.0 force field was used to study the liposome formation of soy lecithin. Prior to the initial coordinate preparation, the coarse-grained structure of phospholipids (DPPC, POPC, and PIPC), cholesterol, and 4-*n*-butylresorcinol were prepared. The GC-MS analytical results were used to determine the composition of soy lecithin and to construct the initial coordinates according to the composition. The initial coordinate was prepared by a random placement of phospholipid and cholesterol in a $22 \times 22 \times 22 \text{ nm}^3$ box using the gmx insert-molecule module of GROMACS 2019.1 package on Centos 7.4. Three concentrations of lipids were used i.e. 2400, 2800, and 3200 lipids/boxes. Each concentration of lipid had two treatments, i.e. by adding 4-*n*-butylresorcinol and without 4-*n*-butylresorcinol (negative controls). The molecule numbers of 4-*n*-butylresorcinol being added were 123, 143, and 164 molecules for 2400, 2800, and 3200 lipids, respectively. In total, there are six different initial coordinates in a duplicate experiment. After the lipid was randomly placed, the box configuration was edited using a gmx edit conf module and solvated with 78,299 water beads using a gmx solvate module generating a system consisted of lipids and water with the size of $24 \times 24 \times 24 \text{ nm}^3$. The initial coordinates were then subjected to energy minimization using the gmx mdrun module with the steepest descent method. The resulting minimized system was then simulated under NPT condition for 240 ns using 40 fs time step, 323 K temperature, and reaction-field long-range electrostatic treatment. The

temperature was controlled using the V-rescale method while the pressure was controlled using the Berendsen barostat. The MD results were then processed with gmx energy module to generate the potential energy data and gmx gyrate module to generate the radius of gyration data. The data were then plotted using Grace. The trajectory file was visualized using VMD 1.9.3 to understand the dynamic of the liposome assembly and the morphology of the liposome.

Empirical observations

The formulation of nanoliposomes with and without 4-*n*-butylresorcinol was carried out by dispersing an amount of soy lecithin over re-distilled water using the heating method and combined with sonication. The formula was composed of soy lecithin and other ingredients, as presented in Table 1. The empirical observation was approached by formulating the nanoliposomes using the theoretical data of soy lecithin through *in silico* modeling of 3200 phospholipid molecules. The phospholipid being used in this study was soy lecithin having the temperature transition between 50-60 °C (De Leeuw et al., 2009). The temperature in the formulation was set at 50 °C along with 30 min sonication and 37 kHz of a bath sonicator. Soy lecithin was dispersed in 100 ml of re-distilled water at 50 °C until the nanoparticle system met the requirement of a polydispersity index (< 0.3). The 50° C temperature used during the formulation was conducted to set the temperature condition close to the *in silico* modeling parameters. The soy lecithin dispersed solution was then blended at a high speed. This aimed to minimize the bilayer fragment sheets that had been formed. The solution was then homogenized using Ultraturac® for 1 min at four scales and sonicated for 30 min (Dwiastuti et al., 2016; Dwiastuti et al., 2018) until the particle with <100 nm in size were observed. The product was then left to cool down at room temperature before morphological and particle size testing were conducted. After that, the product was stored in a refrigerator at 4-8 °C.

The conversion of soy lecithin from molecular modeling into empirical observation

The soy lecithin used in the empirical observation was based on the number of phospholipid molecules in nanoliposomes modeling. The concentration of phospholipid in the simulation of 3200 phospholipid molecules was 387.12 mg/mL, which equals to 0.38712 g/mL.

The amount of soy lecithin used

$$\begin{aligned}
 &: \frac{0.38712 \text{ g/mL} \times (\text{the box size in modeling})}{100 \text{ nm}} \\
 &: 0.38712 \text{ g/mL} \times \frac{(20 \text{ nm})}{100 \text{ nm}} \\
 &: 0.077424 \text{ g/mL} \sim 7.74 \text{ g/100 mL}
 \end{aligned}$$

According to the conversion above, the soy lecithin used was 7.74 g in 100 mL of re-distilled water.

In this study, the expected empirical size of nanoliposomes was 100 nm. However, due to the limitation in facility (computer speed and power), the simulation of

nanoliposomes formation with particle size of 100 nm was difficult to conduct in the *in silico* modeling. Hence, we converted the particle size into 20 nm which is equivalent to 0.077424 g/mL (7.74 g/100 mL) of soy lecithin where the simulation could be performed.

Determination of the particle size

The determination and distribution process in the system were conducted using particle size analyzer instrument, Horiba SZ-100, based on the scattering dynamic light principle with 25.0 °C temperature and 90° angle. 0.50 µL of the sample was transferred into a 25 mL volumetric flask, and then re-distilled water was added up to the calibration sign. A volume of 2.00 mL of aliquot was transferred into cuvet for the measurement of 100,000 particles.

Determination of a lamellar particle

The lamellar particle was observed using Transmission Electron Microscopy (TEM) JEM-1400 (Faculty of Natural Sciences, Gadjah Mada University, Indonesia). The determination of the lamellar particle was carried out on lipid nanoparticle with and without the presence of 4-*n*-butylresorcinol. A volume of 0.5 mL of the formula was added into 1 mL of re-distilled water, and then dropped into an object plate. The electromagnetic transmission light was subjected to the plate, and then the particle morphology was observed at a suitable magnitude. Morphological results of the sample were displayed on the camera screen with no staining process. The results were then visualized as the lamellar particle shape.

Results and discussion

The GC-MS of soy lecithin

GC-MS is used as this technique is suitable for lipid identification, in which the lipids are usually presented in an ester form (Naviglio et al., 2017). Ester is usually volatile; therefore, GC-MS is one of the suitable techniques to identify lipid both for its qualitative as well as quantitative prediction (Chiu & Kuo, 2020). Furthermore, GC-MS is a fast and cheap method, providing the compounds library which could easily predict the identified peak in GC chromatogram which corresponds to the mass spectrum when it is available in the database. The % similarity in the MS results enhances the power of this instrument to be utilized in determining the lipid. The ion source is electron impact 70 MeV resulting in fragmentation of the parent molecule into its most stable peak (base peak) accordingly. However, this technique has a disadvantage as the thermal stability of the compounds becomes the major issue (Koo et al., 2013). LC-MS could be the alternative method to reduce this thermal instability. Unfortunately, we could not perform this method at the moment due to our limitation. The soy lecithin contents were analyzed using GC-MS method. The retention time,

percentage of peak area, and the base peaks as well as their interpretation are presented in Table 2.

In general, it was found that there were two primary compounds in soy lecithin, namely fatty acid and sterol (Table 3). The detected fatty acids were recognized as palmitic acid, oleic acid, and linoleic acid; whereas the sterol contents were recognized as clonasterol, 5-ergosterol, and stigmasterol.


Table 3 shows that only several lipids were detected and calculated by the GC-MS method due to their thermal instability during the analytical process. Therefore, the percentage of total area calculation was still below 100%. It can be seen that palmitate content was set at 54.32% since palmitate was found as a major compound in soy lecithin. Finally, the number of molecules for each lipid can be determined by multiplying the percentage and the total lipid, which can be seen in Table 4.

The potential energy calculation

The potential energy in the modeling of 2400, 2800, and 3200 phospholipid molecules were calculated. It can be noted that by increasing 4-*n*-butylresorcinol, the potential energy decreased (Table 5). The temperature was set to 50 °C during the simulation due to the protocol published in the previous study by Siwko et al. (2009). Figure 1 depicted the graphical plot of potential energy-time dependent which occurred during the simulation of 2400, 2800, and 3200 lipid molecules. The energy calculation involved in this study during the simulation processes were kinetic and potential energies.

The potential energy involved in the liposome formation describes that without 4-*n*-butylresorcinol (blue lines) (Figure 1), it tends to have higher potential energy than with 4-*n*-butylresorcinol (green lines). This indicates that a system having 4-*n*-butylresorcinol is more stable than the one without this active ingredient. The results are in line with the study performed by Siwko et al., in which the presence of resorcinol increased the liposome structural stability by joining the bilayer membrane while inducing a proper arrangement of the phospholipid acyl chain leading to a lipid head group hydration.

The potential energies employed during the liposome formation are electrostatic and vdW energies, which are visualized in Figure 2. Figure 2a indicates that the electrostatic energy declines along with the increasing number of the phospholipid molecules as well as the 4-*n*-butylresorcinol, and so does the vdW energy in Figure 2b. The total energy system in Figure 2d is the cumulative energy of kinetics (Figure 2c) and the potential energy of electrostatic (Figure 2a). From these graphs, the electrostatic and vdW are two of the most potential energies involved during the nanoliposome modeling. Although the electrostatic seems to contribute the most stable energy with the system having the largest molecule number of phospholipids, the presence of 4-*n*-butylresorcinol is predicted to increase the system stability by inducing the phospholipid acyl chain to stabilize the forming bilayer membrane. The vdW also contributes into the system stability as this energy profile (Figure 2b) is identical with the electrostatic energy (Figure 2a). In conjunction,

Table 2. GC-MS analysis results of soy lecithin.

No	Peak #	Retention Time (minute)	Area (%)	Base peak (MW)	Interpreted as (MW)
1	2	24.38	1.49	74.05	Methyl palmitate (270)
2	5	25.73	5.42	43.10	Palmitic acid (256)
3	6	28.15	1.50	55.05	Methyl oleate (296)
4	9	29.35	11.17	55.10	Linoleic acid (280)
5	13	35.58	5.25	43.05	Glycerol 1-palmitate (330)
6	16	38.52	14.09	55.10	2-hydroxy-1-(hydroxymethyl)ethyl ester linolein (354)
7	18	45.28	1.89	43.10	Cholesteryl myristate
8	19	48.13	4.85	43.05	5-ergosterol (400)
9	20	48.90	3.98	55.10	Stigmasterol (412)
10	21	50.49	10.09	43.05	Clonasterol (414)

Note: only lipids and positive compound are shown.

Table 3. Soy lecithin primary component profiles.

Component	Peak #	% Area	Total % Area
Fatty acid			
Palmitate	2, 5, 13	1.49 + 5.42 + 5.25	12.16
Oleate	6	1.50	1.50
Linoleate	9, 16	11.17 + 14.09	25.26
		Total fatty acids	38.92
Sterol			
5-ergosterol	19	4.85	4.85
Stigmasterol	20	3.98	3.98
Clonasterol	21	10.09	10.09
		Total sterol	18.92

the total energy (Figure 2d) is in proportion with the potential energy as depicted in Figure 2a and b, whereas the kinetic energy does not have the same trend. This different trend could be due to the kinetic energy being a fraction from the total energy system, so this could be neglected. In other words, the selected system is considered from the most stable energy profile, in which potential system employing electrostatic and vdW are the two best ones.

The radius of gyration of in silico modeling simulation

The radius of gyration can be described as a unit for the radius average from the molecule to the centre of mass to determine the liposome diameter. The radius of gyration allows to measure the nanoliposome formation during the simulation of 2400, 2800, and 3200 lipid molecules with and without the presence of 4-*n*-butylresorcinol (Figure 3). The nanoliposomes morphological results of the simulation of 2400, 2800, and 3200 lipid molecules and their snapshots of nanoliposomes formation results during simulation can be

Table 4. Number of lipid molecules for each system in molecular modeling simulation.

System #	Number of lipids	DPPC	POPC	PIPC	Cholesterol
1	2400	1304	36	606	454
2	2800	1521	42	707	530
3	3200	1738	48	808	606

Note. DPPC = dipalmitoyl-phosphatidylcholine; POPC = palmitoyloleoyl-phosphatidylcholine; PIPC = palmitoyllinoleoyl-phosphatidylcholine.

seen in Figures 4 and 5, respectively. The liposome radii being formed during the simulation are shown in Figure 3a–c with a duplicate simulation. The results indicate that the liposome radii increases along with the increasing number of both phospholipid and 4-*n*-butylresorcinol molecules.

The increasing liposome radii results in the increasing of the liposome size as its morphology is depicted in Figure 4. The shape of simulation having 2400 phospholipid molecules with 4-*n*-butylresorcinol in the first replication is more likely less spherical than the second one (Figure 4a). The similar results occur in the system where 2800 phospholipid molecules with 4-*n*-butylresorcinol show less imperfect spherical shape. Interestingly, the system with 3200 phospholipid molecules with 4-*n*-butylresorcinol shows the most spherical shape among others, indicating its efficient atomic interactions between phospholipid and 4-*n*-butylresorcinol. The inefficient atomic interactions could form the double connection between 4-*n*-butylresorcinol and its corresponding phospholipid and other phospholipid leading to the final imperfect spherical structure. From this simulation, the system with 3200 phospholipid molecules is the most stable and reliable system predicted to produce liposome structure that will be used in the empirical study.

Table 5. The comparison between the time, the potential energy, and the shape of liposome being formed during the simulation with the number of lipids and 4-*n*-butylresorcinol treatment variation.



Number of lipids	<i>n</i> -butyl resorcinol	T enclosed (ns)	E1 ($\times 10^6$ kJ/mol)	E2 ($\times 10^6$ kJ/mol)	E3 ($\times 10^6$ kJ/mol)	Shape
2400	No	90	-2.947	-2.962	-2.963	Rod
		19	-2.942	-2.948	-2.952	Spherical
	Yes	56	-2.954	-2.968	-2.969	Elliptical
		31	-2.947	-2.953	-2.957	Spherical
2800	No	40	-2.941	-2.949	-2.951	Spherical with torus-shaped cavity
		20	-2.948	-2.956	-2.959	Spherical
	Yes	68	-2.947	-2.957	-2.959	Elliptical with U-shaped cavity
		18	-2.950	-2.955	-2.958	Spherical
3200	No	12	-2.937	-2.939	-2.945	Spherical
		24	-2.939	-2.941	-2.945	Spherical with crescent-shaped cavity
	Yes	32	-2.945	-2.953	-2.955	Spherical with torus-shaped cavity
		14	-2.950	-2.954	-2.958	Spherical

T = The time until liposome is fully enclosed.

E1 = The energy at 10 ns.

E2 = The energy when liposome is fully enclosed.

E3 = The energy at 240 ns.

**Figure 1.** The graph plotting potential energy-time which occurred during the simulation of a) 2400, b) 2800, and c) 3200 lipid molecules. Note: Simulation with and without 4-*n*-butylresorcinol are shown by two green lines and two blue lines, respectively, as each treatment was performed in duplicate.**Figure 2.** The graph plotting potential energy-time which occurred during the simulation of a) electrostatic, b) vdW, c) kinetic, and d) total energy.




Figure 3. The gyration radius of the nanoliposomes being formed during the simulation of a) 2400, b) 2800, and c) 3200 of lipid molecules with (green) and without (blue) the presence of 4-*n*-butylresorcinol.




Figure 4. Nanoliposomes morphological results of the simulation of a) 2400, b) 2800, and c) 3200 lipid molecules with two replications for each lipid molecule number.

The size of nanoliposome with the presence of 4-*n*-Butylresorcinol

The size of nanoliposome resulted from both molecular dynamics simulations and empirical observations were compared and presented in Table 6. In this presentation, the empirical liposome was prepared from the selected system only (3200 phospholipid molecules with 4-*n*-butylresorcinol) because this system showed the most stable energy profile as well as its morphology. The distribution curve of the nanoliposome size can be seen in Figure 6, demonstrating the population of liposome particle size with three times replications which shows the consecutive mode values i.e. 87.60, 87.50, and 87.60 nm with the standard deviation value of 0.06. The mode value is chosen to represent the liposome size distribution because it represents the largest frequency of the liposome size, which is generated in the empirical study. The distribution curves exhibit a similar pattern so that it indicates a good repeatability in the particle size of the liposome.

The morphology of 3200 phospholipid molecules

The morphological observation of 3200 phospholipid molecules from the simulation model was compared to the empirical observation without the presence of 4-*n*-butylresorcinol,

which can be seen in Figure 7. In addition, the interaction between drug and the membrane model is depicted in Figure 8.

The GC-MS method was utilized to determine the chemical composition of the soy lecithin as listed in Table 2. The fatty acids of the phospholipid and sterol were used as the platform in the nanoliposomes modeling. The identified compositions of fatty acid from the GC-MS were palmitic acid, oleic acid, and linoleic acid. Theoretically, it is appropriate with the composition of soy lecithin (Perkins, 1995; Van Hoogevest et al., 2013). The sterol being composed in the soy lecithin was 5-ergosterol, stigmasterol, and clionasterol (Table 3). The percentage of fatty acid and sterol as calculated in Table 3 was used as the basic calculation to decide the composition of the nanoliposomes modeling.

The nanoliposomes modeling with and without the presence of 4-*n*-butylresorcinol used three variations of the lipid molecule number, i.e. 2400, 2800, and 3200 molecules. This observation was aimed to predict the possibility of the nanoliposomes formation with and without the presence of 4-*n*-butylresorcinol by modeling on a few lipid molecules used. The decision of selecting the number of lipid molecules was applied and referred to the study conducted by Koshiyama and Wada (2016). The results indicated a formation of nanoliposomes from modeling using 2400, 2800 and 3200 molecules inside a 22 nm³ size box (Figure 2).




Figure 5. Snapshots of nanoliposomes formation results during the simulation.

Table 6. Results of nanoliposome size based on simulation and empirical observation for 4-*n*-butylresorcinol.

Nanoliposome sizes based on the <i>in silico</i> simulations (3200 molecules)		Nanoliposome sizes based on the empirical observation		
Replication	Nanoliposome sizes (nm)	Replication	Calculation mode (nm)	Polydispersity Index
1	86.60	1	87.60	0.28
2	87.43	2	87.50	0.18
		3	87.60	0.24
Mean ($\pm$ SD)	87.01 ($\pm$ 0.59)	Mean ($\pm$ SD)	87.57 ($\pm$ 0.06)	

According to the graphical energy presented in Figures 1 and 2, the potential energies were stronger for a lower lipid number. This trend appears in the lower number of lipids and the higher number of water, which contributes to stronger electrostatic and vdW interactions. In terms of the velocity and stability, both the energy graph as well as the radius of gyration graph show that at a higher number of lipids, the liposomes are formed faster and quickly stabilized. It is also important to note that the slower liposome formation tends to give the phospholipid bilayer a chance to form a multiple connection, leading to an imperfect sphere or a bilayer bridge in the middle of the liposome (see Figure 4b). Therefore, the 3200 lipids molecule was subjected to the empirical nanoliposomes formulation with and without the

presence of 4-*n*-butylresorcinol. The formation of nanoliposomes in the molecular modeling involves kinetic and potential energies in the form of electrostatic and vdW interactions. According to the energy calculation results in the modeling simulation at 50 °C, the nanoliposome system was stable at about 100 ns (Figures 1 and 2). Nanoliposome structures were generated in the modeling simulation with electrostatic energy. It may result from the interaction between electronegative atoms in the phosphate group of the phospholipid with the hydrogen atom from the water molecules around the system. Furthermore, an electrostatic interaction between polar groups of phospholipid during simulation also occurs. The lower vdW energy indicates a similar graphical profile with lower electrostatic and total




Figure 6. The nanoliposome size distribution curves in triplicate measurements exhibit a similar population with their consecutive mode values i.e. 87.60 (a), 87.50 (b), and 87.60 nm (c) having polydispersity index (PI) of 0.281, 0.179, and 0.237, respectively.




Figure 7. The nanoliposome morphology being formed from a) the modeling and b) the empirical observation on 3200 lipid molecules.

energy in the system. The interaction between non-polar group during the simulation resulted in vdW energy. With the longer simulation time, the electrostatic, vdW, and the total energy were decreased due to the system stabilization in order to achieve the nanoliposomes structure (Figure 5) for simulation, with and without 4-*n*-butylresorcinol in three different molecules number (2400, 2800, and 3200). The system stabilization has been achieved when the unilamellar nanoliposomes was formed in 100 ns. Moreover, the electrostatic, vdW, and total energies decreased in both systems with and without 4-*n*-butylresorcinol during the simulation. The vdW energy depicted interactions between non-polar

groups of phospholipid molecules and non-polar group of 4-*n*-butylresorcinol. Chemical properties of the non-polar model compound might affect the vdW interaction at the initial and at the end of the simulation. The interaction between 4-*n*-butylresorcinol-lipid and lipid-membrane during the simulation process resulted in similar characteristic results with the previous study (Siwko et al., 2009), where the hydroxyl (-OH) group of 4-*n*-butylresorcinol interacted with the hydrophilic group of the phospholipid compound (Figure 7). The presence of 4-*n*-butylresorcinol increased the vdW interaction between non-polar groups, whereas the electrostatic interaction in phosphate group with hydrogen




Figure 8. The interaction of the model compound in the lipid membrane.

atom from water was decreased since there was several hydrogen bonding between the two phospholipid molecules. The morphological results of the simulation (Figure 5) indicated that nanoliposomes were formed into unilamellar structure. The simulation with 3200 phospholipid molecules resulted in a similar form of nanoliposomes compared to the empirical observation results (Figure 6).

The conversion of the phospholipid from the simulation was done using a concentration of 3200 molecules by 0.38712 g/mL with the conversion factor of 0.2. This factor was a ratio between the box size used in the simulation (20 nm) and the actual box that should be used in the empirical observation, i.e. 100 nm. The parameter being observed in the nanoliposomes product was the size and the morphology of the nanoliposomes being formed. The nanoliposomes size in the molecular modeling of 3200 lipid molecules was 87.01 nm (± 0.59), whereas the size from the empirical study was 87.57 (± 0.06). These results indicated that there was no significant difference for the size and the morphology of nanoliposomes formed with and without the presence of 4-*n*-butylresorcinol. However, this study has some limitations in that no physicochemical parameter evaluation and empirical studies for 2400 and 2800 phospholipid molecules are conducted, which prompts us to do these experiments in our future study.

Conclusions

Molecular dynamics simulation was proven to be the tool that strongly predicts the dynamics behavior of 4-*n*-butylresorcinol upon nanoliposome formulation. The program was able to predict the number of phospholipid molecule during the formulation which resulted in agreement between *in silico* and empirical study. It was found that phospholipid molecules with the number of 3200 were selected in the formulation process of nanoliposome containing 4-*n*-butylresorcinol. The modeling and empirical studies resulted in the molecules size of 87.01 (± 0.59) and 87.57 (± 0.06) nm, respectively. These significant results indicated the connection between the *in silico* modeling and the empirical observation in formulating nanoliposome.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

We gratefully acknowledge The Ministry of Research and Technology of Higher Education for funding under the Research Grant with contract letter no: 029/Penel./LPPM-USD/IV/2019.

References

- Akbarzadeh, A., Rezaei-Sadabady, R., Davaran, S., Joo, S. W., Zarghami, N., Hanifehpour, Y., Samiei, M., Kouhi, M., & Nejati-Koshki, K. (2013). Liposome: Classification, preparation, and applications. *Nanoscale Research Letters*, 8(1), 1–9. <https://doi.org/10.1186/1556-276X-8-102>
- Boisseau, P., & Loubaton, B. (2011). Nanomedicine, nanotechnology in medicine. *Comptes Rendus Physique*, 12(7), 620–636. <https://doi.org/10.1016/j.crhy.2011.06.001>
- Cern, A., Barenholz, Y., Tropsha, A., & Goldblum, A. (2014). Computer-aided design of liposomal drugs: In silico prediction and experimental validation of drug candidates for liposomal remote loading. *Journal of Controlled Release: official Journal of the Controlled Release Society*, 173, 125–131. <https://doi.org/10.1016/j.jconrel.2013.10.029>
- Cern, A., Marcus, D., Tropsha, A., Barenholz, Y., & Goldblum, A. (2017). New drug candidates for liposomal delivery identified by computer modeling of liposomes' remote loading and leakage. *Journal of Controlled Release: Official Journal of the Controlled Release Society*, 252, 18–27. <https://doi.org/10.1016/j.jconrel.2017.02.015>
- Chiu, H. H., & Kuo, C. H. (2020). Gas chromatography-mass spectrometry-based analytical strategies for fatty acid analysis in biological samples. *Journal of Food and Drug Analysis*, 28(1), 60–73. <https://doi.org/10.1016/j.jfda.2019.10.003>
- Coradini, K., Lima, F. O., Oliveira, C. M., Chaves, P. S., Athayde, M. L., Carvalho, L. M., & Beck, R. (2014). Co-encapsulation of resveratrol and curcumin in lipid-core nanocapsules improves their in vitro antioxidant effects. *European Journal of Pharmaceutics and Biopharmaceutics*, 88(1), 178–185. <https://doi.org/10.1016/j.ejpb.2014.04.009>
- De Leeuw, J., De Vrijlder, H. C., Bjerring, P., & Neumann, H. A. M. (2009). Liposomes in dermatology today. *Journal of the European Academy of Dermatology and Venereology: JEADV*, 23(5), 505–516. <https://doi.org/10.1111/j.1468-3083.2009.03100.x>
- Dogra, P., Butner, J. D., Chuang, Y. L., Caserta, S., Goel, S., Brinker, C. J., Cristini, V., & Wang, Z. (2019). Mathematical modeling in cancer nanomedicine: A review. *Biomedical Microdevices*, 21(2), 1–23. <https://doi.org/10.1007/s10544-019-0380-2>

- Dwiastuti, R., Noegrohati, S., Istyastono, E. P. Marchaban, (2016). Formulation and physical properties observations of soy lecithin liposome containing 4-n-butylresorcinol. In *AIP Conference Proceedings* (Vol. 1755, No. 1, p. 160005). AIP Publishing LLC.
- Dwiastuti, R., Radifar, M., Marchaban, M., Noegrohati, S., & Istyastono, E. P. (2018). Molecular dynamics simulations and empirical observations on soy lecithin liposome preparation. *Indonesian Journal of Chemistry*, 16(2), 222–228. <https://doi.org/10.22146/ijc.21167>
- Eloy, J. O., de Souza, M. C., Petrilli, R., Barcellos, J. P. A., Lee, R. J., & Marchetti, J. M. (2014). Liposomes as carriers of hydrophilic small molecule drugs: Strategies to enhance encapsulation and delivery. *Colloids and Surfaces. B, Biointerfaces*, 123, 345–363. <https://doi.org/10.1016/j.colsurfb.2014.09.029>
- Hashemzadeh, H., Javadi, H., & Darvishi, M. H. (2020). Study of Structural stability and formation mechanisms in DSPC and DPSM liposomes: A coarse-grained molecular dynamics simulation. *Scientific Reports*, 10(1), 1837. <https://doi.org/10.1038/s41598-020-58730-z>
- Hudiyanti, D., Radifar, M., Raharjo, T. J., Narsito, N., & Noegrohati, S. (2014). A coarse-grained molecular dynamics simulation using NAMD package to reveal aggregation profile of phospholipids self-assembly in water. *Journal of Chemistry*, 2014, 1–6. <https://doi.org/10.1155/2014/273084>
- Huh, S. Y., Shin, J. W., Na, J. I., Huh, C. H., Youn, S. W., & Park, K. C. (2010). Efficacy and safety of liposome-encapsulated 4-n-butylresorcinol 0.1% cream for the treatment of melasma: a randomized controlled split-face trial. *The Journal of Dermatology*, 37(4), 311–315. <https://doi.org/10.1111/j.1346-8138.2010.00787.x>
- Jämbeck, J. P. M., Eriksson, E. S. E., Laaksonen, A., Lyubartsev, A. P., & Eriksson, L. A. (2014). Molecular dynamics studies of liposomes as carriers for photosensitizing drugs: development, validation, and simulations with a coarse-grained model. *Journal of Chemical Theory and Computation*, 10(1), 5–13. <https://doi.org/10.1021/ct400466m>
- Khemis, A., Kaifa, A., Queille-Roussel, C., Duteil, L., & Ortonne, J. P. (2007). Evaluation of efficacy and safety of rucinol serum in patients with melasma: A randomized controlled trial. *British Journal of Dermatology*, 156(5), 997–1004. <https://doi.org/10.1111/j.1365-2133.2007.07814.x>
- Kolbe, L., Mann, T., Gerwat, W., Batzer, J., Ahlheit, S., Scherner, C., Wenck, H., & Stäb, F. (2013). 4-n-butylresorcinol, a highly effective tyrosinase inhibitor for the topical treatment of hyperpigmentation. *Journal of the European Academy of Dermatology and Venerology*, 27, 19–23. <https://doi.org/10.1111/jdv.12051>
- Koo, I., Kim, S., & Zhang, X. (2013). Comparative analysis of mass spectral matching-based compound identification in gas chromatography-mass spectrometry. *Journal of Chromatography. A*, 1298, 132–138. <https://doi.org/10.1016/j.chroma.2013.05.021>
- Koshiyama, K., & Wada, S. (2016). Collapse of a lipid-coated nanobubble and subsequent liposome formation. *Scientific Reports*, 6(1), 1–8. <https://doi.org/10.1038/srep28164>
- Laouini, A., Jaafar-Maalej, C., Limayem-Blouza, I., Sfar, S., Charcosset, C., & Fessi, H. (2012). Preparation, characterization and applications of liposomes: State of the art. *Journal of Colloid Science and Biotechnology*, 1(2), 147–168. <https://doi.org/10.1166/jcsb.2012.1020>
- Li, Y., & Hou, T. (2010). Computational simulation of drug delivery at molecular level. *Current Medicinal Chemistry*, 17(36), 4482–4491. <https://doi.org/10.2174/092986710794182935>
- Love, A. R., Kerschner, J. L., Barratt, M. J., & Zhou, Y. (2005). U.S. Patent No. 6863897. U.S. Patent and Trademark Office.
- Maghraby, G. M. E., Williams, A. C., & Barry, B. W. (2010). Skin delivery of 5-fluorouracil from ultra-deformable and standard liposomes in-vitro. *Journal of Pharmacy and Pharmacology*, 53(8), 1069–1077. <https://doi.org/10.1211/0022357011776450>
- Maghraby, G. M. E., Williams, A. C., & Barry, B. W. (2010). Skin delivery of oestradiol from deformable and traditional liposomes: Mechanistic studies. *Journal of Pharmacy and Pharmacology*, 51(10), 1123–1134. <https://doi.org/10.1211/0022357991776813>
- Martien, R., Adhyatmika, A., Irianto, I. D., Farida, V., & Sari, D. P. (2012). Perkembangan teknologi nanopartikel sebagai sistem penghantaran obat. *Majalah Farmaseutik*, 8(1), 133–144.
- Mozafari, M. R., Reed, C. J., & Rostron, C. (2007). Prospects of anionic nanolipoplexes in nanotherapy: Transmission electron microscopy and light scattering studies. *Micron (Oxford, England: 1993)*, 38(8), 787–795. <https://doi.org/10.1016/j.micron.2007.06.007>
- Naviglio, D., Dellagrecia, M., Ruffo, F., Andolfi, A., & Gallo, M. (2017). Rapid analysis procedures for triglycerides and fatty acids as pentyl and phenethyl esters for the detection of butter adulteration using chromatographic techniques. *Journal of Food Quality*, 2017, 1–11. <https://doi.org/10.1155/2017/9698107>
- Perkins, E. G. (1995). Composition of soybeans and soybean products. In *Practical handbook of soybean processing and utilization* (pp. 9–28). AOCS Press.
- Puri, A., Loomis, K., Smith, B., Lee, J. H., Yavlovich, A., Heldman, E., & Blumenthal, R. (2009). Lipid-based nanoparticles as pharmaceutical drug carriers: From concepts to clinic. *Critical Reviews™ in Therapeutic Drug Carrier Systems*, 26(6), 523–580.
- Reverchon, E., & Adami, R. (2006). Nanomaterials and supercritical fluids. *The Journal of Supercritical Fluids*, 37(1), 1–22. <https://doi.org/10.1016/j.supflu.2005.08.003>
- Risselada, H. J., & Marrink, S. J. (2009). Curvature effects on lipid packing and dynamics in liposomes revealed by coarse grained molecular dynamics simulations. *Physical Chemistry Chemical Physics: PCCP*, 11(12), 2056–2067. <https://doi.org/10.1039/b818782g>
- Shen, K. C., Kakumanu, S., Beckett, C. D., & Laugharn, J. A. Jr (2015). Use of Adaptive Focused Acoustics™ ultrasound in controlling liposome formation. *Ultrasonics Sonochemistry*, 27, 638–645. <https://doi.org/10.1016/j.ultsonch.2015.04.027>
- Silva, R., Ferreira, H., Little, C., & Cavaco-Paulo, A. (2010). Effect of ultrasound parameters for unilamellar liposome preparation. *Ultrasonics Sonochemistry*, 17(3), 628–632. <https://doi.org/10.1016/j.ultsonch.2009.10.010>
- Singh, M., Manikandan, S., & Kumaraguru, A. K. (2011). Nanoparticles: A new technology with wide applications. *Research Journal of Nanoscience and Nanotechnology*, 1(1), 1–11. <https://doi.org/10.3923/rjnn.2011.1.11>
- Siwko, M. E., de Vries, A. H., Mark, A. E., Kozubek, A., & Marrink, S. J. (2009). Disturb or stabilize? A molecular dynamics study of the effects of resorcinolic lipids on phospholipid bilayers. *Biophysical Journal*, 96(8), 3140–3153. <https://doi.org/10.1016/j.bpj.2009.01.040>
- Trisilowati, & Mallet, D. G. (2012). In silico experimental modeling of cancer treatment. *ISRN Oncology*, 2012, 1–8. <https://doi.org/10.5402/2012/828701>
- Van Hoogevest, P., Prusseit, B., & Wajda, R. (2013). Phospholipids: Natural functional ingredients and actives for cosmetic products. *SOFW-Journal*, 8(9), 14.
- Wang, S., Zhang, J., Jiang, T., Zheng, L., Wang, Z., Zhang, J., & Yu, P. (2011). Protective effect of Coenzyme Q(10) against oxidative damage in human lens epithelial cells by novel ocular drug carriers. *International Journal of Pharmaceutics*, 403(1–2), 219–229. <https://doi.org/10.1016/j.jipharm.2010.10.020>
- Zhao, L., Wei, Y. M., Zhong, X. D., Liang, Y., Zhang, X. M., Li, W., Li, B., Wang, Y., & Yu, Y. (2009). PK and tissue distribution of docetaxel in rabbits after iv administration of liposomal and injectable formulations. *Journal of Pharmaceutical and Biomedical Analysis*, 49(4), 989–996. <https://doi.org/10.1016/j.jpba.2009.01.016>