



Sri Hartati Yuliani <srihartati.yuliani@gmail.com>

[JPSC] Submission Acknowledgement

Enade Perdana Istyastono <libusd@gmail.com>
To: Sri Hartati Yuliani <srihartati.yuliani@gmail.com>

Mon, Jan 13, 2020 at 7:25 PM

The following message is being delivered on behalf of Journal of Pharmaceutical Sciences and Community (J Pharm Sci Community).

Sri Hartati Yuliani:

Thank you for submitting the manuscript, "The Effect of Tempe Extract Gel on Wound Healing in Diabetes Rat: Overview of Tissue Collagen, Wound Closure, Epithelialization and Capillarization" to Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community). With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL:

<https://e-journal.usd.ac.id/index.php/JFSK/author/submission/2357>

Username: syuliani

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Enade Perdana Istyastono
Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community)

Journal of Pharmaceutical Sciences and Community (J Pharm Sci Community)

To send an email to JPSC, please make sure to use the email:

editorial.jpssc@usd.ac.id or visit our site in:

<http://e-journal.usd.ac.id/index.php/JFSK>

(Not just clicking the "reply" button).

Thank you.

The Effect of Tempeh Extract Gel on Wound Healing in Diabetes Rat: Overview of Tissue Collagen, Wound Closure, Epithelialization and Capillarization

Ignatius Adi Kurniawan¹, Rini Dwiastuti¹, Sri Hartati Yuliani^{1*}

1. Magister Pharmacy Program, Sanata Dharma University, Yogyakarta, Indonesia

^{*)} srihartatiyuliani@usd.ac.id

ABSTRACT

The development of wound healing preparations in diabetes conditions needs to be done because the wound healing process in the conditions requires a longer healing process. The aim of this study is to measure the effect of *tempeh* extract gel containing genistein to wound healing in diabetes rats by observing the response parameters of collagen expression, wound closure speed, epithelialization, and capillarization of wound tissue.

The results of testing the wound closure using a base gel treatment, tempeh extract gel 2.5%, 5%, 7.5% and without treatment, indicated that treated group of tempeh extract gel 5% and 7.5% has a significant effect on diabetes wound healing in the test animals. This indication is strengthened by the results of the non-treatment group statistical test with the 5% and 7.5% tempeh extract gel treatment groups significantly difference in influencing the wound closure response, epithelialization and collagen formation in tissue. The results of the statistical treatment group without treatment with the base treatment group without extract gel and tempeh extract gel 2.5% showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. Statistical test results of capillarization responses indicated the results were not significantly different from all treatment groups.

Keywords: gel, tempeh extract, wound healing, diabetes, wound closure

Commented [WU1]: Tempeh or tempe???

Commented [WU2]: Abstract should state about method used.

*Corresponding author: corresponding author's name
Email: corresponding author's email

INTRODUCTION

Diabetes Mellitus (DM) is a metabolic disorder caused by a decrease in the amount of insulin production from pancreatic cells. It is a chronic metabolic disorder in which the pancreas doesn't produce enough insulin or the body cannot use the insulin effectively that is produced. This condition could increase in glucose concentration in the blood (hyperglycemia). Indonesia ranks number 7 for the number of people with DM in the world. The cases of diabetes ulcer were found in 15% of people with DM in the world (Utami and Karim 2014). DM conditions can cause disruption of the wound healing process. Longer wound healing time is influenced by increased matrix metalloproteinase (MMP) activity and decreased fibroblast activity in forming collagen (Sharp and Jane 2011).

Tempeh is a food ingredient made from soy that is very easy to get in Indonesia. The process of boiling and processing soybeans into tempeh can increase levels of isoflavones from soybeans (Utari 2010). Genistein is one of the isoflavone compounds contained in soy that can be extracted and isolated (Agrawal et al. 2015). Genistein is known to have an influence on wound healing through various mechanisms. Administration of preparations containing genistein can accelerate the wound healing process in experimental animals (Emmerson et al. 2010). The treatment of genistein is known to influence the wound healing process in test animals with DM conditions (Tie et al. 2013). Genistein contained in *tempeh* extract needs to be developed into a topical dosage form. So it is necessary to conduct a test to measure of tempeh extract effect in topical preparations on the wound healing process in experimental animals DM. The effect of administration seen from the amount of collagen formed in wound healing.

The development of wound healing preparations in diabetes conditions needs to be done because the wound healing process in diabetes conditions often encountered obstacles. It causes the healing process takes longer. These obstacles include an increase in the potential for infection due to decreased neutrophil migration and prolonged inflammation. In this study, the observation of wound healing activities in diabetic rats was done by giving *tempeh* extract gel. This study aims to look at the effect of the application tempeh extract gel preparations to wound healing in diabetic rats with qualitative and quantitative observation parameters. The measured parameters to be observed were the collagen expression response, wound closure, epithelialization, and capillarization of wound tissue in the presence of tempeh extract gel containing genistein. The preparations used in this study are hydrogels, which are a form of preparation that can be used for wound care. This preparation has several

benefits in closing wounds, maintaining moisture and can be formulated to deliver medicinal ingredients (Gupta et al. 2011).

METHODS

Materials and Instrument

Genistein-standardized Tempeh Extract with levels of 429,055 µg / mL, Ethanol, Water, Buffer Citrate. Streptozotocin (Cayman Chemical_USA), injection ketamine (Generic®), Neutral Buffer Formalin 10%, xylene. Test animals of male Wistar strain rats aged 2 months with a weight range of 150-180g obtained from the Center for Food and Nutrition of Gadjah Mada University Yogyakarta, Masson Thricrome (Bio-Optica_Italia).

Test Animal

The test animals are male Wistar strain rats placed in individual cages maintained at room conditions $25 \pm 2^{\circ}\text{C}$ with a relative air humidity of 75%. The weight control of test animals is carried out by weighing test animals from time to time. Test animal bodies weight between 170-200g.

Induction of animal diabetic try

Experimental animals were induced using Streptozotocin (STZ) with a single dose of 45mg/kg weight. Induction is carried out by injection of i.p STZ solution in citrate buffer (0.1M) pH 4.5. The STZ solution was made 1 hour before induction by dissolving STZ in a cold buffer solution (Etuk 2010).

Measurement of blood glucose levels in experimental animals

Glucose level measurements were performed after Three days of induction using STZ. Blood sampling is done 4 days after STZ induction treatment. The rat blood was taken from the retro-orbital plexus and held in a 1.5ml microtube (let stand for 30 minutes). Blood samples were centrifuged for 15 minutes at 4,000 rpm. The obtained serum was measured glucose levels based on the workings of the GOD FS reagent (Diasys®). Blood sugar levels used in this study were >200 mg/dl (Martsiningsih 2016).

Wound making and measurement of test responses

Experimental animals were anesthetized using ketamine injection at a dose of 50 mg/kg, by administering intramuscular (i.m) (Kintoko 2017). After the animal tried to lose consciousness, the rat's back hair was shaved and wounded using a 5 mm diameter punch biopsy (Júnior et al. 2011).

a. Grouping of Test Animals.

Commented [WU3]: Should mention about ethical clearance approval

Test animals were grouped into five groups. The each group made replication of three test animals, the division of the observation group is as follows:

- DM1 group : Basis Treatment
- DM2 group : Formula 1 (F1) with an extract content of 2.5%
- DM3 group : Formula 2 (F2) with an extract content of 5%
- DM4 group : Formula 3 (F3) with an extract content of 7.5%
- DM5 group : No Treatment

Administration of preparations containing genistein gel preparation was done every 12 hours, in the morning and evening. The amount of weight applied was as much as ± 25 mg in the wound area.

b. Observation of collagen expression and epithelialization

Injured skin samples were fixed using 10% formalin neutral buffer. Then the sample was put in distilled water for 15 minutes and conditioned dehydrated with the addition of ethanol in stages, 70%, 80%, 96%. The Samples were clarified using xylol and printed on paraffin blocks. The slide cutting was done using a microtome with a thickness of 5-6 μ m. Paraffin pieces were stretched in warm water to prevent creases. The preparation was removed using a beaker and dried in 60°C oven. HE staining was carried out on preparations to evaluate closure of dermis tissue and perform the remodeling stage (Febram, 2010). Collagen observations were carried out by Masson Thricrome staining based on the workings of the Masson Thricrome Goldner (Bio-Optica®) kit (Suvik 2012). Furthermore, observations were made using a microscope with a magnification of 10 x 10 times to see the collagen and epithelialization scores formed by calculating the scores as in table I.

Table I. Histological assessment

The degree of collagen formation	The degree of epithelialization	Total new blood vessel formation	Score
Collagen appears very dense / small visual field (density <75%)	High epithelialization / microscope field of view	More than 4 new blood vessels / microscope field of view.	4
Collagen appears solid / small visual field (density> 50% - 75%)	Normal epithelialization / microscope field of view	2-4 new blood vessels / microscope field of view	3
Collagen appears to be less dense / small visual field (density> 25% - 50%)	Low epithelialization / microscope field of view	1-2 new blood vessels / microscope field of view	2
Collagen appears to be very dense / small visual field (density <25%)	There is no epithelialization / microscope field of view	There are no new blood vessels / microscopic field of view	1

c. Observation of wound closure

The observation of wound closure was done by measuring the diameter of the initial wound area and wound area on the day of observation. The diameter was obtained from the average of two wound areas on the right and left-back of the test animal by measuring the diameter of the wound (Kintoko 2017). The percentage measurements of wound closure were calculated using the following formula:

$$\text{Wound closure (\%)} = \frac{\text{Initial wound area} - \text{Wound area on the day}}{\text{Initial wound area}} \times 100\%$$

d. Data analysis

The results of the research data were analyzed using the Shapiro-Wilk test to determine the normality of data distribution. If abnormal data distribution was found, a Kruskal-Wallis analysis was used and a significant difference in results was followed by a Post-Hoc Test.

RESULTS AND DISCUSSION

a. Blood glucose levels in experimental animals

Diabetes in rat test animals is a condition in which the rats have blood glucose levels above 200mg / dl. In this study measurements of serum blood levels of rats were carried out to measure blood glucose levels on days 0, 3, 7 and 14. The test results obtained blood sugar levels of test animals on measurements of 0, 3, 7, 14 were >200mg /dl. This condition was created to ensure that test animals experience diabetes during the wound healing test period to describe the process of wound healing in a diabetes condition.

High glucose levels in diabetes has an effect on the wound healing process. High blood glucose causes the decreasing of neutrophils and macrophages migration, leading to increase potency for infection and prolongation of inflammatory conditions. The expression of TNF- α (Tumor Necrosis Factor-Alpha) will increase during ulcers. The high TNF- α has result in an increase in apoptosis and decrease fibroblast proliferation (Rosyid 2016). Decreasing the number of fibroblasts causes the synthesis of collagen in the wound area will also decrease. The high expression of MMP (Matrix Metalloproteinase) in a diabetes condition has worsen the wound healing process. MMP has cause some ECM (extracellular matrix) degradation, including collagen.

Commented [WU4]: Use reference

b. Response toward wound closure, collagen synthesis, epithelialization, and tissue capillarization

The stages of the wound healing process are grouped into four stages namely, hemostasis, inflammation, proliferation, and remodeling (Velmar et al. 2009). Hemostasis stage occurs in the initial stages after the injury. Inflammation in the wound area is characterized by increased neutrophil and macrophage activity in the wound area. Formation of new blood vessels and fibroblasts occurs at the stage of proliferation. In addition, epithelial tissue begins to form from the edge of the wound. In the remodeling stage, collagen synthesized by fibroblasts will bond together and increasingly crowded. At this stage, the epithelial tissue formed thickens and closes the wound surface area (Baltzis et al. 2014).

The speed of wound closure is indicated by the closure of the wound surface caused by the process of epithelial cell formation on the wound surface. From the results of experiments on the gel base group, there are test subjects who did not experience the wound healing process. This condition is shown by not decreasing the size of the wound diameter. The highest rate of wound closure was achieved in the 7.5% extract gel group.

From the results of the measurement of wound closure speed, the calculation results are obtained as in table II. The results of statistical analysis using the Kruskal-Wallis test ($p < 0.05$), showed a significant difference in the treatment group of 5% extract gel and 7.5% extract gel against the group without treatment. In the 2.5% extraction gel treatment group and the base treatment group, there was no significant difference in the group without treatment.

Genistein is the main isoflavone compound contained in *tempeh* extract. These compounds belong to the phytoestrogen group found in soybean seeds. Estrogen affects the speed of wound closure by increasing mitogenesis and migration of keratinocytes. Estrogen interacts with Estrogen Receptor Beta (ER β) thereby affecting the pattern of differentiation and proliferation activity of keratinocytes (Hornig et al., 2017). Keratinocytes has migrate from the edge of the wound area to the center. Increased proliferation and migration activity of keratinocytes will accelerate the narrowing of the wound area resulting in wound closure.

Table II. Result of wound closure, collagen synthesis, epithelialization, and capillarization.

Group	Wound Closure (%)	Collagen Score	Epithelialization Score	Capillarization Score
DM1	16,23±4,07	1,11±0,33	6,00±2,64	8,33±4,61
DM2	48,99 ± 7,72	1,22±0,44	6,66±0,57	9,66±3,21
DM3	91,89±14,05 *	3,00±0,50*	7,33±2,31*	8,66±3,05
DM4	94,68 ±9,21*	3,67±0,50*	7,33±1,53*	7,66±3,21
DM5	33,82 ±10.30	1,67±1,00	5,00±1,73	5,66±2,51

Information: *significant difference ($p < 0.05$) to the group without treatment. DM1 Group: Basis Treatment, DM2 group: Formula 1 (F1) with an extract content of 2.5%, DM3 group: Formula 2 (F2) with 5% extract content, DM4 Group: Formula 3 (F3) with an extract content of 7.5%, DM5 Group: No Treatment.

Histological observation of tissue samples was made to observe the wound healing process from the process of epithelial tissue formation, blood vessel formation, and collagen. This observation is followed by a scoring assessment of the formation of epithelial cells, blood vessels and collagen.

Wound closure can occur when epithelial cells, formed due to keratinocyte differentiation, migrate over the provisional matrix headed to the middle of the wound. Epithelial cells will meet in the middle area of the wound, so that cell migration will stop. Then, the formation of bacialis membrane will start accordingly. The thickness of the epithelial tissue is done by measuring the image using the help of Image Raster. The appearance of epithelial tissue can be shown by the arrows in figure 1.




Figure1. Microscopic images of epithelialization in each group

The results of statistical analysis using ($p < 0.05$) showed a significant difference in the degree of epithelialization from the formula 5% extract content and 7.5% extract content

of the treatment group. Whereas in the test group with basis application and giving formula 2.5% extract content did not have significant differences in the group without treatment.

Giving *tempeh* extract with levels of 5% extract content and 7.5% extract content has an influence on the degree of epithelialization in the wound healing process. *Tempeh* extract contains genistein which has a mechanism for increasing the degree of epithelialization that depends on keratinocyte cells, fibroblasts, and macrophages. Migration of epithelial cells into the wound surface is one of the keys in the process of re-epithelialization. The speed of cell migration is influenced by the presence of estrogen (Emmerson et al., 2010).

Formation of new blood vessels has a major role in the process of wound healing. The presence of blood vessels in the wound area will increase the wound healing process. Blood vessels contribute to the supply of blood, oxygen, and nutrients to the injured area. Blood vessels will be seen by the presence of red blood cells in the observation of histological preparations as shown in Figure 2. From the results of the study found, there were no significant differences in the formation of blood vessels from each test group. Giving *tempeh* extract gel preparations in the base treatment group, levels of 2.5%, 5%, and 7.5% extract content did not provide significantly different results compared to the group without treatment.




Figure 2. Capillary microscopic depictions of each group.

Collagen is one of the proteins that make up the skin. Collagen fibers have a strong resistance to pressure. The more collagen found in the tissue, the stronger the tissue is against pressure. Collagen observations were carried out by staining using Masson Trichrome Goldner. The results of collagen staining provide a visualization of the green color in the histological preparations as shown in Figure 3. The more green color that appears shows the greater amount of collagen contained in the test tissue.

Commented [WU5]: Use reference

From the results of collagen analysis, the results showed a significant difference in collagen expression in the test group. The 5% formula group and 7.5% extract content formula group had significant differences in the expression of collagen groups without treatment. The test group giving the basis and formula 2.5% extract content did not show any significant difference to the group without treatment.

The content of genistein contained in *tempeh* extract has an influence on collagen synthesis. From previous studies, genistein has an effect on collagen synthesis by binding to estrogen receptors and inhibiting tyrosine kinase (Irrera et al. 2017). The effect of genistein in *tempeh* extract on collagen synthesis is thought to be related to the mechanism of shortening the inflammatory phase so that the synthesis of fibroblasts in skin tissue increases. Fibroblasts are cells that produce collagen, so an increase in the number of fibroblasts in the tissue causes an increase in the amount of collagen.




Figure3. Microscopic picture of collagen in each group

In the condition of diabetes, the formed collagen is degraded due to high MMP in the tissue. The reduced amount of collagen will make the tissue more fragile. Hence, the wound that has been covered more easily open again. With the high amount of collagen in the wound area, it is expected to increase the tissue's resistance to pressure on the skin.

Commented [WU6]: Avoid assumption, use reference

From the results of testing the wound closure using a gel base treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without gel treatment, indicating the treatment group gel 5% and 7.5% *tempeh* extract content treatment can affect diabetes wound healing in test animals. This is based on the results of the statistical test of the untreated group with the treatment group of gel 5% *tempeh* extract content and 7.5% *tempeh* extract content treatment showing significantly different results in influencing the wound closure response, epithelialization and collagen formation. While the statistical results of the untreated group with the base treatment group without extracts and gel 2.5% *tempeh* extract content treatment showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. The results of statistical

tests in influencing capillarization responses indicate the results were not significantly different from all treatment groups.

CONCLUSION

The results of testing the wound closure using a base gel treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without treatment, indicate the treatment group gel 5% *tempeh* extract content and 7.5% *tempeh* extract content treatment has a significant effect on diabetes wound healing in test animals based on the results of statistical tests. The untreated group with the treatment base gel without extract and gel 2.5% *tempeh* extract content treatment showed no statistically significant test results in influencing the wound closure response, epithelialization and collagen formation. In addition, all treatment groups did not show significant capillarization responses.

ACKNOWLEDGEMENT

The authors would like to express their gratitude to the Directorate General of Higher Education (KEMENRISTEK DIKTI) for funding this research through the Master Thesis Research scheme with research contract No: 029 / Penel./ LPPM-USD / IV / 2019, and to Maria Yolanda Intansari and Yosua Hengky Purwanto for assisting the laboratory technicians in this research.

REFERENCES

- Agrawal, S.S., Saxena, S., and Sharma, A., 2015. Phytoestrogen “ Genistein ”: Its Extraction and Isolation from Soybean Seeds. *International Journal of Pharmacognosy and Phytochemical Research*, 7 (6), 1121–1126.
- Anwar, K., 2018. Aktivitas Gel Ekstrak Etanol Umbi Akar Tawas Ut (*Ampelocissus rubiginosa* L .) Terhadap Penyembuhan Luka Insisi Pada Tikus Wistar. *Traditional Medicine Journal*, 23 (April), 30–39.
- Baltzis, D., Eleftheriadou, I., and Veves, A., 2014. Pathogenesis and Treatment of Impaired Wound Healing in Diabetes Mellitus: New Insights. *Advances in Therapy*, 31 (8), 817–836.
- Emmerson, E., Campbell, L., Ashcroft, G.S., and Hardman, M.J., 2010. The phytoestrogen

- genistein promotes wound healing by multiple independent mechanisms. *Molecular and Cellular Endocrinology*, 321 (2), 184–193.
- Etuk, E.U., 2010. Animals models for studying diabetes mellitus Department of Pharmacology , College of Health Sciences , Usmanu Danfodiyo University ., *Agriculture and Biology Journal of North America*, 1 (2), 130–134.
- Febram, B., 2010. Aktivitas sediaan salep ekstrak batang pohon pisang ambon (musa paradisiaca var sapientum) dalam proses persembuhan luka pada mencit (*Mus musculus albinus*), *Majalah Obat Tradisional*, 15(3), pp. 121–137.
- Gupta, B., Agarwal, R., and Alam, M.S., 2011. Hydrogels for wound healing applications. Biomedical Hydrogels: *Biochemistry*, Manufacture and Medical Applications. Woodhead Publishing Limited.
- Horng, H. et al. (2017) ‘Estrogen Effects on Wound Healing’, *International Journal of Molecular science*, pp. 1–14. doi: 10.3390/ijms18112325.
- Irrera, N., Pizzino, G., Anna, R.D., Vaccaro, M., Arcoraci, V., Squadrito, F., Altavilla, D., and Bitto, A., 2017. Dietary Management of Skin Health : The Role of Genistein. *MDPI*, 1–10.
- Júnior, M.C.S., Colchon, P.H., and Portes, A.G., 2011. Histological evaluation on Brazilian green propolis effect in tissue repair of wistar rats cutaneous wounds Histological Evaluation on Brazilian Green Propolis Effect in Tissue Repair of Wistar Rats Cutaneous Wounds, *Lat. Am. J. Pharm.* 30 (2): 383-7 (2011).
- Kintoko, 2017. Effect of Diabetes Condition on Topical Treatment of Binahong Leaf Fraction in Wound Healing Process Pengaruh Kondisi Diabetes pada Pemberian Topikal Fraksi Daun Binahong dalam. *Majalah Obat Tradisional*, 22 (August), 103–110.
- Martsiningsih, M.A., 2016. Gambaran Kadar Glukosa Darah Metode GOD-PAP (Glucose Oksidase – Peroxidase Aminoantypirin) Sampel Serum dan Plasma EDTA (Etylen Diamin Terta Acetat), *Jurnal Teknologi Laboratorium*, Vol.5, No.1, Maret 2016, pp. 45 ~ 48.
- Rosyid, F.N., 2016. Role of TNF- α in Diabetic Ulcer Healing Process. In: *The 2nd International Conference on Science, Technology, and Humanity*. 142–149.
- Sharp, A.N.U. and Jane, C., 2011. Diabetes and its effects on wound healing. *Nursing Standard*, 25 (45), 41–47.
- Suvik, A., 2012. The Use Of Modified Masson’s Trichrome Staining In Collagen Evaluation In Wound Healing Study, *Malaysian Journal of Veterinary Research*, Volume 3 No.1 January 2012, 39-47.

- Tie, L., An, Y., Han, J., Xiao, Y., Xiaokaiti, Y., Fan, S., Liu, S., Chen, A.F., and Li, X., 2013. Genistein accelerates refractory wound healing by suppressing superoxide and FoxO1/iNOS pathway in type 1 diabetes. *Journal of Nutritional Biochemistry*, 24 (1), 88–96.
- Utami, D.T. and Karim, D., 2014. Diabetes mellitus dengan ulkus diabetikum, *Jom Psik* Vol. 1 No. 2 Oktober 2014 1–7.
- Utari, 2010. Tempe Terhadap Kadar Isoflavon, *PGM* 2010, 33(2): 148-153.
- Velmar, T., Bailey, T., and Smrkolj, V., 2009. The wound healing process: An overview of the cellular and molecular mechanisms. *Journal of International Medical Research*, 37 (5), 1528–1542.
- Yuliani, S.H., Gani, M.R., Istyastono, E.P., and Riswanto, F.D.O., 2018. Optimization of genistein and daidzein extraction from a tempeh-fermented product of soybean, *Journal of Pharmacy & Pharmacognosy Research*, 6 (4), 231-241, 2018, 231–241.

The Effect of Tempeh Extract Gel on Wound Healing in Diabetes Rat: Overview of Tissue Collagen, Wound Closure, Epithelialization and Capillarization

Ignatius Adi Kurniawan¹, Rini Dwiastuti¹, Sri Hartati Yuliani^{1*}

1. Magister Pharmacy Program, Sanata Dharma University, Yogyakarta, Indonesia

^{*)} srihartatiyuliani@usd.ac.id

Commented [D01]: Make it consistence: tempe or tempeh

Commented [D02]: diabetic

ABSTRACT

The development of wound healing preparations in diabetes conditions needs to be done because the wound healing process in the conditions requires a longer healing process. The aim of this study is to measure the effect of *tempeh* extract gel containing genistein to wound healing in diabetes rats by observing the response parameters of collagen expression, wound closure speed, epithelialization, and capillarization of wound tissue.

The results of testing the wound closure using a base gel treatment, tempeh extract gel 2.5%, 5%, 7.5% and without treatment, indicated that treated group of tempeh extract gel 5% and 7.5% has a significant effect on diabetes wound healing in the test animals. This indication is strengthened by the results of the non-treatment group statistical test with the 5% and 7.5% tempeh extract gel treatment groups significantly difference in influencing the wound closure response, epithelialization and collagen formation in tissue. The results of the statistical treatment group without treatment with the base treatment group without extract gel and tempeh extract gel 2.5% showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. Statistical test results of capillarization responses indicated the results were not significantly different from all treatment groups.

Keywords: gel, tempeh extract, wound healing, diabetes, wound closure

*Corresponding author: corresponding author's name
Email: corresponding author's email

INTRODUCTION

Diabetes Mellitus (DM) is a metabolic disorder caused by a decrease in the amount of insulin production from pancreatic cells. It is a chronic metabolic disorder in which the pancreas doesn't produce enough insulin or the body cannot use the insulin effectively that is produced. This condition could increase in glucose concentration in the blood (hyperglycemia). Indonesia ranks number 7 for the number of people with DM in the world. The cases of diabetes ulcer were found in 15% of people with DM in the world (Utami and Karim 2014). DM conditions can cause disruption of the wound healing process. Longer wound healing time is influenced by increased matrix metalloproteinase (MMP) activity and decreased fibroblast activity in forming collagen (Sharp and Jane 2011).

Tempeh is a food ingredient made from soy that is very easy to get in Indonesia. The process of boiling and processing soybeans into tempeh can increase levels of isoflavones from soybeans (Utari 2010). Genistein is one of the isoflavone compounds contained in soy that can be extracted and isolated (Agrawal et al. 2015). Genistein is known to have an influence on wound healing through various mechanisms. Administration of preparations containing genistein can accelerate the wound healing process in experimental animals (Emmerson et al. 2010). The treatment of genistein is known to influence the wound healing process in test animals with DM conditions (Tie et al. 2013). Genistein contained in *tempeh* extract needs to be developed into a topical dosage form. So it is necessary to conduct a test to measure of tempeh extract effect in topical preparations on the wound healing process in experimental animals DM. The effect of administration seen from the amount of collagen formed in wound healing.

The development of wound healing preparations in diabetes conditions needs to be done because the wound healing process in diabetes conditions often encountered obstacles. It causes the healing process takes longer. These obstacles include an increase in the potential for infection due to decreased neutrophil migration and prolonged inflammation. In this study, the observation of wound healing activities in diabetic rats was done by giving *tempeh* extract gel. This study aims to look at the effect of the application tempeh extract gel preparations to wound healing in diabetic rats with qualitative and quantitative observation parameters. The measured parameters to be observed were the collagen expression response, wound closure, epithelialization, and capillarization of wound tissue in the presence of tempeh extract gel containing genistein. The preparations used in this study are hydrogels, which are a form of preparation that can be used for wound care. This preparation has several

Commented [D03]: Which one is appropriate: look at or observe?

benefits in closing wounds, maintaining moisture and can be formulated to deliver medicinal ingredients (Gupta et al. 2011).

METHODS

Materials and Instrument

Genistein-standardized Tempeh Extract with levels of 429,055 µg / mL, Ethanol, Water, Buffer Citrate, Streptozotocin (Cayman Chemical_USA), injection ketamine (Generic®), Neutral Buffer Formalin 10%, xylene. Test animals of male Wistar strain rats aged 2 months with a weight range of 150-180g obtained from the Center for Food and Nutrition of Gadjah Mada University Yogyakarta, Masson Thricrome (Bio-Optica_Italia).

Test Animal

The test animals are male Wistar strain rats placed in individual cages maintained at room conditions $25 \pm 2^{\circ} \text{C}$ with a relative air humidity of 75%. The weight control of test animals is carried out by weighing test animals from time to time. Test animal bodies weight between 170-200g.

Induction of animal diabetic try

Experimental animals were induced using Streptozotocin (STZ) with a single dose of 45mg/kg weight. Induction is carried out by injection of i.p STZ solution in citrate buffer (0.1M) pH 4.5. The STZ solution was made 1 hour before induction by dissolving STZ in a cold buffer solution (Etuk 2010).

Measurement of blood glucose levels in experimental animals

Glucose level measurements were performed after Three days of induction using STZ. Blood sampling is done 4 days after STZ induction treatment. The rat blood was taken from the retro-orbital plexus and held in a 1.5ml microtube (let stand for 30 minutes). Blood samples were centrifuged for 15 minutes at 4,000 rpm. The obtained serum was measured glucose levels based on the workings of the GOD FS reagent (Diasys®). Blood sugar levels used in this study were >200 mg/dl (Martsiningsih 2016).

Wound making and measurement of test responses

Experimental animals were anesthetized using ketamine injection at a dose of 50 mg/kg, by administering intramuscular (i.m) (Kintoko 2017). After the animal tried to lose consciousness, the rat's back hair was shaved and wounded using a 5 mm diameter punch biopsy (Júnior et al. 2011).

a. Grouping of Test Animals.

Commented [D04]: Citrate buffer

Commented [D05]: Is there any ethical clearance approval?

Commented [D06]: Which one is appropriate: try or test/trial?

Commented [D07]: Body weight

Test animals were grouped into five groups. The each group made replication of three test animals, the division of the observation group is as follows:

- DM1 group : Basis Treatment
- DM2 group : Formula 1 (F1) with an extract content of 2.5%
- DM3 group : Formula 2 (F2) with an extract content of 5%
- DM4 group : Formula 3 (F3) with an extract content of 7.5%
- DM5 group : No Treatment

Administration of preparations containing genistein gel preparation was done every 12 hours, in the morning and evening. The amount of weight applied was as much as ± 25 mg in the wound area.

b. Observation of collagen expression and epithelialization

Injured skin samples were fixed using 10% formalin neutral buffer. Then the sample was put in distilled water for 15 minutes and conditioned dehydrated with the addition of ethanol in stages, 70%, 80%, 96%. The Samples were clarified using xylol and printed on paraffin blocks. The slide cutting was done using a microtome with a thickness of 5-6 μ m. Paraffin pieces were stretched in warm water to prevent creases. The preparation was removed using a beaker and dried in 60°C oven. HE staining was carried out on preparations to evaluate closure of dermis tissue and perform the remodeling stage (Febram, 2010). Collagen observations were carried out by Masson Thricrome staining based on the workings of the Masson Thricrome Goldner (Bio-Optica®) kit (Suvik 2012). Furthermore, observations were made using a microscope with a magnification of 10 x 10 times to see the collagen and epithelialization scores formed by calculating the scores as in table I.

Table I. Histological assessment

The degree of collagen formation	The degree of epithelialization	Total new blood vessel formation	Score
Collagen appears very dense / small visual field (density <75%)	High epithelialization / microscope field of view	More than 4 new blood vessels / microscope field of view.	4
Collagen appears solid / small visual field (density> 50% - 75%)	Normal epithelialization / microscope field of view	2-4 new blood vessels / microscope field of view	3
Collagen appears to be less dense / small visual field (density> 25% - 50%)	Low epithelialization / microscope field of view	1-2 new blood vessels / microscope field of view	2
Collagen appears to be very dense / small visual field (density <25%)	There is no epithelialization / microscope field of view	There are no new blood vessels / microscopic field of view	1

Commented [DO8]: Stages or levels?

c. Observation of wound closure

The observation of wound closure was done by measuring the diameter of the initial wound area and wound area on the day of observation. The diameter was obtained from the average of two wound areas on the right and left-back of the test animal by measuring the diameter of the wound (Kintoko 2017). The percentage measurements of wound closure were calculated using the following formula:

$$\text{Wound closure (\%)} = \frac{\text{Initial wound area} - \text{Wound area on the day}}{\text{Initial wound area}} \times 100\%$$

Commented [DO9]: State the formula number(1)

d. Data analysis

The results of the research data were analyzed using the Shapiro-Wilk test to determine the normality of data distribution. If abnormal data distribution was found, a Kruskal-Wallis analysis was used and a significant difference in results was followed by a Post-Hoc Test.

Commented [DO10]: Is there any software used in this section?

RESULTS AND DISCUSSION

a. Blood glucose levels in experimental animals

Diabetes in rat test animals is a condition in which the rats have blood glucose levels above 200mg / dl. In this study measurements of serum blood levels of rats were carried out to measure blood glucose levels on days 0, 3, 7 and 14. The test results obtained blood sugar levels of test animals on measurements of 0, 3, 7, 14 were >200mg /dl. This condition was created to ensure that test animals experience diabetes during the wound healing test period to describe the process of wound healing in a diabetes condition.

High glucose levels in diabetes has an effect on the wound healing process. High blood glucose causes the decreasing of neutrophils and macrophages migration, leading to increase potency for infection and prolongation of inflammatory conditions. The expression of TNF- α (Tumor Necrosis Factor-Alpha) will increase during ulcers. The high TNF- α has result in an increase in apoptosis and decrease fibroblast proliferation (Rosyid 2016). Decreasing the number of fibroblasts causes the synthesis of collagen in the wound area will also decrease. The high expression of MMP (Matrix Metalloproteinase) in a diabetes condition has worsen the wound healing process. MMP has cause some ECM (extracellular matrix) degradation, including collagen.

b. Response toward wound closure, collagen synthesis, epithelialization, and tissue capillarization

The stages of the wound healing process are grouped into four stages namely, hemostasis, inflammation, proliferation, and remodeling (Velmar et al. 2009). Hemostasis stage occurs in the initial stages after the injury. Inflammation in the wound area is characterized by increased neutrophil and macrophage activity in the wound area. Formation of new blood vessels and fibroblasts occurs at the stage of proliferation. In addition, epithelial tissue begins to form from the edge of the wound. In the remodeling stage, collagen synthesized by fibroblasts will bond together and increasingly crowded. At this stage, the epithelial tissue formed thickens and closes the wound surface area (Baltzis et al. 2014).

The speed of wound closure is indicated by the closure of the wound surface caused by the process of epithelial cell formation on the wound surface. From the results of experiments on the gel base group, there are test subjects who did not experience the wound healing process. This condition is shown by not decreasing the size of the wound diameter. The highest rate of wound closure was achieved in the 7.5% extract gel group.

From the results of the measurement of wound closure speed, the calculation results are obtained as in table II. The results of statistical analysis using the Kruskal-Wallis test ($p < 0.05$), showed a significant difference in the treatment group of 5% extract gel and 7.5% extract gel against the group without treatment. In the 2.5% extraction gel treatment group and the base treatment group, there was no significant difference in the group without treatment.

Genistein is the main isoflavone compound contained in *tempeh* extract. These compounds belong to the phytoestrogen group found in soybean seeds. Estrogen affects the speed of wound closure by increasing mitogenesis and migration of keratinocytes. Estrogen interacts with Estrogen Receptor Beta (ER β) thereby affecting the pattern of differentiation and proliferation activity of keratinocytes (Hornig et al., 2017). Keratinocytes has migrate from the edge of the wound area to the center. Increased proliferation and migration activity of keratinocytes will accelerate the narrowing of the wound area resulting in wound closure.

Table II. Result of wound closure, collagen synthesis, epithelialization, and capillarization.

Group	Wound Closure (%)	Collagen Score	Epithelialization Score	Capillarization Score
DM1	16,23±4,07	1,11±0,33	6,00±2,64	8,33±4,61
DM2	48,99 ± 7,72	1,22±0,44	6,66±0,57	9,66±3,21
DM3	91,89±14,05 *	3,00±0,50*	7,33±2,31*	8,66±3,05
DM4	94,68 ±9,21*	3,67±0,50*	7,33±1,53*	7,66±3,21
DM5	33,82 ±10.30	1,67±1,00	5,00±1,73	5,66±2,51

Information: *significant difference ($p < 0.05$) to the group without treatment. DM1 Group: Basis Treatment, DM2 group: Formula 1 (F1) with an extract content of 2.5%, DM3 group: Formula 2 (F2) with 5% extract content, DM4 Group: Formula 3 (F3) with an extract content of 7.5%, DM5 Group: No Treatment.

Histological observation of tissue samples was made to observe the wound healing process from the process of epithelial tissue formation, blood vessel formation, and collagen. This observation is followed by a scoring assessment of the formation of epithelial cells, blood vessels and collagen.

Wound closure can occur when epithelial cells, formed due to keratinocyte differentiation, migrate over the provisional matrix headed to the middle of the wound. Epithelial cells will meet in the middle area of the wound, so that cell migration will stop. Then, the formation of bacialis membrane will start accordingly. The thickness of the epithelial tissue is done by measuring the image using the help of Image Raster. The appearance of epithelial tissue can be shown by the arrows in figure 1.




Figure1. Microscopic images of epithelialization in each group

The results of statistical analysis using ($p < 0.05$) showed a significant difference in the degree of epithelialization from the formula 5% extract content and 7.5% extract content

Commented [D011]: Check the decimal places, in English they should be separated using (.) not (,)

Commented [D012]: This figure should be presented clearly. The quality of labels in the bottom should be improved. It is recommended to authors to label the figure with A, B, C, and D followed by the brief caption to make it informative.

of the treatment group. Whereas in the test group with basis application and giving formula 2.5% extract content did not have significant differences in the group without treatment.

Giving *tempeh* extract with levels of 5% extract content and 7.5% extract content has an influence on the degree of epithelialization in the wound healing process. *Tempeh* extract contains genistein which has a mechanism for increasing the degree of epithelialization that depends on keratinocyte cells, fibroblasts, and macrophages. Migration of epithelial cells into the wound surface is one of the keys in the process of re-epithelialization. The speed of cell migration is influenced by the presence of estrogen (Emmerson et al., 2010).

Formation of new blood vessels has a major role in the process of wound healing. The presence of blood vessels in the wound area will increase the wound healing process. Blood vessels contribute to the supply of blood, oxygen, and nutrients to the injured area. Blood vessels will be seen by the presence of red blood cells in the observation of histological preparations as shown in Figure 2. From the results of the study found, there were no significant differences in the formation of blood vessels from each test group. Giving *tempeh* extract gel preparations in the base treatment group, levels of 2.5%, 5%, and 7.5% extract content did not provide significantly different results compared to the group without treatment.




Figure 2. Capillary microscopic depictions of each group.

Collagen is one of the proteins that make up the skin. Collagen fibers have a strong resistance to pressure. The more collagen found in the tissue, the stronger the tissue is against pressure. Collagen observations were carried out by staining using Masson Trichrome Goldner. The results of collagen staining provide a visualization of the green color in the histological preparations as shown in Figure 3. The more green color that appears shows the greater amount of collagen contained in the test tissue.

Commented [D013]: Similar comment with Figure 1. This figure should be presented clearly. The quality of labels in the bottom should be improved. It is recommended to authors to label the figure with A, B, C, and D followed by the brief caption to make it informative.

From the results of collagen analysis, the results showed a significant difference in collagen expression in the test group. The 5% formula group and 7.5% extract content formula group had significant differences in the expression of collagen groups without treatment. The test group giving the basis and formula 2.5% extract content did not show any significant difference to the group without treatment.

The content of genistein contained in *tempeh* extract has an influence on collagen synthesis. From previous studies, genistein has an effect on collagen synthesis by binding to estrogen receptors and inhibiting tyrosine kinase (Irrera et al. 2017). The effect of genistein in *tempeh* extract on collagen synthesis is thought to be related to the mechanism of shortening the inflammatory phase so that the synthesis of fibroblasts in skin tissue increases. Fibroblasts are cells that produce collagen, so an increase in the number of fibroblasts in the tissue causes an increase in the amount of collagen.




Figure3. Microscopic picture of collagen in each group

In the condition of diabetes, the formed collagen is degraded due to high MMP in the tissue. The reduced amount of collagen will make the tissue more fragile. Hence, the wound that has been covered more easily open again. With the high amount of collagen in the wound area, it is expected to increase the tissue's resistance to pressure on the skin.

From the results of testing the wound closure using a gel base treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without gel treatment, indicating the treatment group gel 5% and 7.5% *tempeh* extract content treatment can affect diabetes wound healing in test animals. This is based on the results of the statistical test of the untreated group with the treatment group of gel 5% *tempeh* extract content and 7.5% *tempeh* extract content treatment showing significantly different results in influencing the wound closure response, epithelialization and collagen formation. While the statistical results of the untreated group with the base treatment group without extracts and gel 2.5% *tempeh* extract content treatment showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. The results of statistical

Commented [D014]: This figure should be presented clearly. The quality of labels in the bottom should be improved. It is recommended to authors to label the figure with A, B, C, and D followed by the brief caption to make it informative.

tests in influencing capillarization responses indicate the results were not significantly different from all treatment groups.

CONCLUSION

The results of testing the wound closure using a base gel treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without treatment, indicate the treatment group gel 5% *tempeh* extract content and 7.5% *tempeh* extract content treatment has a significant effect on diabetes wound healing in test animals based on the results of statistical tests. The untreated group with the treatment base gel without extract and gel 2.5% *tempeh* extract content treatment showed no statistically significant test results in influencing the wound closure response, epithelialization and collagen formation. In addition, all treatment groups did not show significant capillarization responses.

Commented [D015]: In this section, it is better not to include the numerical results value. You can summarize the final results with brief statement as effective as possible.

ACKNOWLEDGEMENT

The authors would like to express their gratitude to the Directorate General of Higher Education (KEMENRISTEK DIKTI) for funding this research through the Master Thesis Research scheme with research contract No: 029 / Penel./ LPPM-USD / IV / 2019, and to Maria Yolanda Intansari and Yosua Hengky Purwanto for assisting the laboratory technicians in this research.

REFERENCES

- Agrawal, S.S., Saxena, S., and Sharma, A., 2015. Phytoestrogen “ Genistein ”: Its Extraction and Isolation from Soybean Seeds. *International Journal of Pharmacognosy and Phytochemical Research*, 7 (6), 1121–1126.
- Anwar, K., 2018. Aktivitas Gel Ekstrak Etanol Umbi Akar Tawas Ut (*Ampelocissus rubiginosa* L .) Terhadap Penyembuhan Luka Insisi Pada Tikus Wistar. *Traditional Medicine Journal*, 23 (April), 30–39.
- Baltzis, D., Eleftheriadou, I., and Veves, A., 2014. Pathogenesis and Treatment of Impaired Wound Healing in Diabetes Mellitus: New Insights. *Advances in Therapy*, 31 (8), 817–836.
- Emmerson, E., Campbell, L., Ashcroft, G.S., and Hardman, M.J., 2010. The phytoestrogen

- genistein promotes wound healing by multiple independent mechanisms. *Molecular and Cellular Endocrinology*, 321 (2), 184–193.
- Etuk, E.U., 2010. Animals models for studying diabetes mellitus Department of Pharmacology , College of Health Sciences , Usmanu Danfodiyo University ., *Agriculture and Biology Journal of North America*, 1 (2), 130–134.
- Febraam, B., 2010. Aktivitas sediaan salep ekstrak batang pohon pisang ambon (musa paradisiaca var sapientum) dalam proses persembuhan luka pada mencit (*Mus musculus albinus*), *Majalah Obat Tradisional*, 15(3), pp. 121–137.
- Gupta, B., Agarwal, R., and Alam, M.S., 2011. Hydrogels for wound healing applications. Biomedical Hydrogels: *Biochemistry*, Manufacture and Medical Applications. Woodhead Publishing Limited.
- Horng, H. et al. (2017) ‘Estrogen Effects on Wound Healing’, *International Journal of Molecular science*, pp. 1–14. doi: 10.3390/ijms18112325.
- Irrera, N., Pizzino, G., Anna, R.D., Vaccaro, M., Arcoraci, V., Squadrito, F., Altavilla, D., and Bitto, A., 2017. Dietary Management of Skin Health : The Role of Genistein. *MDPI*, 1–10.
- Júnior, M.C.S., Colchon, P.H., and Portes, A.G., 2011. Histological evaluation on Brazilian green propolis effect in tissue repair of wistar rats cutaneous wounds Histological Evaluation on Brazilian Green Propolis Effect in Tissue Repair of Wistar Rats Cutaneous Wounds, *Lat. Am. J. Pharm.* 30 (2): 383-7 (2011).
- Kintoko, 2017. Effect of Diabetes Condition on Topical Treatment of Binahong Leaf Fraction in Wound Healing Process Pengaruh Kondisi Diabetes pada Pemberian Topikal Fraksi Daun Binahong dalam. *Majalah Obat Tradisional*, 22 (August), 103–110.
- Martsiningsih, M.A., 2016. Gambaran Kadar Glukosa Darah Metode GOD-PAP (Glucose Oksidase – Peroxidase Aminoantypirin) Sampel Serum dan Plasma EDTA (Etylen Diamin Terta Acetat), *Jurnal Teknologi Laboratorium*, Vol.5, No.1, Maret 2016, pp. 45 ~ 48.
- Rosyid, F.N., 2016. Role of TNF- α in Diabetic Ulcer Healing Process. In: *The 2nd International Conference on Science, Technology, and Humanity*. 142–149.
- Sharp, A.N.U. and Jane, C., 2011. Diabetes and its effects on wound healing. *Nursing Standard*, 25 (45), 41–47.
- Suvik, A., 2012. The Use Of Modified Masson’s Trichrome Staining In Collagen Evaluation In Wound Healing Study, *Malaysian Journal of Veterinary Research*, Volume 3 No.1 January 2012, 39-47.

- Tie, L., An, Y., Han, J., Xiao, Y., Xiaokaiti, Y., Fan, S., Liu, S., Chen, A.F., and Li, X., 2013. Genistein accelerates refractory wound healing by suppressing superoxide and FoxO1/iNOS pathway in type 1 diabetes. *Journal of Nutritional Biochemistry*, 24 (1), 88–96.
- Utami, D.T. and Karim, D., 2014. Diabetes mellitus dengan ulkus diabetikum, *Jom Psik* Vol. 1 No. 2 Oktober 2014 1–7.
- Utari, 2010. Tempe Terhadap Kadar Isoflavon, *PGM* 2010, 33(2): 148-153.
- Velnar, T., Bailey, T., and Smrkolj, V., 2009. The wound healing process: An overview of the cellular and molecular mechanisms. *Journal of International Medical Research*, 37 (5), 1528–1542.
- Yuliani, S.H., Gani, M.R., Istyastono, E.P., and Riswanto, F.D.O., 2018. Optimization of genistein and daidzein extraction from a tempeh-fermented product of soybean, *Journal of Pharmacy & Pharmacognosy Research*, 6 (4), 231-241, 2018, 231–241.

The Effect of Tempe Extract Gel on Wound Healing in Diabetes Rat: Overview of Tissue Collagen, Wound Closure, Epithelialization and Capillarization

Ignatius Adi Kurniawan¹, Rini Dwiastuti¹, Sri Hartati Yuliani^{1*}

1. Magister Program of Pharmacy Sanata Dharma University, Yogyakarta, Indonesia

*) srihartatiyuliani@usd.ac.id

ABSTRACT

The development of wound healing preparations in diabetes conditions needs to be done because the wound healing process in the respective conditions requires a longer time. The aim of this study is to measure the effect of *tempeh* extract gel containing genistein on wound healing in diabetes rats by observing the response parameters of collagen expression, wound closure speed, epithelialization, and capillarization of wound tissue.

The results of testing wound closure using a base gel treatment, *tempeh* extract gel 2.5%, 5%, 7.5% and without treatment, indicated that the treated group of *tempeh* extract gel 5% and 7.5% has a significant effect on diabetes wound healing in the tested animals. This indication is strengthened by the results of the non-treatment group statistical test with the 5% and 7.5% *tempeh* extract gel treatment groups significant difference in influencing the wound closure response, epithelialization and collagen formation in tissue. The results of the statistical in group without treatment with the base treatment group without extract gel and *tempeh* extract gel 2.5% showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. Statistical test results on capillarization responses indicated that the results were not significantly different from all treatment groups.

Keywords: gel, *tempeh* extract, wound healing, diabetes, wound closure

INTRODUCTION

Diabetes Mellitus (DM) is a metabolic disorder caused by a decrease in the amount of insulin production from pancreatic cells. It is a chronic metabolic disorder in which the pancreas does not produce enough insulin or the body cannot use the available insulin effectively. This condition could increase glucose concentration in the blood (hyperglycemia). Indonesia ranks number 7 for the number of people with DM in the world. The cases of diabetes ulcer were found in 15% of people with DM in the world (Utami and Karim, 2014). DM conditions can cause disruption on wound healing process. Longer wound healing time is influenced by increasing matrix metalloproteinase (MMP) activity and decreasing fibroblast activity in forming collagen (Sharp and Jane, 2011).

Tempeh is food made of soy that is very easy to get in Indonesia. The process of boiling and processing soybeans into *tempeh* can increase levels of isoflavones from soybeans (Utari, 2010). Genistein is one of the isoflavone compounds contained in soybeans that can be extracted and isolated (Agrawal et al., 2015). Genistein is known to have an influence on wound healing through various mechanisms. Administration of gel containing genistein can accelerate the wound healing process in experimental animals (Emmerson et al., 2010). The treatment of genistein is known to influence the wound healing process in test animals with DM conditions (Tie et al., 2013). Genistein contained in *tempeh* extract needs to be developed into a topical dosage form. So, it is necessary to conduct a test to measure *tempeh* extract effect in topical preparations on wound healing process in experimental animals with DM. The effect of administration is seen from the amount of collagen formed in wound healing.

The development of wound healing preparations in diabetes conditions needs to be done because wound healing process in diabetes conditions often encounters obstacles. It causes healing process to take a longer period. These obstacles include an increasing potential for infection due to decreasing neutrophil migration and prolonged inflammation. In this study, the observation of wound healing activities in diabetic rats was done by giving *tempeh* extract gel. This study aims to look at the effect of *tempeh* extract gel preparations to wound healing in diabetic rats with qualitative and quantitative observation parameters. The measured parameters to be observed were the collagen expression response, wound closure, epithelialization, and capillarization of wound tissue in the presence of *tempeh* extract gel containing genistein. The preparations used in this study were hydrogels, a form of

preparation that can be used for wound care. This preparation has several benefits in closing wounds, maintaining moisture, and delivering medicinal ingredients (Gupta et al., 2011).

METHODS

Materials and Instrument

The materials included enistein-standardized *tempeh* extract with levels of 429,055 µg / mL, ethanol, water, buffer citrate, streptozotocin (Cayman Chemical_USA), injection ketamine (Generic®), neutral buffer formalin 10%, and xylene. Test animals of male Wistar strain rats aged 2 months with a weight range of 150-180g was obtained from the Center for Food and Nutrition of Gadjah Mada University Yogyakarta, masson thricrome (Bio-Optica_Italia).

Test Animal

The tested animals were male Wistar strain rats placed in individual cages maintained at room conditions of $25 \pm 2^{\circ} \text{C}$ with a relative air humidity of 75%. The weight control of the tested animals was carried out by weighing them from time to time. Tested animals' bodies weighed between 170-200g.

Induction of animal diabetic try

Experimental animals were induced with Streptozotocin (STZ) with a single dose of 45mg/kg. Induction was carried out by injection of i.p STZ solution in citrate buffer (0.1M) pH 4.5. The STZ solution was made an hour before the induction by dissolving STZ in a cold buffer solution (Etuk, 2010).

Measurement of blood glucose levels in experimental animals

Glucose level measurements were performed after three days of induction with STZ. Blood sampling was done 4 days after STZ induction treatment. The rat blood was taken from the retro-orbital plexus, held in a 1.5ml microtube, and let standing for 30 minutes. Blood samples were centrifuged for 15 minutes at 4,000 rpm. The obtained serum was measured on its glucose levels based on the workings of the GOD FS reagent (Diasys®). Blood sugar levels used in this study were >200 mg/dl (Martsiningsih 2016).

Wound making and measurement of test responses

Experimental animals were anesthetized with ketamine injection at a dose of 50 mg/kg by administering intramuscular (i.m) (Kintoko, 2017). After the animal lost consciousness, the rat's back hair was shaved and wounded using a 5 mm diameter punch biopsy (Júnior et al., 2011).

a. Grouping of Tested Animals.

Tested animals were grouped into five. Each group made replication of three tested animals. The division of the observation group is as follows:

- DM1 group : Basis Treatment
- DM2 group : Formula 1 (F1) with extract content of 2.5%
- DM3 group : Formula 2 (F2) with extract content of 5%
- DM4 group : Formula 3 (F3) with extract content of 7.5%
- DM5 group : No Treatment

Administration of preparations containing genistein gel was done every 12 hours in the morning and evening. The amount applied was as much as ± 25 mg in the wound area.

b. Observation of collagen expression and epithelialization

Injured skin samples were fixed using 10% of formalin neutral buffer. Then, the sample was put in distilled water for 15 minutes and conditioned dehydrated with the addition of ethanol in stages; 70%, 80%, and 96%. The samples were clarified using xylol and printed on paraffin blocks. The slide cutting was done using a microtome with a thickness of 5-6 μm . Paraffin pieces were stretched in warm water to prevent creases. The preparation was removed using a beaker and dried in 60°C oven. HE staining was carried out on preparations to evaluate closure of dermis tissue and perform the remodeling stage (Febram, 2010). Collagen observations were carried out by Masson Thricrome staining based on the workings of the Masson Thricrome Goldner (Bio-Optica®) kit (Suvik, 2012). Furthermore, observations were made using a microscope with a magnification of 10 x 10 times to see the collagen and epithelialization scores. The result is shown in table I.

Table I. Histological assessment

Parameter and description	Score
The degree of collagen formation	
a. Collagen appears very dense / small visual field (density <75%)	4
b. Collagen appears solid / small visual field (density > 50% - 75%)	3
c. Collagen appears to be less dense / small visual field (density > 25% -50%)	2
d. Collagen appears to be very dense / small visual field (density <25%)	1
The degree of epithelialization	
a. High epithelialization / microscope field of view	4
b. Normal epithelialization / microscope field of view	3
c. Low epithelialization / microscope field of view	2
d. There is no epithelialization / microscope field of view	1
Total new blood vessel formation	
a. More than 4 new blood vessels / microscope field of view	4
b. 2-4 new blood vessels / microscope field of view	3
c. 1-2 new blood vessels / microscope field of view	2
d. There are no new blood vessels / microscopic field of view	1

c. Observation of wound closure

The observation of wound closure was done by measuring the diameter of the initial wound area and wound area on the day of observation. The diameter was obtained from the average of two wound areas on the right and left-back of the tested animal (Kintoko, 2017). The percentage measurements of wound closure were calculated using the following formula:

$$\text{Wound closure (\%)} = \frac{\text{initial wound area} - \text{wound area on the day}}{\text{initial wound area}} \times 100\%$$

d. Data analysis

The results of the research data were analyzed using the Shapiro-Wilk test to determine the normality of data distribution. If abnormal data distribution was found, a Kruskal-Wallis analysis was used and a significant difference in results was followed by a Post-Hoc Test.

RESULTS AND DISCUSSION

a. Blood glucose levels in experimental animals

Diabetes in rat tested animals is a condition in which the rats have blood glucose levels above 200mg / dl. In this study measurements of serum blood levels of rats were carried out to measure blood glucose levels on day 0, 3, 7 and 14. The blood sugar level tests on the animals within measurements of 0, 3, 7, 14 were >200mg /dl. This condition was created to ensure that the tested animals experienced diabetes during wound healing test period.

High glucose level in diabetes has an effect on the wound healing process. High blood glucose causes decreasing neutrophils and macrophages migration leading to increasing potency for infection and prolongation of inflammatory conditions. The expression of TNF- α (Tumor Necrosis Factor-Alpha) will increase during ulcers. The high TNF- α increases apoptosis and decreases fibroblast proliferation (Rosyid, 2016). Decreasing number of fibroblasts causes the synthesis of collagen in the wound area will also decrease. The high expression of MMP (Matrix Metalloproteinase) in a diabetes condition has worsen the wound

healing process. MMP has cause some ECM (extracellular matrix) degradation including collagen.

b. Response toward wound closure, collagen synthesis, epithelialization, and tissue capillarization

The stages of the wound healing process are grouped into four stages namely, hemostasis, inflammation, proliferation, and remodeling (Velmar et al., 2009). Hemostasis stage occurs in the initial stage after the injury. Inflammation in the wound area is characterized by the increase of neutrophil and macrophage activity in the wound area. Formation of new blood vessels and fibroblasts occurs at the stage of proliferation. In addition, epithelial tissue begins to form from the edge of the wound. In the remodeling stage, collagen synthesized by fibroblasts will bond together and crowd increasingly. At this stage, the epithelial tissue forms thickens and closes the wound surface area (Baltzis et al., 2014).

The speed of wound closure is indicated by the closure of the wound surface caused by the process of epithelial cell formation on the wound surface. From the results of experiments on the gel base group, there are test subjects who did not experience the wound healing process. This condition is shown by the the size of the wound diameter which did not decrease. The highest rate of wound closure was achieved in the 7.5% extract gel group.

From the results of the measurement of wound closure speed, the calculation results are obtained as in table II. The results of statistical analysis using the Kruskal-Wallis test ($p < 0.05$) shows a significant difference in the treatment group of 5% extract gel, and 7.5% extract gel against the group without treatment. In 2.5% extraction gel treatment group and base treatment group, there was no significant difference in the group without treatment.

Genistein is the main isoflavone compound contained in *tempeh* extract. The compound belongs to the phytoestrogen group found in soybean seeds. Estrogen affects the speed of wound closure by increasing mitogenesis and migration of keratinocytes. Estrogen interacts with Estrogen Receptor Beta (ER β) thereby affecting the pattern of differentiation and proliferation activity of keratinocytes (Horng et al., 2017). Keratinocytes has migrated from the edge of the wound area to the center. Increased proliferation and migration activity of keratinocytes will accelerate the narrowing of the wound area resulting in wound closure.


Table II. Result of wound closure, collagen synthesis, epithelialization, and capillarization.

Group	Wound Closure (%) ±SD	Collagen Score ± SD	Epithelialization Score ± SD	Capillarization Score ± SD
DM1	16,23±4,07	1,11±0,33	6,00±2,64	8,33±4,61
DM2	48,99 ± 7,72	1,22±0,44	6,66±0,57	9,66±3,21
DM3	91,89±14,05 *	3,00±0,50*	7,33±2,31*	8,66±3,05
DM4	94,68 ±9,21*	3,67±0,50*	7,33±1,53*	7,66±3,21
DM5	33,82 ±10.30	1,67±1,00	5,00±1,73	5,66±2,51

Information: *significant difference (p <0.05) to the DM5 group.

Histological observation of tissue samples was made to observe the wound healing process from the process of epithelial tissue formation, blood vessel formation, and collagen. This observation is followed by a scoring assessment of the formation of epithelial cells, blood vessels and collagen.

Wound closure can occur when epithelial cells, formed due to keratinocyte differentiation, migrate over the provisional matrix headed to the middle of the wound. Epithelial cells will meet in the middle area of the wound, so that cell migration will stop. Then, the formation of bacialis membrane will start accordingly. The thickness of the epithelial tissue is done by measuring the image using the help of image raster. The appearance of epithelial tissue can be shown by the arrows in figure 1.

**Figure1.** Microscopic images of epithelialization in each group

The results of statistical analysis using (p <0.05) showed a significant difference in the degree of epithelialization from the formula 5% extract content and 7.5% extract content of the treatment group. Whereas in the test group with basis application, giving formula 2.5% extract content did not have significant differences in the group without treatment.

Giving *tempeh* extract with levels of 5% and 7.5% extract contents has an influence on the degree of epithelialization in the wound healing process. *Tempeh* extract contains

genistein which has a mechanism for increasing the degree of epithelialization that depends on keratinocyte cells, fibroblasts, and macrophages. Migration of epithelial cells into the wound surface is one of the keys in the process of re-epithelialization. The speed of cell migration is influenced by the presence of estrogen (Emmerson et al., 2010).

Formation of new blood vessels has a major role in the process of wound healing. The presence of blood vessels in the wound area will increase the wound healing process. Blood vessels contribute to the supply of blood, oxygen, and nutrients to the injured area. Blood vessels will be seen by the presence of red blood cells in the observation of histological preparations as shown in Figure 2. From the results of the study, there were no significant differences in the formation of blood vessels from each test group. Giving *tempeh* extract gel preparations in the base treatment group, levels of 2.5%, 5%, and 7.5% extract content did not provide significantly different results compared to the group without treatment.




Figure 2. Capillary microscopic depictions of each group.

Collagen is one of the proteins that make up the skin. Collagen fibers have a strong resistance to pressure. The more collagen found in the tissue, the stronger the tissue is against pressure. Collagen observations were carried out by staining using Masson Trichrome Goldner. The results of collagen staining provide a visualization of the green color in the histological preparations as shown in Figure 3. The more green color that appears shows the greater amount of collagen contained in the test tissue.

From the results of collagen analysis, it showed a significant difference in collagen expression in the test group. The 5% formula group and 7.5% extract content formula group had significant differences in the expression of collagen groups without treatment. The test group giving the basis and formula 2.5% extract content did not show any significant difference to the group without treatment.

The content of genistein contained in *tempeh* extract has an influence on collagen synthesis. From previous studies, genistein has an effect on collagen synthesis by binding to estrogen receptors and inhibiting tyrosine kinase (Irrera et al., 2017). The effect of genistein

in *tempeh* extract on collagen synthesis is thought to be related to the mechanism of shortening the inflammatory phase so that the synthesis of fibroblasts in skin tissue increases. Fibroblasts are cells that produce collagen, so an increase in the number of fibroblasts in the tissue causes an increase in the amount of collagen.




Figure 3. Microscopic picture of collagen in each group

In the condition of diabetes, the formed collagen is degraded due to high MMP in the tissue. The reduced amount of collagen will make the tissue more fragile. Hence, the wound that has been covered becomes more easily open again. With the high amount of collagen in the wound area, it is expected to increase the tissue's resistance to pressure on the skin.

From the results of testing, the wound closure using a gel base treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without gel treatment, indicating the treatment group gel 5% and 7.5% *tempeh* extract content treatment can affect diabetes wound healing in test animals. This is based on the results of the statistical test of the untreated group with the treatment group of gel 5% *tempeh* extract content and 7.5% *tempeh* extract content treatment showing significantly different results in influencing the wound closure response, epithelialization and collagen formation. While the statistical results of the untreated group with the base treatment group without extracts and gel 2.5% *tempeh* extract content treatment showed no significant difference in influencing the wound closure response, epithelialization and collagen formation. The results of statistical tests in influencing capillarization responses indicate the results were not significantly different from all treatment groups.

CONCLUSION

The results of testing the wound closure using a base gel treatment, gel 2.5% *tempeh* extract content, gel 5% *tempeh* extract content, gel 7.5% *tempeh* extract content and without treatment, indicate that the treatment group gel 5% *tempeh* extract content and 7.5% *tempeh*

extract content treatment have a significant effect on diabetes wound healing in test animals based on the results of statistical tests. The untreated group with the treatment base gel without extract and gel 2.5% *tempeh* extract content treatment showed no statistically significant test results in influencing the wound closure response, epithelialization and collagen formation. In addition, all treatment groups did not show significant capillarization responses.

ACKNOWLEDGEMENT

The authors would like to express their gratitude to the Directorate General of Higher Education (KEMENRISTEK DIKTI) for funding this research through the Master Thesis Research scheme with research contract No: 029 / Penel./ LPPM-USD / IV / 2019, and to Maria Yolanda Intansari and Yosua Hengky Purwanto for assisting the laboratory technicians in this research.

REFERENCES

- Agrawal, S.S., Saxena, S., and Sharma, A., 2015. Phytoestrogen “ Genistein ”: Its Extraction and Isolation from Soybean Seeds. *International Journal of Pharmacognosy and Phytochemical Research*, 7 (6), 1121–1126.
- Anwar, K., 2018. Aktivitas Gel Ekstrak Etanol Umbi Akar Tawas Ut (*Ampelocissus rubiginosa* L .) Terhadap Penyembuhan Luka Insisi Pada Tikus Wistar. *Traditional Medicine Journal*, 23 (April), 30–39.
- Baltzis, D., Eleftheriadou, I., and Veves, A., 2014. Pathogenesis and Treatment of Impaired Wound Healing in Diabetes Mellitus: New Insights. *Advances in Therapy*, 31 (8), 817–836.
- Emmerson, E., Campbell, L., Ashcroft, G.S., and Hardman, M.J., 2010. The phytoestrogen genistein promotes wound healing by multiple independent mechanisms. *Molecular and Cellular Endocrinology*, 321 (2), 184–193.
- Etuk, E.U., 2010. Animals models for studying diabetes mellitus Department of Pharmacology , College of Health Sciences , Usmanu Danfodiyo University ., *Agriculture and Biology Journal of North America*, 1 (2), 130–134.
- Febam, B., 2010. Aktivitas sediaan salep ekstrak batang pohon pisang ambon (*musa paradisiaca* var *sapientum*) dalam proses persembuhan luka pada mencit (*Mus musculus albinus*), *Majalah Obat Tradisional*, 15(3), pp. 121–137.

- Gupta, B., Agarwal, R., and Alam, M.S., 2011. Hydrogels for wound healing applications. Biomedical Hydrogels: *Biochemistry, Manufacture and Medical Applications*. Woodhead Publishing Limited.
- Horng, H. et al. (2017) 'Estrogen Effects on Wound Healing', *International Journal of Molecular science*, pp. 1–14. doi: 10.3390/ijms18112325.
- Irrera, N., Pizzino, G., Anna, R.D., Vaccaro, M., Arcoraci, V., Squadrito, F., Altavilla, D., and Bitto, A., 2017. Dietary Management of Skin Health : The Role of Genistein. *MDPI*, 1–10.
- Júnior, M.C.S., Colchon, P.H., and Portes, A.G., 2011. Histological evaluation on Brazilian green propolis effect in tissue repair of wistar rats cutaneous wounds Histological Evaluation on Brazilian Green Propolis Effect in Tissue Repair of Wistar Rats Cutaneous Wounds, *Lat. Am. J. Pharm.* 30 (2): 383-7 (2011).
- Kintoko, 2017. Effect of Diabetes Condition on Topical Treatment of Binahong Leaf Fraction in Wound Healing Process Pengaruh Kondisi Diabetes pada Pemberian Topikal Fraksi Daun Binahong dalam. *Majalah Obat Tradisional*, 22 (August), 103–110.
- Martsiningsih, M.A., 2016. Gambaran Kadar Glukosa Darah Metode GOD-PAP (Glucose Oksidase – Peroxidase Aminoantipirin) Sampel Serum dan Plasma EDTA (Ethylen Diamin Terta Acetat), *Jurnal Teknologi Laboratorium*, Vol.5, No.1, Maret 2016, pp. 45 ~ 48.
- Rosyid, F.N., 2016. Role of TNF- α in Diabetic Ulcer Healing Process. *In: The 2nd International Conference on Science, Technology, and Humanity*. 142–149.
- Sharp, A.N.U. and Jane, C., 2011. Diabetes and its effects on wound healing. *Nursing Standard*, 25 (45), 41–47.
- Suvik, A., 2012. The Use Of Modified Masson's Trichrome Staining In Collagen Evaluation In Wound Healing Study, *Malaysian Journal of Veterinary Research*, Volume 3 No.1 January 2012, 39-47.
- Tie, L., An, Y., Han, J., Xiao, Y., Xiaokaiti, Y., Fan, S., Liu, S., Chen, A.F., and Li, X., 2013. Genistein accelerates refractory wound healing by suppressing superoxide and FoxO1/iNOS pathway in type 1 diabetes. *Journal of Nutritional Biochemistry*, 24 (1), 88–96.
- Utami, D.T. and Karim, D., 2014. Diabetes mellitus dengan ulkus diabetikum, *Jom Psik* Vol. 1 No. 2 Oktober 2014 1–7.
- Utari, 2010. Tempe Terhadap Kadar Isoflavon, *PGM* 2010, 33(2): 148-153.
- Velmar, T., Bailey, T., and Smrkolj, V., 2009. The wound healing process: An overview of

the cellular and molecular mechanisms. *Journal of International Medical Research*, 37 (5), 1528–1542.

Yuliani, S.H., Gani, M.R., Istyastono, E.P., and Riswanto, F.D.O., 2018. Optimization of genistein and daidzein extraction from a *tempeh*-fermented product of soybean, *Journal of Pharmacy & Pharmacognosy Research*, 6 (4), 231-241, 2018, 231–241.



Sri Hartati Yuliani <srihartati.yuliani@gmail.com>

Proofing manuscript

J. Pharm. Sci. Community <editorial.jpssc@usd.ac.id>
To: Sri Hartati Yuliani <srihartati.yuliani@gmail.com>
Cc: "J. Pharm. Sci. Community" <editorial.jpssc@usd.ac.id>

Fri, Apr 24, 2020 at 8:24 AM

Sri Hartati Yuliani:

Your submission "The Effect of Tempe Extract Gel on Wound Healing in Diabetes Rat: Overview of Tissue Collagen, Wound Closure, Epithelialization and Capillarization" for Jurnal Farmasi Sains dan Komunitas (Journal of Pharmaceutical Sciences and Community) has been through the first step of copyediting, and is available for you to review by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and click on the File that appears in Step 1.
3. Open the downloaded submission.
4. Review the text, including copyediting proposals and Author Queries.
5. Make any copyediting changes that would further improve the text.
6. When completed, upload the file in Step 2.
7. Click on METADATA to check indexing information for completeness and accuracy.
8. Send the COMPLETE email to the editor and copyeditor.

Submission URL: <https://e-journal.usd.ac.id/index.php/JFSK/author/submissionEditing/2357>

Username: syuliani

This is the last opportunity to make substantial copyediting changes to the submission. The proofreading stage, that follows the preparation of the galleys, is restricted to correcting typographical and layout errors.

If you are unable to undertake this work at this time or have any questions, please contact me. Thank you for your contribution to this journal.

Michael Raharja Gani
Faculty of Pharmacy, Universitas Sanata Dharma
Phone 085643207613
michaelvaynardgani@gmail.com