



Digital Receipt

This receipt acknowledges that Turnitin received your paper. Below you will find the receipt information regarding your submission.

The first page of your submissions is displayed below.

Submission author: Bernadeta Wuri Harini
Assignment title: Quick Submit
Submission title: Development of Curcuminoid Content Measurement Equip...
File name: ment_Equipment_Device_in_Turmeric_Curcuma_domestica_...
File size: 202.37K
Page count: 7
Word count: 1,984
Character count: 10,197
Submission date: 11-May-2021 03:22PM (UTC+0700)
Submission ID: 1583447693

Development of Curcuminoid Content Measurement Equipment Device in Turmeric (*Curcuma domestica*) Rhizomes

Bernadeta Wuri Harini¹⁾, Rini Dwiastuti²⁾, Lucia Widi Wijayanti³⁾
¹⁾Electrical Engineering Department, Sanata Dharma University,
wuribernard@usd.ac.id

²⁾Pharmacy Department, Sanata Dharma University, rindwiastuti83@gmail.com
³⁾Biology Department, Sanata Dharma University, lwijayanti@yahoo.com

Abstract

Turmeric (Curcuma domestica/kunyit, Indo) is containing curcumin a medical compound for some diseases. The most important part of this plant is it's rhizomes.

It has developed the measurement equipment device with an aim to empower Turmeric (*Curcuma domestica*) farmers in the turmeric production centre area, Wonosobo, Wonogiri, Inogiri, Magelang, and Klaten. This research showed that 33,3% farmers in that area do the sorting according to size and 23,8% according to water content. There is not any practice to sort according to curcumin content. It is caused by the lack of knowledge and practical equipment among the farmers. This is the reason why this research is carried out.

The measurement equipment device is using the principal of spectrophotometer. From the scanning test of standard curcumin using visible-spectrophotometer, we get 405 nm for the maximum wavelength of absorbance. The wavelength is in violet range. Therefore, the device will be developed using light source of violet LED (Light Emitting Diode) and violet Laser. The result of this research showed that both of the light source can be used to measure the absorbance of curcuminoid content of turmeric.

Keywords: Curcuminoid, Curcuma Domestica, LED, Laser, violet

1. Introduction

Turmeric (Curcuma domestica/kunyit, Indo) is containing curcumin a medical compound for some diseases. The most important part of this plant is it's rhizomes. Curcumin is a phenolic compound derived from the rhizome isolation of plant turmeric (*Curcuma domestica* Rhizome) containing desmetoksikurkumin, curcumin and bisdesmetoksikurkumin, which is often referred to as the curcuminoids. The main content of curcuminoids is curcumin, a yellow orange. Directed toward the development of medicinal plants to meet domestic industry, pharmaceuticals, cosmetics, household industry, carrying herbs and exports. There is a lot of data and literature that indicates that the compound curcumin in turmeric (*Curcuma domestica*) has great potential in the pharmacological activity of anti-inflammatory, anti immunodeficiency, anti-virus (bird flu virus), anti-bacterial, anti-fungal, anti-oxidant, anti-carcinogenic and anti-infective (Joe *et al.*, 2004; Chattopadhyay *et al.*, 2004).

It has developed the measurement equipment device with an aim to empower Turmeric (*Curcuma domestica*) farmers in the turmeric production centre area, Wonosobo, Wonogiri, Inogiri, Magelang, and Karanganyar. The measurement equipment device is using the principal of spectrophotometer. Some researchers have examined the absorption of curcumin by using spectrophotometer. According to Harada, T., maximum wavelength of absorption of curcumin is 427-329 nm (Harada, T., 2011), while according to Jagannathan, R., maximum wavelength for absorption of curcumin in methanol is about 420nm (Jagannathan, R., 2011). From the scanning test of standard curcumin using visible spectrophotometer, we get 422 nm for the maximum wavelength of absorbance. From all this research can be seen that the

53

Development of Curcuminoid Content Measurement Equipment Device in Turmeric (*Curcuma domestica*) Rhizomes

by Harini Bernadeta Wuri

Submission date: 11-May-2021 03:22PM (UTC+0700)

Submission ID: 1583447693

File name: ment_Equipment_Device_in_Turmeric_Curcuma_domestica_Rhizomes.pdf (202.37K)

Word count: 1984

Character count: 10197

2

Development of Curcuminoid Content Measurement Equipment Device in Turmeric (*Curcuma domestica*) Rhizomes

6

Bernadeta Wuri Harini¹⁾, Rini Dwiastuti²⁾, Lucia Wiwid Wijayanti³⁾

¹⁾Electrical Engineering Department, Sanata Dharma University,
wuribernard@usd.ac.id

²⁾Pharmacy Department, Sanata Dharma University, rinidwiastuti83@gmail.com

³⁾Biology Department, Sanata Dharma University, l.wijayanti@yahoo.com

Abstract

Turmeric (Curcuma domestica/kunyit, Indo) is containing curcumin a medical compound for some diseases. The most important part of this plant is it's rhizomes.

It has developed the measurement equipment device with an aim to empower Turmeric (*Curcuma domestica*) farmers in the turmeric production centre area, Wonosobo, Wonogiri, Imogiri, Magelang, and Kulonprogo. This research showed that 33.3% farmers in that area do the sorting according to size and 23.8% according to water content. There is not any practice to sort according to curcumin content. It is caused by the lack of knowledge and practical equipment among the farmers. This is the reason why this research is carried out.

The measurement equipment device is using the principal of spectrophotometer. From the scanning test of standard curcumin using visible spectrophotometer, we get 405 nm for the maximum wavelength of absorbance. The wavelength is in violet range. Therefore, the device will be developed using light source of violet LED (Light Emitting Diode) and violet Laser. The result of this research showed that both of the light source can be used to measure the absorbance of curcuminoid content of turmeric.

Keywords: Curcuminoid, Curcuma Domestica, LED, Laser, violet

1. Introduction

Turmeric (Curcuma domestica/kunyit, Indo) is containing curcumin a medical compound for some diseases. The most important part of this plant is it's rhizomes. Curcumin is a phenolic compound derived from the rhizome isolation of plant turmeric (*Curcuma domestica* Rhizome) containing desmetoksikurkumin, curcumin and bisdesmetoksikurkumin, which is often referred to as the curcuminoids. The main content of curcuminoids is curcumin, a yellow orange. Directed toward the development of medicinal plants to meet domestic industry, pharmaceuticals, cosmetics, household industry, carrying herbs and exports. There is a lot of data and literature that indicates that the compound curcumin in turmeric (*Curcuma domestica*) has great potential in the pharmacological activity of anti-inflammatory, anti immunodeficiency, anti-virus (bird flu virus), anti-bacterial, anti-fungal, anti-oxidant, anti-carcinogenic and anti-infective (Joe *et al.*, 2004; Chattopadhyay *et al.*, 2004).

It has developed the measurement equipment device with an aim to empower Turmeric (*Curcuma domestica*) farmers in the turmeric production centre area, Wonosobo, Wonogiri, Imogiri, Magelang, and Karanganyar. The measurement equipment device is using the principal of spectrophotometer. Some researchers have examined the absorption of curcumin by using spectrophotometer. According to Harada, T., maximum wavelength of absorption of curcumin is 427-329 nm (Harada, T., 2011), while according to Jagannathan, R., maximum wavelength for absorption of curcumin in methanol is about 420nm (Jagannathan, R., 2011). From the scanning test of standard curcumin using visible spectrophotometer, we get 422 nm for the maximum wavelength of absorbance. From all this research can be seen that the

wavelength of absorption of curcumin present in purple light spectrum range, which is between 380-435 nm (Jones, A.Z., accessed in Oct 11, 2012). Therefore, the device will be developed using light source of violet LED (Light Emitting Diode) and violet Laser.

2. Research Method

The stages of this study are as follows:

- a. Survey of farmers capabilities in sorting turmeric and collect samples of 5 turmeric turmeric producing region is in Central Java and Yogyakarta, from the village of Jumantono Karanganyar, Wonogiri district, village Pandansari Magelang, Wonosobo, and village Kunden Imogiri
- b. Preparation of standard curve of curcumin liquor to obtain a standard curve equation


$$y = bx + a$$

(1)


where y = absorbance

x = levels of substances

How to create a solution for standard curve is as follows:



- c. Preparation of samples from various places planting turmeric in the following ways



- d. Prototyping curcuminoids measurement device using purple LED and laser light sources.
The values of b and that is obtained on standard curve equation used to calculate the levels of curcumin in turmeric samples. At first photodetector receives the voltage of the beam of purple light when no molecular absorber between the light source and a photodetector. Then a solution containing curcumin is placed between the light source

and a photodetector. The light through the absorbing molecule to be received by a photodetector. Voltage difference between the before and after it is placed a solution of curcumin is the absorbance value (y). With the obtained values of a, b and y from these measurements, you will get the levels of curcumin in solution (x) according to standard curve equation above by the formula

$$x = \frac{y - a}{b}$$

(2)

Figure 1 shows the block diagram of this system.



Figure 1. Block diagram

3. Results and Analysis

A. Survey Result

Figure 2 shows the turmeric agricultural area in ¹Karanganyar, Wonogiri, Wonosobo, Magelang and Imogiri.



Figure 2. The turmeric agricultural area (a) Karanganyar (b) Wonogiri (c) Wonosobo (d) Magelang (e) Imogiri

Table 1. The results of the survey capability of farmers

Statement	%Yes	%No
Do sorting the harvest that is sold	30.43	69.57
Sorting criterion is based on the size of	30.43	69.57
Sorting criterion is based on the level of drought / water content	21.74	78.26
Sorting criterion is based on the amount of curcumin content	4.54	95.46
Farmers have to test the levels of curcumin in turmeric crop	0	100

This research showed that 30.43% farmers in that area do the sorting according to size and 21.79% according to water content. Farmers who have been sorting turmeric by curcumin levels only 4.54%. It is caused by the lack of knowledge and practical equipment among the farmers. This is the reason why this research is carried out.

B. Prototyping Curcuminoids Measurement Device



Figure 3. Curcuminoids measurement device prototype

C. Preparation of Standard Curve and Samples of Turmeric

Figure 4 shows the cuvette, the solution for the standard curve and samples from different growing areas in Central Java and Yogyakarta. Table 2 shows the results of measurements of standard curves on tools that have been made and compared with results obtained with a standard spectrophotometer. These results are then illustrated in Figure 5.



Figure 4. The sample

Table 2. The results of absorbance measurements for standard curve

NO	Curcumin solution	LED voltage (Volt)	Laser voltage (Volt)	Spectrophotometer
1	1 ppm	0.16	0.06	0.156
2	2 ppm	0.35	0.08	0.306
3	3 ppm	0.78	0.1	0.461
4	4 ppm	1.73	0.11	0.612
5	5 ppm	2.5	0.12	0.764




Figure 5. Standart curve of curcumin solution

From Figure 5 it appears that the results obtained from measurements using a standard spectrophotometer produced a linear graph with the level of linearity equal to 1, while for LED to laser at 0.940 and at 0.969. It appears that the results of measurements with a laser light source produced more linear than the LED. This is because the laser beam is more focused than LED beam. The linearity level could be improved by making the cover along the beam of light, so the measurement is not affected by the light from outside sources of light. From the chart above also obtained standard curve equation as follows:

f. Spectrophotometer

$$y = 0.152x + 0.003 \quad (3)$$

g. LED

$$y = 0.606x - 0.714 \quad (4)$$

h. Laser

$$y = 0.015x + 0.049 \quad (5)$$

In table 3 are shown the results of measurements of the sample absorbance planting from different regions by using three device. From these results is then used to measure the levels of curcumin in turmeric according to equation 2 with a and b as in equation 3-5, which are shown in Table 4.

Table 3. The results of turmeric absorbance measurements

NO	Region	LED voltage (Volt)	Laser voltage (Volt)	Spectrophotometer
1	Karanganyar	2.51	0.11	0.609
2	Magelang	2.38	0.13	0.848
3	Imogiri	2.32	0.13	0.812
4	Wonosobo	1.35	0.118	0.532
5	Wonogiri	2.15	0.12	0.784

Table 4. The results of the curcuminoid levels calculation of turmeric sample (x)

NO	Region	Curcuminoid Levels (x)			Error (%)	
		LED	Laser	Spectrophotometer	LED	Laser
1	Karanganyar	5.32	4.07	3.99	33.44	2.00
2	Magelang	5.11	5.40	5.56	8.16	2.86
3	Imogiri	5.01	5.40	5.32	5.93	1.46
4	Wonosobo	3.41	4.60	3.48	2.14	32.17
5	Wonogiri	4.73	4.73	5.14	8.02	7.88

From Table 4 it appears that the level of curcuminoid (x) using LED and laser differ from those generated by a standard spectrophotometer with error between 1.46 to 33.44%. The smallest error resulting from the measurement of turmeric in Imogiri area using a laser light source. Greatest error resulting from the measurement of turmeric in Karanganyar area using LED light source. From the above calculations it appears that the measuring instrument with a laser light source produces measurements with an average error smaller (9.28%) than LED source (11.54%). It shows the performance of the laser light source is better than LED. The order from largest to smallest levels as follows::

- LED: Karanganyar - Magelang - Imogiri - Wonogiri - Wonosobo
- Laser: Magelang - Imogiri - Wonogiri - Wonosobo - Karanganyar
- Spektrofotometer standar: Magelang - Imogiri - Wonogiri - Karanganyar - Wonosobo

The difference in results is due to the measurement is still in an open condition, so there is likely to be affected by external light. In addition, samples were measured after two hours the samples were extracted and in the outdoor environment without closed. This allows the change in turmeric samples. Hygiene factors cuvette and cuvette position when put in place will influence the measurement cuvette. Therefore prototype of curcuminoid levels device is still needed improvement, in good repair and measurement tools.

4. Conclusion.

The result of this research showed that both of the light source can be used to measure the absorbance of curcuminoid content of turmeric. Error that is generated by laser source (9.28%) is smaller than that is produced by LED source (11.54%).

5. References.

Chattopadhyay, I., Biswas, K., Bandyopadhyay, U. and Banerjee, R.K., 2004., *Tumeric and Curcumin : Biological actions and medicinal applications*. *Current Science*. 87 (1) : 44 - 53.

Harada, T., 2011, Cooperative Binding and Stabilization of the Medicinal Pigment Curcumin by Diamide Linked γ -Cyclodextrin Dimers: A Spectroscopic Characterization, *The Journal of Physical Chemistry*, pages 1268–1274
http://www.laserchemistry.adelaide.edu.au/Kee/Papers_pdf/jp1096025_si_001.pdf, was accessed on Oct 11 2012

Joe, B.; M. Vijaykumar and B.R. Lokesh, 2004, *Biological properties of curcumin-cellular and molecular mechanisms of action*. *Critical Review in Food Science and Nutrition* 44 (2) : 97 - 112.

Jones, A.Z., *The Visible Light Spectrum*,
<http://physics.about.com/od/lightoptics/a/vislightspec.htm>, was accessed on Oct 11 2012

Ramya Jagannathan, 2011, Solubilising Curcumin, Synthesizing Gold Nanoparticles And Their Anti-Oxidant Property ,
http://shodhganga.inflibnet.ac.in/bitstream/10603/2446/16/16_chapter5.pdf , was accessed on Oct 11 2012 420 nm

6. Acknowledgements

Thanks is given to: Lian Chrismatsy, Oktovianus Ferryandi and Marito Dos Santos who helped research.

Development of Curcuminoid Content Measurement Equipment Device in Turmeric (*Curcuma domestica*) Rhizomes

ORIGINALITY REPORT

10%	10%	6%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	repository.akprind.ac.id	4%
	Internet Source	
2	repository.usd.ac.id	3%
	Internet Source	
3	journal.sttnas.ac.id	1%
	Internet Source	
4	www.intechopen.com	1%
	Internet Source	
5	Vadivambal, Rajagopal, and Digvir Jayas. "Imaging in the visible spectrum", Bio-Imaging, 2015.	1%
	Publication	
6	www.scribd.com	<1%
	Internet Source	

Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	Off		